

## **On Hermitian interpolation of first order data with locally generated $C^1$ -splines over triangular meshes**

By László L. Stachó

**Abstract.** Given a system of triangles in the plane  $\mathbb{R}^2$  along with given data of function and gradient values at the vertices, we describe the general pattern of local linear methods involving only four smooth standard shape functions which results in a spline function fitting the given value and gradient data value with  $C^1$ -coupling along the edges of the triangles. We characterize their invariance properties with relevance for the construction of interpolation surfaces over triangularizations of scanned 3D data. The numerically simplest procedures among them leaving invariant all polynomials of 2-variables with degree 0 resp 1 involve only polynomials of 5-th resp. 6-th degree, but the characterizations give rise to a huge variety of procedures with non-polynomial shape functions.

### **1. Introduction**

Recently [5] we published a  $C^1$ -spline interpolation method over 2-dimensional triangular meshes with polynomials of 5-th degree with low operational costs: using first order data (value and gradient) at the mesh vertices for input. The result relies on the following basic tool: given a non-degenerate triangle with vertices  $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3 \in \mathbb{R}^2$  along with three values  $f_1, f_2, f_3 \in \mathbb{R}$  respectively three linear functionals  $A_1, A_2, A_3 \in \mathcal{L}(\mathbb{R}^2, \mathbb{R})$  and three vectors  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3 \in \mathbb{R}^2$  such that  $\mathbf{u}_k \not\parallel \mathbf{p}_i - \mathbf{p}_j$ , we can construct a polynomial  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$  of 5-th degree of the

---

*Mathematics Subject Classification:* 65D07, 41A15, 65D15.

*Key words and phrases:*  $C^1$ -spline, triangular mesh, 2D Hermite interpolation, reduced side derivatives (RSD), shape function, range shift property, affinity invariance.

form  $F = F_0 - H$  where

$$\begin{aligned} F_0(\mathbf{x}) &= \sum_{i=1}^3 \left[ \Phi(\lambda_i(\mathbf{x}))f_i + \Theta(\lambda_i(\mathbf{x}))A_i(\mathbf{x} - \mathbf{p}_i) \right], \\ H(\mathbf{x}) &= 30\lambda_1(\mathbf{x})^2\lambda_2(\mathbf{x})^2\lambda_3(\mathbf{x})^2 \sum_{k=1}^3 \lambda_k(\mathbf{x})^{-1} \frac{M_k \mathbf{u}_k}{G_k \mathbf{u}_k} \end{aligned} \quad (1.1)$$

in terms of the *barycentric weights*  $\lambda_1, \lambda_2, \lambda_3 : \mathbb{R}^2 \rightarrow \mathbb{R}$  of the triangle  $\mathbf{T} = \text{Conv}\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$  associated with the vertices and the *shape functions*

$$\Phi(t) = t^3(10 - 15t + 6t^2), \quad \Theta(t) = t^3(4 - 3t) \quad (1.2)$$

for a Hermite interpolation  $F_0$  on  $\mathbf{T}$  with the data  $f_i, A_i$  ( $i = 1, 2, 3$ ). The correction term  $H$  is defined by means of the linear functionals  $\mathbb{R}^2 \rightarrow \mathbb{R}$

$$\begin{aligned} G_k \mathbf{u} &= \lambda'_k \mathbf{u} = \left. \frac{d}{dt} \right|_{t=0} \lambda_k(\mathbf{x} + t\mathbf{u}) = \lambda_k(\mathbf{x} + \mathbf{u}) - \lambda_k(\mathbf{x}), \\ M_k \mathbf{u} &= \sum_{\{i,j\}=\{1,2,3\}\setminus\{k\}} [G_i \mathbf{u}] (30f_i + 12A_i(\mathbf{p}_j - \mathbf{p}_i)). \end{aligned}$$

It is worth to notice that the function  $F_0$  in (1.1) restricted to the edge  $[\mathbf{p}_i, \mathbf{p}_k]$  depends only on the terms  $f_i, f_j, A_i, A_j$  and the impact of the correction by adding  $H$  results in the *reduced side derivatives (RSD)*

$$F'(\mathbf{x}_t)\mathbf{u}_k = \Theta(t)A_i\mathbf{u}_k + \Theta(1-t)A_j\mathbf{u}_k \quad \text{for } \mathbf{x}_t = t\mathbf{p}_i + (1-t)\mathbf{p}_j.$$

As a consequence, if we choose any family  $\{\mathbf{u}_{\mathbf{E}} : \mathbf{E} \text{ being a mesh edge}\}$  of vectors such that  $\mathbf{u}_{\{\mathbf{E}\}} \not\parallel \mathbf{E}$ , by applying the construction (1.1) with the associated data over every triangle of the mesh, we obtain a  $\mathcal{C}^1$ -smooth (continuously differentiable) function on the union of the mesh triangles.

The shape functions  $\Phi, \Theta$  and also the RSD method described above appeared in [5] without heuristic introduction. Actually they arose from our earlier study [6] with somewhat restrictive postulates on the possible polynomial  $\mathcal{C}^1$ -interpolations over triangular meshes based on computer algebraic analysis of techniques developed in [8],[4],[3],[2].

Our goal here is to characterize all RSD methods with shape functions  $\Psi_0, \Psi_1 \in \mathcal{C}^1([0, 1])$  instead of (1.2). This requires a completely different approach as that in [6] relying heavily upon polynomial identities. From our main result

Theorem 3.5 it turns out that there is a very general pattern behind (1.1) in the form

$$\begin{aligned}
F_0 &= \sum_{k=1}^3 \left[ \Psi_0(\lambda_k) f_k + \Psi_1(\lambda_k) A_k(\mathbf{x} - \mathbf{p}_k) \right], \\
H &= \sum_{\{\ell, m, n\}=\{1,2,3\}} \left\{ f_\ell \frac{G_\ell \mathbf{u}_n}{G_n \mathbf{u}_n} \chi_0(\lambda_\ell, \lambda_m, \lambda_n) + \right. \\
&\quad \left. + A_\ell(\mathbf{p}_m - \mathbf{p}_\ell) \frac{G_\ell \mathbf{u}_n}{G_n \mathbf{u}_n} \chi_1(\lambda_\ell, \lambda_m, \lambda_n) \right\}.
\end{aligned} \tag{1.3}$$

If  $\Psi_0, \Psi_1$  fulfill the minimal necessary condition (2.1) for giving rise to a Hermite interpolation of the type  $F_0$  in (1.1) of first order then we can find plenty of suitable functions  $\chi_0, \chi_1 \in C_0^1(\mathbb{R}_+^3)$ , namely those satisfying only the not too restrictive conditions (3.6), (3.7), in order for  $H$  being an RSD correction for  $F_0$ . This degree of freedom enables various canonical constructions for the modifiers  $\chi_0, \chi_1$  in terms of suitable shape functions  $\Psi_0, \Psi_1$  to ensure specific geometric properties of the interpolation operator  $\mathfrak{F} : \{(\mathbf{p}_k, f_k, A_k) : k = 1, 2, 3\} \mapsto [F_0 + H$  by (1.3)].

We complete the paper with the investigation of two essential properties of the interpolation  $\mathfrak{F}$  having obvious importance in applications used to construct smooth surfaces in the form of the graph of a function  $\mathbb{R}^2 \rightarrow \mathbb{R}$  passing through a finite family of points in  $\mathbb{R}^3$  (obtained as vertices of a triangularization from scanned data):

(i) The *range shift property*  $\mathfrak{F}\{(\mathbf{p}_k, 1, 0) : k = 1, 2, 3\} \equiv 1$ , that is all constant functions remain invariant when interpolated with  $\mathfrak{F}$ . In this case the graph of  $\mathfrak{F}\{(\mathbf{p}_k, f_k + c, f'(\mathbf{p}_k)) : k = 1, 2, 3\}$  is just a shifted form of the graph of  $\mathfrak{F}\{(\mathbf{p}_k, f_k, (\mathbf{p}_k)) : k = 1, 2, 3\}$  for any function  $f \in C^1(\mathbb{R}^2)$ .

(ii) *Affinity invariance*:  $\mathfrak{F}\{(\mathbf{p}_k, A\mathbf{x} + b, A) : k = 1, 2, 3\} \equiv A\mathbf{x} + b$ . That is all affine (constant+linear) functions remain invariant when interpolated with  $\mathfrak{F}$ . In particular if the graph of a function  $\text{Conv}\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\} \rightarrow \mathbb{R}$  is a 3D plane triangle then the function obtained with interpolation by  $\mathfrak{F}$  has the same graph.

In Section 4 we give a parametric classification of the procedures with range shift property. Concerning affinity invariance, we have no complete results yet: in Section 5 we provide a several necessary and sufficient algebraic conditions. As a by no means obvious fact, it turns out that the procedure (1.1) with the polynomials (1.2) has range shift property but fails to be affinity invariant. We also provide an affinity invariant procedure with polynomials of 6-th degree: namely that of the form (1.3) with  $\Psi_0 = \Psi_1 = \Phi$  and  $\chi_0(t_1, t_2, t_3) = 30t_1^2t_2^2t_3$ ,  $\chi_1(t_1, t_2, t_3) = 30t_1^2t_2^3t_3 + 15t_1^2t_2^2t_3^2$ .

## 2. Preliminaries

For standard terminology,  $\mathbb{R}^N$  resp.  $\mathbb{R}_+^N$  stand for the set of all real resp. non-negative  $N$ -tuples  $[\xi_1, \dots, \xi_N]$ . Actually, we shall only be interested in the cases of dimensions  $N = 1, 2, 3$ .

Given a subset  $\Omega$  of  $\mathbb{R}^N$ , we write  $\mathcal{C}(\Omega)$  for the family of all continuous functions with domain  $\Omega$ . If the interior  $\Omega^\circ$  of  $\Omega$  is dense in  $\Omega$ , we define  $\mathcal{C}^1(\Omega)$  as the set consisting of all functions  $f \in \mathcal{C}(\Omega)$  with continuous partial derivatives  $D_1f, \dots, D_Nf$  on  $\Omega^\circ$  admitting a continuous extension to  $\Omega$  with the value denoted also by  $D_kf(\mathbf{x})$  at the points  $\mathbf{x} \in \Omega \setminus \Omega^\circ$ . As a folklore consequence of Whitney's extension theorem [7], if  $\Omega$  is closed in  $\mathbb{R}^N$  then every function  $f \in \mathcal{C}^1(\Omega)$  can be regarded as the restriction of a continuously differentiable function defined on  $\mathbb{R}^N$ .

For any function  $f \in \mathcal{C}^1(\Omega)$ , we write  $f'$  for its *Fréchet derivative* defined at any point  $\mathbf{x} \in \Omega$  as the linear functional

$$f'(\mathbf{x})\mathbf{u} = \sum_{k=1}^N D_kf(\mathbf{x})v_k \quad (\mathbf{u} = [v_1, \dots, v_N]).$$

In particular  $f'(\mathbf{x})\mathbf{u} = \lim_{t \rightarrow 0} t^{-1} [f(\mathbf{x} + t\mathbf{u}) - f\mathbf{u}]$  is the familiar directional derivative at the points  $\mathbf{x} \in \Omega^\circ$ . If  $\Omega$  is a closed set, we define  $\mathcal{C}_0^k(\Omega) = \{f \in \mathcal{C}^k(\Omega) : f(\mathbf{x}) = 0 \ (\mathbf{x} \in \Omega \setminus \Omega^\circ)\}$  ( $k = 0, 1$ ).

Given any subset  $\mathbf{P} \subset \mathbb{R}^2$ , we write  $\text{Conv}(\mathbf{P})$  for its convex hull. If  $\mathbf{P} = \{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$  and the triangle  $\mathbf{T} = \text{Conv}(\mathbf{P})$  is non-degenerate, the *barycentric weights* (weight functions associated to the vertices) [1] are the functions

$$\lambda_i = \lambda_{\mathbf{p}_i}^{\mathbf{T}} : \mathbf{x} \mapsto \frac{\det[\mathbf{x} - \mathbf{p}_j, \mathbf{x} - \mathbf{p}_k]}{\det[\mathbf{p}_i - \mathbf{p}_j, \mathbf{p}_i - \mathbf{p}_k]} \quad \text{with } (i, j, k) \in S_3$$

where

$$S_3 = \{(1, 2, 3), (2, 1, 3), (2, 3, 1), (3, 2, 1), (3, 1, 2), (1, 3, 2)\}$$

denotes the set of all permutations of the indices 1, 2, 3. For the sake of brevity, in the sequel we omit the background parameters in most formulas (like writing  $\lambda_i = \lambda_{\mathbf{p}_i}^{\mathbf{T}}$  above) without danger of confusion. It is well-known that the mapping  $\mathbf{x} \mapsto [\lambda_1(\mathbf{x}), \lambda_2(\mathbf{x}), \lambda_3(\mathbf{x})]$  is a homeomorphism between  $\mathbf{T}$  and the *3D-unit simplex*

$$\Delta_3 := \{[t_1, t_2, t_3] \in \mathbb{R}_+^3 : t_1 + t_2 + t_3 = 1\},$$

moreover  $[\lambda_1(\mathbf{x}), \lambda_2(\mathbf{x}), \lambda_3(\mathbf{x})]$  is the unique triple of non-negative numbers such that  $\mathbf{x} = \sum_k \lambda_k(\mathbf{x})\mathbf{p}_k$  and  $\sum_k \lambda_k(\mathbf{x}) = 1$ . Furthermore  $\lambda_1, \lambda_2, \lambda_3 : \mathbb{R}^2 \rightarrow \mathbb{R}$  are affine (linear+constant) functions with the necessarily constant Fréchet derivatives

$$G_i\mathbf{u} := \lambda'_i(\mathbf{x}) = \left. \frac{d}{dt} \right|_{t=0} \lambda_i(\mathbf{x} + t\mathbf{u}) \quad \text{independently of } \mathbf{x}.$$

Given any pair  $\Psi = [\Psi_0, \Psi_1]$  with  $\Psi_0, \Psi_1 \in \mathcal{C}^1([0, 1])$  such that

$$\Psi_0(0) = \Psi'_0(0) = \Psi_1(0) = \Psi'_1(0) = \Psi'_0(1) = 0, \quad \Psi_0(1) = \Psi_1(1) = 1 \quad (2.1)$$

we introduce the *basic triangular interpolation of first order with the shape functions*  $\Psi_0, \Psi_1$  as the operator

$$\mathfrak{F}_0^\Psi : \{(\mathbf{p}_k, f_k, A_k)\}_{k=1}^3 \mapsto \sum_{k=1}^3 \left\{ \Psi_0(\lambda_k) f_k + \Psi_1(\lambda_k) A_k(\mathbf{x} - \mathbf{p}_k) \right\} \quad (2.2)$$

defined for all *first order function germs*  $\{(\mathbf{p}_k, f_k, A_k) : k = 1, 2, 3\}$  with  $\mathbf{T} = \text{Conv}\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\} \subset \mathbb{R}^2$  being a non-degenerate triangle,  $f_k \in \mathbb{R}$  and  $A_k \in \mathcal{L}(\mathbb{R}^2, \mathbb{R})$  in terms of the weights  $\lambda_k = \lambda_k^\mathbf{T}$ . By definition, the domain of the function  $F_0 = \mathfrak{F}_0^\Psi \{(\mathbf{p}_k, f_k, A_k)\}_{k=1}^3$  is only the triangle  $\mathbf{T}$  and  $F_0 \in \mathcal{C}^1(\mathbf{T})$ . By straightforward calculation, for its Fréchet derivative we have

$$F'_0 = \sum_{k=1}^3 \left\{ [\Psi'_0(\lambda_k) f_k + [\Psi'_1(\lambda_k) A_k(\mathbf{x} - \mathbf{p}_k)] G_k + \Psi_1(\lambda_k) A_k] \right\}. \quad (2.3)$$

*Remark. 2.4.* (a) Conditions (2.1) do not restrict the value of  $\Psi'_1(1)$ .

(b) In view of (2.3), it is not hard to see that (2.1) is sufficient and necessary for  $F_0$  to satisfy the relations

$$F_0(\mathbf{p}_i) = f_i, \quad F'_0(\mathbf{p}_i) = A_i \quad (i = 1, 2, 3) \quad (2.5)$$

under any choice of the first order function germs  $\{(\mathbf{p}_k, f_k, A_k)\}_{k=1}^3$ .

(c) We can express the terms  $A_k(\mathbf{x} - \mathbf{p}_k)$  appearing in (1.3), (2.2) resp. (2.5) in the form of linear combination of the weights due to the identity  $\mathbf{x} = \sum_{i=1}^3 \lambda_i(\mathbf{x}) \mathbf{p}_i$  as

$$A_k(\mathbf{x} - \mathbf{p}_k) = \sum_{i=1}^3 A_k(\mathbf{p}_i - \mathbf{p}_k) \lambda_i.$$

*Definition. 2.6.* Henceforth we say that  $\Psi = [\Psi_0, \Psi_1]$  is a *pair of admissible shape functions* if  $\Psi_0, \Psi_1 \in \mathcal{C}^1([0, 1])$  and (2.1) holds. We also say for short that  $\{(\mathbf{p}_k, f_k, A_k)\}_{k=1}^3$  is an *admissible function germ* if  $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3 \in \mathbb{R}^2$  form a non-degenerate triangle,  $f_k \in \mathbb{R}$  and  $A_k \in \mathcal{L}(\mathbb{R}^2, \mathbb{R})$ .

As an immediate consequence of (2.3), since

$$\begin{aligned} \lambda_i(\mathbf{p}_j) &= \delta_{ij}, & G_i(\mathbf{p}_j - \mathbf{p}_k) &= \delta_{ij} - \delta_{ik}, \\ \lambda_i(\mathbf{x}_t^k) &= t, & \lambda_j(\mathbf{x}_t^k) &= 1 - t, & \lambda_k(\mathbf{x}_t^k) &= 0, \\ \mathbf{x}_t^k - \mathbf{p}_i &= (1 - t)(\mathbf{p}_j - \mathbf{p}_i), & \mathbf{x}_t^k - \mathbf{p}_j &= t(\mathbf{p}_i - \mathbf{p}_j), \end{aligned} \quad (2.7)$$

we obtain the following observation.

**Lemma. 2.8.** *Given an admissible pair  $\Psi = [\Psi_0, \Psi_1]$  of shape functions along with an admissible function germ  $\mathfrak{g} = \{(\mathbf{p}_k, f_k, A_k)\}_{k=1}^3$ , if  $i, j, k \in \{1, 2, 3\}$  are three different indices then, at the generic point*

$$\mathbf{x}_t^k := t\mathbf{p}_+(1-t)\mathbf{p}_j \quad (0 \leq t \leq 1) \quad (2.9)$$

*of the edge  $[\mathbf{p}_i, \mathbf{p}_j]$  in the triangle  $\mathbf{T} = \text{Conv}\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$ , for the function  $F_0 = \mathfrak{F}_0^\Psi \mathfrak{g}$  we have*

$$\begin{aligned} F_0(\mathbf{x}_t^k) &= \Psi_0(t)f_i + \Psi_1(t)(1-t)A_i(\mathbf{p}_j - \mathbf{p}_i) + \\ &\quad + \Psi_0(1-t)f_j + \Psi_1(1-t)tA_j(\mathbf{p}_i - \mathbf{p}_j); \\ F_0'(\mathbf{x}_t^k) &= [\Psi_0'(t)f_i + \Psi_1'(t)(1-t)A_i(\mathbf{p}_j - \mathbf{p}_i)]G_i + \Psi_1(t)A_i + \\ &\quad + [\Psi_0'(1-t)f_j + \Psi_1'(1-t)tA_j(\mathbf{p}_i - \mathbf{p}_j)]G_j + \Psi_1(1-t)A_j. \end{aligned} \quad (2.10)$$

### 3. Generic algorithm of reduced side derivatives (RSD)

In this section we are looking for  $\mathcal{C}^1$ -spline interpolation procedures analogous to those described in [5] but with more general shape functions  $\Psi_0, \Psi_1 \in \mathcal{C}^1[0, 1]$  instead of  $\Phi, \Theta$  there.

Henceforth let  $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3 \in \mathbb{R}^2$  be the vertices of some (arbitrarily fixed) non-degenerate triangle  $\mathbf{T}$  with respective weight functions  $\lambda_m : \mathbb{R}^2 \rightarrow \mathbb{R}$  and derivative weights  $G_m := \lambda_m' \in \mathcal{L}(\mathbb{R}^2, \mathbb{R})$  ( $m=1, 2, 3$ ) and let  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3 \in \mathbb{R}^2$  be given vectors such that  $\mathbf{u}_k \not\parallel \mathbf{p}_i - \mathbf{p}_j$  whenever  $(i, j, k) \in S_3$ .

Furthermore  $\Psi = [\Psi_0, \Psi_1]$  resp.  $\mathfrak{g} = \{(\mathbf{p}_k, f_k, A_k)\}_{k=1}^3$  denote a fixed admissible pair of shape functions and a function germ. As earlier, we write

$$F_0 = \mathfrak{F}_0^\Psi \mathfrak{g}.$$

Our starting point is the following immediate consequence of (2.10):

*Remark. 3.1.* Let  $H \in \mathcal{C}_0^1(\mathbf{T})$ . Then the function  $F := F_0 - H$  coincides with  $F_0$  along the edges of  $\mathbf{T}$  and hence, at the vertices, it has also the properties  $F(\mathbf{p}_i) = f_i$ ,  $F(\mathbf{p}_i) = A_i$  ( $i = 1, 2, 3$ ) analogous to (2.5).

*Definition. 3.2.* We say that the function

$$F = F_0 - H, \quad H \in \mathcal{C}_0^1(\mathbf{T}) \quad (3.3)$$

is an *RSD modification* of  $F_0$  (with respect to the directions  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  along the edges  $[\mathbf{p}_2, \mathbf{p}_3], [\mathbf{p}_3, \mathbf{p}_1], [\mathbf{p}_1, \mathbf{p}_2]$  respectively) if for any index permutation  $(i, j, k) \in S_3$ ,

$$F'(\mathbf{x})\mathbf{u}_k = \Psi_1(\lambda_i(\mathbf{x}))A_i\mathbf{u}_k + \Psi_1(\lambda_j(\mathbf{x}))A_j\mathbf{u}_k \text{ whenever } \mathbf{x} \in [\mathbf{p}_i, \mathbf{p}_j].$$

Conveniently, in this case we refer to  $H$  as an *RSD modifier* of  $F_0$ .

According to Lemma 2.8, we have the following.

**Corollary 3.4.** *A function  $H \in \mathcal{C}_0^1(\mathbf{T})$  is an RSD modifier of  $F_0$  with respect to the directions  $\mathbf{u}_k$  along the edges  $[\mathbf{p}_i, \mathbf{p}_j]$  if and only if*

$$\begin{aligned} H'(\mathbf{x}_t^k)\mathbf{u}_k &= [\Psi_0'(t)f_i + \Psi_1'(t)(1-t)A_i(\mathbf{p}_j - \mathbf{p}_i)][G_i\mathbf{u}_k] + \\ &+ [\Psi_0'(1-t)f_j + \Psi_1'(1-t)tA_j(\mathbf{p}_i - \mathbf{p}_j)][G_j\mathbf{u}_k] \end{aligned}$$

whenever  $\mathbf{x}_t^k = t\mathbf{p}_i + (1-t)\mathbf{p}_j$  with  $(i, j, k) \in S_3$  and  $0 \leq t \leq 1$ .

**Theorem 3.5.** *Let  $\Omega \subset \mathbb{R}^3$  be a set whose interior contains  $\Delta_3 \cap (0, 1)^3$ . Assume  $\chi_0, \chi_1 \in \mathcal{C}^1(\Omega)$  are functions vanishing along the edges  $\Delta_{3,k} = \{(t_1, t_2, t_3) \in \Delta_3 : t_k = 0\}$  ( $k = 1, 2, 3$ ) of  $\Delta_3$  such that*

$$D_3\chi_0(t, 1-t, 0) = \Psi_0'(t), \quad D_3\chi_1(t, 1-t, 0) = \Psi_1'(t) \cdot (1-t) \quad (3.6)$$

with the following marginal conditions on the derivatives

$$\chi_r'(\mathbf{t}) = 0 \quad (\mathbf{t} \in \Delta_{3,1} \cup \Delta_{3,2}); \quad D_m\chi_r(\mathbf{t}) = 0 \quad (\mathbf{t} \in \Delta_{3,3}, m = 1, 2). \quad (3.7)$$

Then the function  $H$  in (1.3) is an RSD modifier for  $F_0$  under any choice of the vectors  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  with  $\mathbf{u}_i \nparallel \mathbf{p}_j - \mathbf{p}_k$  ( $(i, j, k) \in S_3$ ).

PROOF. Consider any permutation  $(i, j, k) \in S_3$ . According to Corollary 3.4, it suffices to see that, at the points  $\mathbf{x}_t = t\mathbf{p}_i + (1-t)\mathbf{p}_j$  we have

$$\begin{aligned} H(\mathbf{x}_t) &= 0, \\ H'(\mathbf{x}_t)\mathbf{u}_k &= [\Psi_0'(t)f_i + \Psi_1'(t)(1-t)A_i(\mathbf{p}_j - \mathbf{p}_i)]G_i\mathbf{u}_k + \\ &+ [\Psi_0'(1-t)f_j + \Psi_1'(1-t)tA_j(\mathbf{p}_i - \mathbf{p}_j)]G_j\mathbf{u}_k. \end{aligned} \quad (3.8)$$

Notice that, in terms of the triples

$$\mathbf{t}_{\ell, m, n} = (\lambda_\ell(\mathbf{x}_t), \lambda_m(\mathbf{x}_t), \lambda_n(\mathbf{x}_t)) \quad ((\ell, m, n) \in S_3),$$

and by setting  $(\ell, m, n)_1 = \ell$ ,  $(\ell, m, n)_2 = m$ ,  $(\ell, m, n)_3 = n$ , we can write

$$\begin{aligned} H(\mathbf{x}_t) &= \sum_{(\ell, m, n) \in \mathcal{S}_3} \frac{G_\ell \mathbf{u}_n}{G_n \mathbf{u}_n} \left[ f_\ell \chi_0(\mathbf{t}_{\ell, m, n}) + A_\ell(\mathbf{p}_m - \mathbf{p}_\ell) \chi_1(\mathbf{t}_{\ell, m, n}) \right], \\ H'(\mathbf{x}_t) \mathbf{u}_k &= \sum_{(\ell, m, n) \in \mathcal{S}_3} \frac{G_\ell \mathbf{u}_n}{G_n \mathbf{u}_n} \left[ f_\ell \left[ \sum_{q=1}^3 D_q \chi_0(\mathbf{t}_{\ell, m, n}) G_{(\ell, m, n)_q} \mathbf{u}_k \right] + \right. \\ &\quad \left. + A_\ell(\mathbf{p}_m - \mathbf{p}_\ell) \left[ \sum_{q=1}^3 D_q \chi_1(\mathbf{t}_{\ell, m, n}) G_{(\ell, m, n)_q} \mathbf{u}_k \right] \right]. \end{aligned}$$

Observe that we have  $\mathbf{t}_{ijk} = (t, 1-t, 0) \in \Delta_{3,3}$ ,  $\mathbf{t}_{jik} = (1-t, t, 0) \in \Delta_{3,3}$ ,  $\mathbf{t}_{ikj} = (t, 0, 1-t) \in \Delta_{3,2}$ ,  $\mathbf{t}_{kij} = (0, t, 1-t) \in \Delta_{3,2}$ ,  $\mathbf{t}_{jki} = (1-t, 0, t) \in \Delta_{3,1}$ ,  $\mathbf{t}_{kji} = (0, 1-t, t) \in \Delta_{3,1}$ . Hence the relation  $H(\mathbf{x}_t) = 0$  is immediate. According to (3, 7), all the terms  $D_q \chi_r(\mathbf{t}_{\ell, m, n})$  vanish except for the cases with  $\lambda_n(\mathbf{x}_t) = 0$  and  $q = 3$  that is for  $(q, r, \ell, m, n) \in \{(3, r, i, j, k), (3, r, j, i, k) : r = 0, 1\}$ . In view of (3.6) we have  $D_3 \chi_0(\mathbf{t}_{ijk}) = \Psi'_0(t)$ ,  $D_3 \chi_0(\mathbf{t}_{jik}) = \Psi'_0(1-t)$ ,  $D_3 \chi_1(\mathbf{t}_{ijk}) = \Psi'_1(t)(1-t)$ ,  $D_3 \chi_1(\mathbf{t}_{jik}) = \Psi'_1(1-t)t$  which completes the proof of (3.8) and hence the theorem.  $\square$

*Definition. 3.9.* We shall say that  $[\Psi_0, \Psi_1, \chi_0, \chi_1]$  is an *RSD tuple* if  $[\Psi_0, \Psi_1]$  is an admissible pair of shape functions and  $\chi_0, \chi_1$  are functions satisfying the requirements of Theorem 3.5.

**Lemma. 3.10.** *Given any function  $h \in \mathcal{C}_0(\mathbb{R}_+^2)$ , there exists  $\chi \in \mathcal{C}_0^1(\mathbb{R}_+^3)$  such that for all  $t_1, t_2, t_3 \geq 0$  we have*

$$\begin{aligned} 0 &= D_m \chi(0, t_2, t_3) = D_m(t_1, 0, t_3) = D_m(t_1, t_2, 0) \quad (m = 1, 2), \\ h(t_1, t_2) &= D_3 \chi(t_2, t_2, 0), \quad 0 = D_3 \chi(0, t_2, t_3) = D_3 \chi(t_1, 0, t_3). \end{aligned}$$

PROOF. For  $\mathbf{t} = (t_1, t_2, t_3) \in \mathbb{R}_+^3$ , define the smoothing

$$\widehat{h}(\mathbf{t}) = \frac{1}{t_3^2} \int_{s_1=t_1}^{t_1+t_3} \int_{s_2=t_2}^{t_2+t_3} h(s_1, s_2) ds_2 ds_1 \quad (t_3 > 0), \quad \widehat{h}(\mathbf{t}) = h(t_1, t_2) \quad (t_3 = 0)$$

of  $h(t_1, t_2, 0)$  and let us fix a function  $\phi \in \mathcal{C}^1([0, 1])$  (e.g.  $\phi(t) = 3t^2 - 2t^3$ ) with bounded derivative and being such that  $\phi(0) = \phi'(0) = \phi'(1) = 0$  and  $\phi(1) = 1$ . We show that the function

$$\chi(t_1, t_2, t_3) = \phi\left(\frac{t_1}{t_1+t_3}\right) \phi\left(\frac{t_2}{t_2+t_3}\right) t_3 \widehat{h}(t_1, t_2, t_3) \quad (t_1, t_2 \geq 0, t_3 > 0)$$



extended with  $\chi(t_1, t_2, 0) = 0$  suits the requirements of the lemma.

It is folklore that, by the Newton-Leibniz theorem, the function  $\widehat{h}$  is continuous, moreover its restriction to  $\mathbb{R}_+^2 \times \mathbb{R}_{++}$  ( $= \{(t_1, t_2, t_3) : t_1, t_2 \geq 0, t_3 > 0\}$ ) is  $\mathcal{C}^1$  smooth. In particular, for any point  $\mathbf{t} = (t_1, t_2, t_3)$  with  $t_3 > 0$  and  $t_1 = 0$  or  $t_2 = 0$  we have  $\chi(\mathbf{t}) = 0$  and  $\chi'(\mathbf{t}) = 0$ . Furthermore, for any point  $\mathbf{t} \in \mathbb{R}_+ \times \mathbb{R}_{++}$ , with the indices  $m = 1, 2$  resp. 3 we can write

$$\begin{aligned} D_m \chi(\mathbf{t}) &= \Phi\left(\frac{t_{3-m}}{t_{3-m}+t_3}\right) \left[ \Phi'\left(\frac{t_m}{t_m+t_3}\right) \frac{t_3^2}{(t_m+t_3)^2} \widehat{h}(\mathbf{t}) + \Phi\left(\frac{t_m}{t_m+t_3}\right) \frac{\partial[t_3 \widehat{h}]}{\partial t_1} \right], \\ D_3 \chi(\mathbf{t}) &= -\Phi'\left(\frac{t_1}{t_1+t_3}\right) \Phi\left(\frac{t_2}{t_2+t_3}\right) \frac{t_1 t_3}{(t_1+t_3)^2} \widehat{h}(\mathbf{t}) - \\ &\quad - \Phi\left(\frac{t_1}{t_1+t_3}\right) \Phi'\left(\frac{t_2}{t_2+t_3}\right) \frac{t_2 t_3}{(t_2+t_3)^2} \widehat{h}(\mathbf{t}) + \Phi\left(\frac{t_1}{t_1+t_3}\right) \Phi\left(\frac{t_2}{t_2+t_3}\right) \frac{\partial[t_3 \widehat{h}]}{\partial t_3}. \end{aligned}$$

Therefore it only remains to prove that given any point  $\mathbf{t}^* = (t_1^*, t_2^*, 0) \in \mathbb{R}_+^3$  we have

$$\begin{aligned} D_m \chi(\mathbf{t}) &\rightarrow 0 \quad (m=1, 2), \quad D_3 \chi(\mathbf{t}) \rightarrow h(t_1^*, t_2^*) \\ \text{whenever } &\mathbb{R}_+^2 \times \mathbb{R}_{++} \ni \mathbf{t} \rightarrow \mathbf{t}^*. \end{aligned} \tag{3.11}$$

Actually, these relation follow from the mean value expressions

$$\begin{aligned} \frac{\partial[t_3 \widehat{h}]}{\partial t_m} &= \frac{1}{t_3} \int_{s_2=t_{3-m}}^{t_{3-m}+t_3} [h(t_m+t_3, s_2) - h(t_m, s_2)] ds_2 = \\ &= h(t_m+t_3, r_{3-m}(\mathbf{t})) - h(t_m, q_{3-m}(\mathbf{t})), \\ \frac{\partial[t_3 \widehat{h}]}{\partial t_3} &= -\frac{1}{t_3^2} \int_{s_1=t_1}^{t_1+t_3} \int_{s_2=t_2}^{t_2+t_3} h(s_1, s_2) ds_2 ds_1 + \\ &\quad + \frac{1}{t_3} \int_{s_2=t_2}^{t_2+t_3} h(t_1+t_3, s_2) ds_2 + \frac{1}{t_3} \int_{s_1=t_1}^{t_1+t_3} h(s_1, t_2+t_3) ds_1 = \\ &= -h(p_1(\mathbf{t}), p_2(\mathbf{t})) + h(t_1+t_3, r_2(\mathbf{t})) + h(r_1(\mathbf{t}), t_2+t_3) \end{aligned}$$

with suitable

$$p_1(\mathbf{t}), q_1(\mathbf{t}), r_1(\mathbf{t}) \in [t_1, t_1+t_3], \quad p_2(\mathbf{t}), q_2(\mathbf{t}), r_2(\mathbf{t}) \in [t_2, t_2+t_3].$$

Indeed, let  $\mathbb{R}_+^2 \times \mathbb{R}_{++} \ni \mathbf{t} \rightarrow \mathbf{t}^* = (0, t_2^*, t_3^*)$ . Then  $\mathbf{p}(\mathbf{t}) \rightarrow (t_1^*, t_2^*)$ ,  $\mathbf{q}_m(\mathbf{t}), \mathbf{r}_m(\mathbf{t}) \rightarrow t_m^*$  ( $m = 1, 2$ ), whence  $\partial[t_3 \widehat{h}]/\partial t_m \rightarrow 0$  ( $m = 1, 2$ ) and  $\partial[t_3 \widehat{h}]/\partial t_3 \rightarrow h(t_1^*, t_2^*)$ . Since the function  $(s, t) \mapsto s/(s+t)$  is analytic on the open half plane  $\{(s, t) \in \mathbb{R}^2 : s+t > 0\}$  containing the rays  $R_1 = \mathbb{R}_{++} \times \{0\}$  resp.  $R_2 = \{0\} \times \mathbb{R}_{++}$ , (3.11) is immediate in the cases  $\mathbf{t}^* \in R_m$  ( $m = 1, 2$ ). If  $\mathbb{R}_+^2 \times \mathbb{R}_{++} \ni \mathbf{t} \rightarrow (0, 0, 0)$ , we can deduce (3.11) from the facts that, for  $\ell = 1, 2, 3$ , the functions  $t_\ell t_3 / (t_\ell + t_3)^2$ ,  $t_3^2 / (t_\ell + t_3)^2$  resp.  $\Phi(t_\ell / (t_\ell + t_3))$ ,  $\Phi'(t_\ell / (t_\ell + t_3))$  are bounded, furthermore  $\partial[t_3 \widehat{h}]/\partial t_\ell \rightarrow h(0, 0) = 0$ .  $\square$

**Corollary. 3.12.** *By Theorem 3.5 and Lemma 3.10, for any admissible pair  $[\Psi_0, \Psi_1]$ , there exist  $\chi_0, \chi_1 \in \mathcal{C}_0^1(\mathbb{R}_+^3 \setminus \{(0, 0, 0)\})$  such that  $[\Psi_0, \Psi_1, \chi_0, \chi_1]$  is an RSD-tuple.*

*Remark. 3.13.* Unfortunately, our convolution type generic construction provided by the proof of Lemma 3.10 is far from being optimal in most cases from numerical view points. In contrast, in view of Theorem 3.5, in many cases we may apply the following satisfactory construction:

**Proposition. 3.14.** *If  $[\Psi_0, \Psi_1]$  is an admissible pair of shape functions and  $h_0, h_1 \in \mathcal{C}_0^1(\mathbb{R}_+^2)$  are functions such that  $h(t, 1-t) = \Psi'_0(t)$  and  $h(t, 1-t) = \Psi'_1(t)(1-t)$  ( $0 \leq t \leq 1$ ) then  $[\Psi_0, \Psi_1, \chi_0, \chi_1]$  is an RSD tuple with*

$$\chi_r(t_1, t_2, t_3) = h_r(t_1, t_2)t_3 \quad (r = 0, 1).$$

*Example. 3.15.* Cases with a factorization

$$\Psi'_0(t) = w_{01}(t)w_{02}(1-t), \quad \Psi'_1(t)(1-t) = w_{11}(t)w_{12}(1-t)$$

for suitable functions  $w_{rk} \in \mathcal{C}_0^1(\mathbb{R}_+)$  such that  $w_{rk}(0) = w'_{rk}(0) = 0$ .

(a) The procedure (1.1) with the shape functions (1.2) corresponds to the case  $[\Psi_0, \Psi_1] = [\Phi, \Theta]$ ,  $\chi_0(t_1, t_2, t_3) = 30t_1^2t_2^2t_3$ ,  $\chi_1(t_1, t_2, t_3) = 12t_1^2t_2^2t_3$  with  $w_{01}(t) = 30t^2$ ,  $w_{11}(t) = 12t^2$ ,  $w_{02}(t) = w_{12}(t) = t^2$  since  $\Phi'(t) = 30t^2(1-t)^2$  and  $\Theta'(t) = 12t^2(1-t)$ .

(b)  $[\Phi, \Phi, 30t_1^2t_2^2t_3, 30t_1^2t_2^2t_3]$  is an RSD tuple by Proposition 3.14, corresponding to the case  $w_{01}(t) = w_{11}(t) = 30t^2$ ,  $w_{02}(t) = t^2$ ,  $w_{12}(t) = t^3$ .

*Remark. 3.16.* It is an easy consequence of Whitney's extension theorem [7] that for any function  $\psi \in \mathcal{C}([0, 1])$  with  $\psi(0) = \psi'(0) = \psi'(1) = 0$  there exists a function  $h \in \mathcal{C}_0^1(\mathbb{R}_+^2)$  such that  $\psi'(t) = h(t, 0-t)$ . Hence every admissible pair  $[\Psi_0, \Psi_1]$  with the property  $\Psi'_0(1) = 0$  admits functions  $h_0, h_1 \in \mathcal{C}(\mathbb{R}_+^2)$  such that  $[\Psi, \Psi_1, h_0(t_1, t_2)t_3, h_0(t_1, t_2)t_3]$  is an RSD tuple.

We complete this section with a brief description of the local approximation operator corresponding to Theorem 3.5 and its use in constructing  $\mathcal{C}^1$ -splines over a 2D triangular mesh analogously as done in [5, Algorithm].

*Definition. 3.17.* Let  $\mathbf{\Pi} = [\Psi_0, \Psi_1, \chi_0, \chi_1]$  be an RSD tuple and write  $\mathbf{\Psi} = [\Psi_0, \Psi_1]$  resp.  $\mathbf{X} = [\chi_0, \chi_1]$ . We introduce the *SDR modification of  $\mathfrak{F}^{\mathbf{\Psi}}$  by means of the complementary shape functions  $\mathbf{X}$*  as the operator

$$\mathfrak{F}^{\mathbf{\Pi}}: \{(\mathbf{p}_k, f_k, A_k, \mathbf{u}_k)\}_{k=1}^3 \mapsto \mathfrak{F}^{\mathbf{\Psi}}\{(\mathbf{p}_k, f_k, A_k)\}_{k=1}^3 - \mathfrak{H}^{\mathbf{X}}\{(\mathbf{p}_k, f_k, A_k, \mathbf{u}_k)\}_{k=1}^3$$

defined for all tuples  $\{(\mathbf{p}_k, f_k, A_k, \mathbf{u}_k)\}_{k=1}^3$  where  $\{(\mathbf{p}_k, f_k, A_k)\}_{k=1}^3$  is a non-degenerate function germ and  $\mathbf{u}_k \parallel \mathbf{p}_i - \mathbf{p}_j$  ( $(i, j, k) \in S_3$ ) and

$$\mathfrak{H}^{\mathbf{X}}\{(\mathbf{p}_k, f_k, A_k, \mathbf{u}_k)\}_{k=1}^3 = \left[ H \text{ in Theorem 3.5} \right].$$

Recall [5] that, given a set  $\mathbf{V} = \{\mathbf{v}_1, \dots, \mathbf{v}_R\}$  of points in  $\mathbb{R}^2$ , by a *triangular mesh* over  $\mathbf{V}$  we mean a family  $\{\mathbf{T}_1, \dots, \mathbf{T}_M\} \subset \mathbb{R}^2$  of non-degenerate triangles of the form

$$\mathbf{T}_m = \text{Conv}\{\mathbf{v}_{i(m,1)}, \mathbf{v}_{i(m,2)}, \mathbf{v}_{i(m,3)}\}, \quad i(m,1) < i(m,2) < i(m,3)$$

with pairwise disjoint interiors whose intersections are either empty or a common vertex or a common edge, furthermore  $\mathbf{V} \subset \bigcup_{m=1}^M \mathbf{T}_m$ . A crucial ingredient of the construction, we enumerate the edges of the mesh triangles  $\mathbf{T}_m$  in the form  $\mathbf{E}_1, \dots, \mathbf{E}_S$  and associate a vector  $\mathbf{u}_s \parallel \mathbf{E}_s$  with each edge  $\mathbf{E}_s$ , furthermore let  $s(m, k)$  denote the index of the *opposite* edge to the vertex  $\mathbf{v}_{m(i,k)}$  in the triangle  $\mathbf{T}_m$ . By replacing  $\Phi, \Theta$  with  $\Psi_0, \Psi_1$ , respectively the functions  $30\lambda_i^2\lambda_j^2\lambda_k$ ,  $12\lambda_i^2\lambda_j^2\lambda_k$  with  $\chi_0, \chi_1$  in a straightforward manner in [5, Proof of Thm. 2], we can conclude the following: Given any germ  $\{(\mathbf{v}_i, f_i, A_i)\}_{i=1}^R$  of first order function data over  $\mathbf{V}$ , the union of the functions

$$F_m = \mathfrak{F}^{\Pi}\left\{(\mathbf{v}_{i(m,k)}, f_{i(m,k)}, A_{i(m,k)}, \mathbf{u}_{s(m,k)})\right\}_{k=1}^3 \in \mathcal{C}^1(\mathbf{T}_m) \quad (m=1, \dots, M)$$

is continuously differentiable on the mesh domain  $\mathbf{D} = \bigcup_{m=1}^M \mathbf{T}_m$ .

#### 4. Range shift property

One of the most frequent applications of splines over triangular meshes is reconstructing approximately a smooth surface with a function graph passing through a family of points. Such procedures correspond to a model of the following pattern: We are given a smooth surface  $\mathbf{S} \subset \mathbb{R}^3$  which can be represented in the form  $\mathbf{S} = \text{graph}(f) = \{[\mathbf{p}, f(\mathbf{p})] : \mathbf{p} \in \mathbb{R}^2\}$  with some (a priori unknown) function  $f \in \mathcal{C}^1(\mathbb{R}^2)$ . Accurate data (obtained with scanner equipments) are available for a finite family of points  $\mathbf{v}_1, \dots, \mathbf{v}_N \in \mathbb{R}^2$  concerning the *first order data*  $f^{[1]}(\mathbf{p}) = [\mathbf{p}, f(\mathbf{p}), f'(\mathbf{p})]$  and we approximate a piece of  $\mathbf{S}$  with the graph of a  $\mathcal{C}^1$ -spline function  $F \in \mathcal{C}^1(\mathbf{D})$  on the domain  $\mathbf{D}$  of a triangular mesh with vertices  $\mathbf{v}_n$ . The procedures used in constructing  $F$  are usually investigated from

the view points of estimating various biases between  $\text{graph}(F)$  and  $\mathbf{S}$ , respectively from exhibiting features of algebraic-geometric character. In this note we only discuss two kinds of the last category in the context of RSD methods: if the the spline graphs are invariant with respect to translations of first order data and the property that plane surfaces remain invariant.

*Definition. 4.1.* Throughout this section  $[\mathbf{P}, \mathbf{U}]$  denotes an admissible pair of triples  $\mathbf{P} = \{\mathbf{p}_i\}_{i=1}^3$ ,  $\mathbf{U} = \{\mathbf{u}_i\}_{i=1}^3$  with  $\mathbf{p}_i, \mathbf{u}_i \in \mathbb{R}^2$ . We write  $\mathbf{T} = \text{Conv}(\mathbf{P})$  and  $\lambda_1, \lambda_2, \lambda_3$  resp.  $G_1, G_2, G_3$  for the weight functions resp. their derivatives associated with the vertices of the triangle  $\mathbf{T}$ . Given any function  $f \in \mathcal{C}^1(\mathbb{R}^2)$ , we shall use the shorthand notations

$$f_{\mathbf{P}, \mathbf{U}}^{[1]} = \{[\mathbf{p}_i, f(\mathbf{p}_i), f'(\mathbf{p}_i), \mathbf{u}_i]\}_{i=1}^3, \quad \mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\Pi} f = \mathfrak{F}^{\Pi} f_{\mathbf{P}, \mathbf{U}}^{[1]}.$$

An RSD-tuple  $\Pi = [\Psi_0, \Psi_1, \chi_0, \chi_1]$  has the *range shift property* if the procedure  $\mathfrak{F}^{\Pi}$  leaves the constant functions invariant, that is if for all admissible pairs  $\mathbf{P}, \mathbf{U}$  and the constant unit function  $\mathbf{1} : \mathbb{R}^2 \ni \mathbf{x} \mapsto 1$  we have

$$\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\Pi} \mathbf{1} = \mathbf{1} |_{\text{Conv}(\mathbf{P})}$$

*Remark. 4.2.* Since  $\mathbf{1}(\mathbf{p}_i) = 1$ ,  $\mathbf{1}'(\mathbf{p}_i)\mathbf{u}_i = 0$ , we can write

$$\begin{aligned} \mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\Pi} \mathbf{1} &= \sum_{i=1}^3 \Psi_0(\lambda_i) - \sum_{(\ell, m, n) \in \mathcal{S}_3} \frac{G_{\ell} \mathbf{u}_n}{G_n \mathbf{u}_n} \chi_0(\lambda_{\ell}, \lambda_m, \lambda_n) = \\ &= \sum_{n=1}^3 \left\{ \Psi_0(\lambda_n) - \left[ \frac{G_{\ell_n} \mathbf{u}_n}{G_n \mathbf{u}_n} \chi_0(\lambda_{\ell_n}, \lambda_{m_n}, \lambda_n) + \frac{G_{m_n} \mathbf{u}_n}{G_n \mathbf{u}_n} \chi_0(\lambda_{m_n}, \lambda_{\ell_n}, \lambda_n) \right] \right\} \end{aligned}$$

with  $(\ell_1, m_1) = (2, 3)$ ,  $(\ell_2, m_2) = (1, 3)$ ,  $(\ell_3, m_3) = (1, 2)$ .

**Lemma. 4.3.** For any  $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$  there exist  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3 \in \mathbb{R}^2$  such that

$$\left[ \frac{G_i \mathbf{u}_j}{G_j \mathbf{u}_j} \right]_{i,j=1}^3 = \begin{bmatrix} 1 & -1 - \alpha_2 & \alpha_3 \\ \alpha_1 & 1 & -1 - \alpha_3 \\ -1 - \alpha_1 & \alpha_2 & 1 \end{bmatrix}.$$

PROOF. A suitable choice is  $\mathbf{u}_1 = \mathbf{p}_1 - \frac{1}{2}[\mathbf{p}_2 + \mathbf{p}_3] + (\frac{1}{2} + \alpha_1)[\mathbf{p}_2 - \mathbf{p}_3]$ ,  $\mathbf{u}_2 = \mathbf{p}_2 - \frac{1}{2}[\mathbf{p}_3 + \mathbf{p}_1] + (\frac{1}{2} + \alpha_2)[\mathbf{p}_3 - \mathbf{p}_1]$ ,  $\mathbf{u}_3 = \mathbf{p}_3 - \frac{1}{2}[\mathbf{p}_1 + \mathbf{p}_2] + (\frac{1}{2} + \alpha_3)[\mathbf{p}_1 - \mathbf{p}_2]$ .  $\square$

**Proposition. 4.4.** An RSD tuple  $\Pi = [\Psi_0, \Psi_1, \chi_0, \chi_1]$  has range shift property if and only if

$$\begin{aligned} \Psi_0(t) + \Psi_0(1-t) &= 1, \quad \Psi_0'(t) = \Psi_0'(1-t) & (0 \leq t \leq 1), \\ \chi_0(t_1, t_2, t_3) &= \chi_0(t_2, t_1, t_3) & ((t_1, t_2, t_3) \in \Delta_3) \end{aligned} \tag{4.5}$$

$$\text{and } \mathbf{1} = \sum_{i=1}^3 \Psi_0(\lambda_i) + \sum_{(\ell,m,n) \in S_3^+} \chi_0(\lambda_\ell, \lambda_m, \lambda_n).$$

PROOF. Consider the points  $\mathbf{x}_t = t\mathbf{p}_1 + (1-t)\mathbf{p}_2$  ( $0 \leq t \leq 1$ ). Since  $\mathbf{1}' \equiv 0$ ,  $\mathbf{1}(\mathbf{x}_t) = 1$ ,  $\lambda_1(\mathbf{x}_t) = t$ ,  $\lambda_2(\mathbf{x}_t) = 1-t$ , and  $\lambda_3(\mathbf{x}_t) = 0$ , by assumption

$$1 = \mathbf{1}(\mathbf{x}_t) = \mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\Pi} \mathbf{1}(\mathbf{x}_t) = \sum_{i=1}^3 \Psi_0(\lambda_i(\mathbf{x}_t)) = \Psi_0(t) + \Psi_0(1-t) + \Psi_0(0)$$

where  $\Psi_0(0) = 0$  due to (2.1). Thus in view of Remark 4.2 and Lemma 4.3, the expression of  $\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\Pi} \mathbf{1}$  has the form

$$\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\Pi} \mathbf{1} = \sum_{n=1}^3 \left\{ \Psi_0(\lambda_n) - [\alpha_n \chi_0(\lambda_{\ell_n}, \lambda_{m_n}, \lambda_n) + (-1 - \alpha_n) \chi_0(\lambda_{m_n}, \lambda_{\ell_n}, \lambda_n)] \right\}$$

where the coefficients  $\alpha_1, \alpha_2, \alpha_3$  may assume any real value. This is possible only

$$\text{if } \mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\Pi} \mathbf{1} = \sum_{i=1}^3 \Psi_0(\lambda_i) + \chi_0(\lambda_\ell, \lambda_m, \lambda_n) + \chi_0(\lambda_\ell, \lambda_m, \lambda_n) + \chi_0(\lambda_\ell, \lambda_m, \lambda_n) \text{ and}$$

$$\chi_0(\lambda_{\ell_n}, \lambda_{m_n}, \lambda_n) = \chi_0(\lambda_{m_n}, \lambda_{\ell_n}, \lambda_n) \quad (n = 1, 2, 3).$$

In particular, for  $n = 3$  and a generic point  $\mathbf{x} = \sum_{k=1}^3 t_k \mathbf{p}_k$  (with  $\sum_k t_k = 1$ ,  $t_k \geq 0$ ) of the triangle  $\text{Conv}(\mathbf{P})$ , we get  $\chi_0(t_1, t_2, t_3) = \chi_0(\lambda_1(\mathbf{x}), \lambda_2(\mathbf{x}), \lambda_3(\mathbf{x})) = \chi_0(\lambda_2(\mathbf{x}), \lambda_1(\mathbf{x}), \lambda_3(\mathbf{x})) = \chi_0(t_2, t_1, t_3)$  which completes the proof.  $\square$

**Lemma. 4.6.** *Given any smooth function  $\phi$  defined on a domain  $\Omega \subset \mathbb{R}^3$  containing  $\Delta_3$  and given any point  $\mathbf{x} \in \mathbf{T}$ , for the function  $f = \phi(\lambda_1, \lambda_2, \lambda_3)$  we have  $f'(\mathbf{x}) = 0$  if and only if*

$$D_1 \phi(\mathbf{t}) = D_2 \phi(\mathbf{t}) = D_3 \phi(\mathbf{t}) \quad \text{where } \mathbf{t} = (\lambda_1(\mathbf{x}), \lambda_2(\mathbf{x}), \lambda_3(\mathbf{x})).$$

PROOF. For any vector  $\mathbf{u} \in \mathbb{R}^2$  we have  $f'(\mathbf{x})\mathbf{u} = \sum_{k=1}^3 D_k \phi(\mathbf{t}) G_k \mathbf{u}$ . Hence the statement is immediate from the relation  $\{(G_1 \mathbf{u}, G_2 \mathbf{u}, G_3 \mathbf{u}) : \mathbf{v} \in \mathbb{R}^2\} = \{(\alpha_1, \alpha_2, \alpha_3) : \sum_k \alpha_k = 0\}$ . (Actually, given any triple  $(i, j, k) \in S_3$  of indices, by taking  $\mathbf{u} = \mathbf{p}_i - \mathbf{p}_j$  we have  $(G_i \mathbf{u} = 1, (G_j \mathbf{u} = -1, (G_k \mathbf{u} = 0$  implying  $D_i \phi(\mathbf{t}) - D_j \phi(\mathbf{t}) = 0$  if  $f'(\mathbf{x})\mathbf{u} = 0$ ).  $\square$

**Corollary. 4.7.** *The RSD tuple  $\Pi = [\Psi_0, \Psi_1, \chi_0, \chi_1]$  has range shift property if and only if (4.5) holds and*

$$\frac{\partial \Sigma}{\partial t_1}(t_1, t_2, t_3) = \frac{\partial \Sigma}{\partial t_2}(t_1, t_2, t_3) = \frac{\partial \Sigma}{\partial t_3}(t_1, t_2, t_3) \quad ((t_1, t_2, t_3) \in \Delta_3), \quad (4.8)$$

$$\Sigma(t_1, t_2, t_3) = \sum_{i=1}^3 \Psi_0(t_i) + \sum_{(\ell,m,n) \in S_3^+} \chi_0(t_\ell, t_m, t_n), \quad S_3^+ = \{(1,2,3), (2,3,1), (3,1,2)\}.$$

PROOF. We established that  $\mathbf{\Pi}$  has range shift property if and only if

$$\Sigma(\lambda_1(\mathbf{x}), \lambda_2(\mathbf{x}), \lambda_3(\mathbf{x})) = 1 \quad (\mathbf{x} \in \text{Conv}(\mathbf{P})). \quad (4.9)$$

Since  $\mathbf{\Pi}$  is an RSD tuple, in any case we have  $\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}}(\mathbf{p}_i) = 1$  ( $i = 1, 2, 3$ ). Hence the relation (4.9) holds if and only if the Fréchet derivative of the function  $\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}} \mathbf{1}$  vanishes, that is if

$$[\Sigma(\lambda_1, \lambda_2, \lambda_3)]'(\mathbf{x}) = 0 \quad (\mathbf{x} \in \text{Conv}(\mathbf{P})). \quad (4.9')$$

An application of Lemma 4.6 with  $\phi = \Sigma$  completes the proof.  $\square$

*Example. 4.10.* The RSD procedures corresponding to tuples of the form  $[t^3(10 - 15t + 6t^2), *, 30t_1^2t_2^2t_3, *]$  as in Examples 3.15 have range shift property. This highly non-trivial fact can easily be established by verifying (4.8) as follows. In such cases, we have

$$\Sigma(t_1, t_2, t_3) = \sum_{i=1}^3 t_i^3(10 - 15t_i + 6t_i^2) + 30(t_1^2t_2^2t_3 + t_2^2t_3^2t_1 + t_3^2t_1^2t_2)$$

and it suffices to show that the polynomial  $\frac{\partial \Sigma}{\partial t_1}$  is symmetric in  $t_1, t_2, t_3$  when replacing  $t_1 = 1 - t_2 - t_3$ . Since  $[t^3(10 - 15t + 6t^2)]' = 30t^2(1-t)^2$ , if  $(t_1, t_2, t_3) \in \Delta_3$  then  $[t_1^3(10 - 15t_1 + 6t_1^2)]' = 30t_1^2(t_2 + t_3)^2$  and

$$\begin{aligned} \frac{1}{30} \frac{\partial \Sigma}{\partial t_1} &= t_1^2(t_2 + t_3)^2 + 2t_1t_2^2t_3 + t_2^2t_3^2 + 2t_3^2t_1t_2 = \\ &= t_1^2t_2^2 + t_1^2t_3^2 + t_2^2t_3^2 + 2t_1t_2t_3(t_1 + t_2 + t_3) = t_1^2t_2^2 + t_1^2t_3^2 + t_2^2t_3^2 + 2t_1t_2t_3. \end{aligned}$$

*Definition. 4.11.* We say that a function  $\Psi \in \mathcal{C}^1([0, 1])$  is *d-symmetric* if

$$\Psi'(t) = \Psi'(1-t) \quad (0 \leq t \leq 1).$$

**Corollary. 4.12.** *If  $\mathbf{\Pi} = [\Psi_0, \Psi_1, \chi_0, \chi_1]$  is an RSD tuple with d-symmetric shape function  $\Psi_0$  then  $\mathbf{\Pi}^{[s]} = [\Psi_0, \Psi_1, \chi_0^{[s]}, \chi_1]$  with*

$$\chi_0^{[s]}(t_1, t_2, t_3) = \frac{1}{2}\chi_0(t_1, t_2, t_3) + \frac{1}{2}\chi_0(t_2, t_1, t_3) \quad (4.13)$$

*is also an RSD tuple such that*

$$\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}^{[s]}} \mathbf{1} = \sum_{(\ell, m, n) \in S_3} \left\{ \frac{1}{2}\Psi_0(\lambda_\ell) + \chi^{[s]}(\lambda_\ell, \lambda_m, \lambda_n) \right\}. \quad (4.14)$$

PROOF. The marginal conditions (3.7) for  $\chi_0^{[s]}$  are immediate. As a consequence of the d-symmetry of  $\Psi_0$ , we have

$$\begin{aligned} D_3\chi_0^{[s]}(t, 1-t, 0) &= \frac{1}{2}D_3\chi_0(t, 1-t, 0) + \frac{1}{2}D_3\chi_0(1-t, t, 0) = \\ &= \frac{1}{2}\Psi_0(t) + \frac{1}{2}\Psi_0(1-t) = \Psi_0(t) \end{aligned}$$

establishing that  $\mathbf{\Pi}^{[s]}$  is an RSD tuple. Due to the symmetry of  $\chi_0^{[s]}$  in the variables  $t_1, t_2$  in Remark 4.2 applied to  $\chi_0^{[s]}$  we can write

$$\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}} \mathbf{1} = \sum_{n=1}^3 \left\{ \Psi_0(\lambda_n) - \left[ \frac{G_{\ell_n} \mathbf{u}_n}{G_n \mathbf{u}_n} + \frac{G_m \mathbf{u}_n}{G_n \mathbf{u}_n} \right] \chi_0(\lambda_{\ell_n}, \lambda_{m_n}, \lambda_n) \right\}.$$

Taking Lemma 4.3 into account, we obtain (4.14).  $\square$

**Theorem. 4.15.** *Let  $[\Psi_0, \Psi_1]$  be an admissible pair of shape functions, with  $\Psi_0$  being d-symmetric. Then we can find  $\chi_0^*, \chi_1^* \in \mathcal{C}_0(\mathbb{R}_+^3)$  such that  $\mathbf{\Pi}^* = [\Psi_0, \Psi_1, \chi_0^*, \chi_1^*]$  is an RSD-tuple with range shift property where  $\chi_0^*$  is symmetric in its first two variables.*

PROOF. According to Theorem 3.5 there exist an RSD tuple of the form  $\mathbf{\Pi} = [\Psi_0, \Psi_1, \chi_0, \chi_1]$  with  $\chi_0, \chi_1 \in \mathcal{C}_0^1(\mathbb{R}_+^3)$ . By Corollary (4.12),  $\mathbf{\Pi}^{[s]} = [\Psi_0, \Psi_1, \chi_0^{[s]}, \chi_1^{[s]}]$  is also an RSD tuple where the complementary shape function  $\chi_0^{[s]}$  is symmetric in its first two variables and such that (4.14) holds. Then, independently of the choice of the vector triple  $\mathbf{U} = [\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3]$ , we can write the difference function of  $\mathbf{1}$  and  $\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}^{[s]}} \mathbf{1}$  in the form

$$\begin{aligned} d &= \mathbf{1} - \mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}^{[s]}} \mathbf{1} = \delta(\lambda_1, \lambda_2, \lambda_3) \quad \text{with} \\ \delta(\mathbf{t}) &= d(\sum_{k=1}^3 t_k \mathbf{p}_k) = \sum_{(\ell, m, n) \in \mathcal{S}_3} \left\{ \frac{1}{6} - \frac{1}{2}\Psi_0(t_\ell) - \chi_0^{[s]}(t_\ell, t_m, t_n) \right\}. \end{aligned} \quad (4.16)$$

Observe that

$$d(\mathbf{x}) = 0, \quad d'(\mathbf{x}) = 0 \quad (\mathbf{x} \in \partial \mathbf{T}). \quad (4.17)$$

Indeed, at the generic point (2.9) of the edge opposite to the vertex  $\mathbf{p}_k$ , by (2.2) we have  $d(\mathbf{x}_t^k) = 1 - [\Psi_0(t) + \Psi_0(1-t)] = 0$ . In particular the functions  $\mathbf{1}$  and  $\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}^{[s]}}$  coincide along the edge  $[\mathbf{p}_i, \mathbf{p}_j]$ , thus  $d'(\mathbf{x}_t^k)(\mathbf{p}_j - \mathbf{p}_i) = 0$  with  $d(\mathbf{x}) = 0$  for  $\mathbf{x} \in \partial \mathbf{T}$ . On the other hand, since trivially  $\mathbf{1}'(\mathbf{p}) = 0$  everywhere, the RSD construction ensures that  $d'(\mathbf{x}_t^k) \mathbf{u}_k = \mathbf{1}'(\mathbf{x}_t^k) \mathbf{u}_k - [\Psi_0(t) \mathbf{1}'(\mathbf{p}_k) \mathbf{u}_k + \Psi_0(1-t) \mathbf{1}'(\mathbf{p}_k) \mathbf{u}_k] = 0$ . Thus

$\delta'(\mathbf{x}_t^k)$  vanishes in two linearly independent directions implying that  $d'(\mathbf{x}_t^k)\mathbf{u} = 0$  ( $\mathbf{u} \in \mathbb{R}^2$ ) i.e.  $d'(\mathbf{x}_t^k) = 0$ .

Notice that  $\mathbf{x} \in \partial\mathbf{T}$  if and only if  $(\lambda_1(\mathbf{x}), \lambda_2(\mathbf{x}), \lambda_3(\mathbf{x})) \in \bigcup_{k=1}^3 \Delta_{3,k}$ . Hence, due to (4.17), we can apply Lemma 4.6 with  $f = d$  and  $\phi = \delta$  to conclude that

$$\delta(\mathbf{t}) = 0, \quad D_1\delta(\mathbf{t}) = D_2\delta(\mathbf{t}) = D_3\delta(\mathbf{t}) \quad \text{whenever } \mathbf{t} \in \bigcup_{k=1}^3 \Delta_{3,k}. \quad (4.17')$$

On the domain  $\Omega := \mathbb{R}_+^3 \setminus \{0\}$ , define

$$\begin{aligned} \chi_0^* &= \chi_0^{[s]} + \frac{1}{6}\widehat{\delta} \quad \text{where} \\ \widehat{\delta}(\mathbf{t}) &= \delta\left(\frac{t_1}{t_1+t_2+t_3}, \frac{t_2}{t_1+t_2+t_3}, \frac{t_3}{t_1+t_2+t_3}\right). \end{aligned}$$

By writing  $s(\mathbf{t}) = t_1 + t_2 + t_3$  for short, for  $k = 1, 2, 3$  we have

$$\begin{aligned} D_k\widehat{\delta}(\mathbf{t}) &= \frac{\partial}{\partial t_k} \delta\left(\frac{t_1}{s(\mathbf{t})}, \frac{t_2}{s(\mathbf{t})}, \frac{t_3}{s(\mathbf{t})}\right) = \\ &= D_1\delta\left(\frac{\mathbf{t}}{s(\mathbf{t})}\right) \frac{\partial}{\partial t_k} \frac{t_1}{s(\mathbf{t})} + D_2\delta\left(\frac{\mathbf{t}}{s(\mathbf{t})}\right) \frac{\partial}{\partial t_k} \frac{t_2}{s(\mathbf{t})} + D_3\delta\left(\frac{\mathbf{t}}{s(\mathbf{t})}\right) \frac{\partial}{\partial t_k} \frac{t_3}{s(\mathbf{t})}. \end{aligned}$$

In particular, if  $0 \neq \mathbf{t} \in \mathbb{R}_+^3$  then  $s(\mathbf{t})^{-1}\mathbf{t} \in \Delta_3$ , furthermore  $s(\mathbf{t})^{-1}\mathbf{t} \in \bigcup_{k=1}^3 \Delta_{3,k}$  whenever  $0 \neq \mathbf{t} \in \partial\mathbb{R}_+^3$ . Thus, as a consequence of (4.17), we get

$$D_k\widehat{\delta}(\mathbf{t}) = \sum_{\ell=1}^3 D_1\delta\left(\frac{\mathbf{t}}{s(\mathbf{t})}\right) \frac{\partial}{\partial t_k} \frac{t_\ell}{s(\mathbf{t})} = D_1\delta\left(\frac{\mathbf{t}}{s(\mathbf{t})}\right) \frac{\partial}{\partial t_k} \frac{s(\mathbf{t})}{s(\mathbf{t})} = 0 \quad (0 \neq \mathbf{t} \in \partial\mathbb{R}_+^3)$$

It follows  $D_k\chi_0^*(\mathbf{t}) = D_k D_k \chi_0(\mathbf{t})$  for  $0 \neq \mathbf{t} \in \partial\mathbb{R}_0^3 = \{(t_1, t_2, t_3) \in \mathbb{R}_+^3 : t_1 t_2 t_3 = 0\}$  and  $k = 1, 2, 3$  whence, in view of Theorem 3.5 we conclude that  $\mathbf{\Pi}^*$  with the choice  $\chi_1^* = \chi_1$  is indeed an RSD tuple.  $\square$

## 5. Affinity invariance

*Definition. 5.1.* An RSD tuple  $\mathbf{\Pi} = [\Psi_0, \Psi_1, \chi_0, \chi_1]$  (and the procedure  $\mathfrak{F}^{\mathbf{\Pi}}$ ) is *affinity invariant* if

$$\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}} f = f|_{\text{Conv}(\mathbf{P})}$$

for all admissible pairs  $[\mathbf{P}, \mathbf{U}]$  (i.e.  $\text{Conv}(\mathbf{P})$  is a non-degenerate triangle,  $\mathbf{p}_\ell - \mathbf{p}_m \parallel \mathbf{p}_n$  ( $(\ell, m, n) \in S_3$ )) and affine (linear+constant) functions  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ .



*Remark. 5.2.* (a) Affinity invariance implies range shift property.

(b) Given any non-degenerate triangle  $\mathbf{T} = \text{Conv}\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\} \subset \mathbb{R}^2$ , any affine function  $\mathbb{R}^2 \rightarrow \mathbb{R}$  is a linear combination of the weights  $\lambda_1, \lambda_2, \lambda_3$ . Hence to verify the affinity invariance of  $\mathbf{\Pi}$ , it suffices to prove the relation  $\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}} \lambda_1(\mathbf{x}) = \lambda_1(\mathbf{x})$  ( $\mathbf{x} \in \text{Conv}(\mathbf{P})$ ) for all admissible pairs  $[\mathbf{P}, \mathbf{U}]$ .

(c) In view of Remark 2.4(c) and the identities  $G_1(\mathbf{x} - \mathbf{p}_k) = \lambda(\mathbf{x}) - \delta_{1k}$  resp.  $G_i(\mathbf{p}_j - \mathbf{p}_k) = \lambda_i(\mathbf{p}_j) - \lambda_i(\mathbf{p}_k) = \delta_{ij} - \delta_{ik}$ , we can write

$$\begin{aligned} \mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}} \lambda_1 &= \sum_{k=1}^3 \left[ \lambda_1(\mathbf{p}_k) \Psi_0(\lambda_k) + G_1(\mathbf{x} - \mathbf{p}_k) \Psi_1(\lambda_k) \right] - \\ &- \sum_{(\ell, m, n) \in S_3} \frac{G_\ell \mathbf{u}_n}{G_n \mathbf{u}_n} \left[ \lambda_1(\mathbf{p}_\ell) \chi_0(\lambda_\ell, \lambda_m, \lambda_n) + G_1(\mathbf{p}_m - \mathbf{p}_\ell) \chi_1(\lambda_\ell, \lambda_m, \lambda_n) \right] = \\ &= \Psi_0(\lambda_1) + \Psi_1(\lambda_1)(\lambda_1 - 1) + \Psi_1(\lambda_2)\lambda_1 + \Psi_1(\lambda_3)\lambda_1 - \\ &- \frac{G_1 \mathbf{u}_3}{G_3 \mathbf{u}_3} \left\{ \chi_0(\lambda_1, \lambda_2, \lambda_3) - \chi_1(\lambda_1, \lambda_2, \lambda_3) \right\} - \frac{G_2 \mathbf{u}_3}{G_3 \mathbf{u}_3} \chi_1(\lambda_2, \lambda_1, \lambda_3) - \\ &- \frac{G_1 \mathbf{u}_2}{G_2 \mathbf{u}_2} \left\{ \chi_0(\lambda_1, \lambda_3, \lambda_2) - \chi_1(\lambda_1, \lambda_3, \lambda_2) \right\} - \frac{G_3 \mathbf{u}_2}{G_2 \mathbf{u}_2} \chi_1(\lambda_3, \lambda_1, \lambda_2) - 0 - 0. \end{aligned}$$

with the ordering (1,2,3), (2,1,3), (1,3,2), (3,1,2), (2,3,1), (3,2,1) for  $S_3$ .

**Lemma. 5.3.** *If the RSD tuple  $\mathbf{\Pi} = [\Psi_0, \Psi_1, \chi_0, \chi_1]$  is affinity invariant then necessarily*

$$\begin{aligned} \Psi_1(t) &= \Psi_0(t) \quad \text{and} \quad \Psi_0(t) + \Psi_0(1-t) = 1 \quad (0 \leq t \leq 1), \\ \chi_0(t_1, t_2, t_3) &= \chi_1(t_1, t_2, t_3) + \chi_1(t_2, t_1, t_3) \quad ((t_1, t_2, t_3) \in \Delta_3). \end{aligned} \tag{5.4}$$

PROOF. Since  $\mathbf{\Pi}$  is an RSD tuple, by Definition 3.2, the directional derivatives  $F'(\mathbf{x}_t^3) \mathbf{u}_3$  along the side  $[\mathbf{p}_2, \mathbf{p}_3]$  with generic point  $\mathbf{x}_t^1 = t\mathbf{p}_2 + (1-t)\mathbf{p}_3$  of the function  $F = \mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}} \lambda_1$  are the linear combinations

$$\begin{aligned} F'(\mathbf{x}_t^1) \mathbf{u}_1 &= \Psi_1(t) F'(\mathbf{p}_2) \mathbf{u}_1 + \Psi_1(1-t) F'(\mathbf{p}_3) \mathbf{u}_1 = \\ &= \Psi_1(t) G_1 \mathbf{u}_1 + \Psi_1(1-t) G_1 \mathbf{u}_1 = [\Psi_1(t) + \Psi_1(1-t)] G_1 \mathbf{u}_1. \end{aligned}$$

Due to the admissibility of  $[\mathbf{P}, \mathbf{U}]$ ,  $G_1 \mathbf{u}_1 \neq 0$ . In case of the affinity invariance of  $\mathbf{\Pi}$ ,  $F = \lambda_1$  with  $F'(\mathbf{x}_t^1) \mathbf{u}_1 = G_1 \mathbf{u}_1$  implying  $\Psi_1(t) + \Psi_1(1-t) = 1$ . Also if  $F = \mathfrak{F}^{\mathbf{\Pi}} \lambda_1 = \lambda_1$  then, as a consequence of the relations (2.7), at the points  $\mathbf{x}_t^3 = t\mathbf{p}_1 + (1-t)\mathbf{p}_2$  we have

$$\begin{aligned} t &= \lambda_1(\mathbf{x}_t^3) = F(\mathbf{x}_t^3) = \Psi_0(t) - \Psi_1(t)(1-t) + \Psi_1(1-t)t, \\ &= [\Psi_0(t) - \Psi_1(t)] + t[\Psi_1(t) + \Psi_1(1-t)] = [\Psi_0(t) - \Psi_1(t)] + t \end{aligned}$$

whence the first statement in (5.4) is immediate.

The proof of the second statement relies upon the fact that the formula for  $\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}}$  in Remark 5.2(c) must be independent of the choice of the vectors in  $\mathbf{U}$ . Using the argument of the proof of Proposition 4.4, this means that the expression

$$\begin{aligned} & \Psi_0(\lambda_1) + \Psi_1(\lambda_1)(\lambda_1 - 1) + \Psi_1(\lambda_2)\lambda_1 + \Psi(\lambda_3)\lambda_1 - \\ & - \alpha_3 \left\{ \chi_0(\lambda_1, \lambda_2, \lambda_3) - \chi_1(\lambda_1, \lambda_2, \lambda_3) \right\} - (-1 - \alpha_3)\chi_1(\lambda_2, \lambda_1, \lambda_3) - \\ & - \alpha_2 \left\{ \chi_0(\lambda_1, \lambda_3, \lambda_2) - \chi_1(\lambda_1, \lambda_3, \lambda_2) \right\} - (-1 - \alpha_2)\chi_1(\lambda_3, \lambda_1, \lambda_2) \end{aligned}$$

must be independent of the scalars  $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ . In particular it follows  $\chi_0(\lambda_1, \lambda_2, \lambda_3) - \chi_1(\lambda_1, \lambda_2, \lambda_3) - \chi_1(\lambda_3, \lambda_1, \lambda_3) = 0$  which completes the proof.  $\square$

*Remark. 5.5.* (a) According to (5.4), affinity invariant tuples are of the form

$$\mathbf{\Pi} = [\Psi, \Psi, \chi_0, \chi_1], \quad \chi_0|_{\Delta_3} = 2\chi_1^{[s]}|_{\Delta_3} \quad (5.6)$$

with a d-symmetric shape function  $\Psi$  and the symmetrization (4.13) of  $\chi_1$ .

(b) From the final argument in the proof of Lemma 5.3 it readily follows the below converse statement.

**Corollary. 5.7.** *Assume  $\mathbf{\Pi} = [\Psi, \Psi, \chi_0, \chi_1]$  is an RSD-tuple with (5.6). Then, independently of the choice of the vectors  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  we get*

$$\mathfrak{F}_{\mathbf{P}, \mathbf{U}}^{\mathbf{\Pi}} \lambda_1 = \sum_{i=1}^3 \Psi(\lambda_i)\lambda_1 + \chi_1(\lambda_2, \lambda_1, \lambda_3) + \chi_1(\lambda_3, \lambda_1, \lambda_2). \quad (5.8)$$

**Proposition. 5.9.** *An RSD tuple  $\mathbf{\Pi}$  is affinity invariant if and only if it is of the form  $\mathbf{\Pi} = [\Psi, \Psi, \chi_0, \chi_1]$  with range shift property, (5.6) and*

$$\chi_1(t_2, t_1, t_3) + \chi_1(t_3, t_1, t_2) = t_1 \sum_{(\ell, m, n) \in S_3} \chi_1(t_\ell, t_m, t_n) \quad ((t_1, t_2, t_3) \in \Delta_3). \quad (5.10)$$

**PROOF.** Necessity: Assume the affinity invariance of  $\mathbf{\Pi}$ . According to 5.5 and Corollary 5.7, we can write  $\mathbf{\Pi} = [\Psi, \Psi, \chi_0, \chi_1]$  with

$$\lambda_1 = \sum_{i=1}^3 \Psi(\lambda_i)\lambda_1 + \chi_1(\lambda_2, \lambda_1, \lambda_3) + \chi_1(\lambda_3, \lambda_1, \lambda_2). \quad (5.10)$$

On the other hand (Remark 5.2(a), Lemma 5.3)  $\mathbf{\Pi}$  has range shift property with (5.6) and  $1 = \sum_{i=1}^3 \Psi(\lambda_i) + \chi_0(\lambda_1, \lambda_2, \lambda_3) + \chi_0(\lambda_2, \lambda_3, \lambda_1) + \chi_0(\lambda_3, \lambda_1, \lambda_2)$  by (4.9). Hence, in view of (5.6) we conclude that

$$1 = \sum_{i=1}^3 \Psi(\lambda_i) + \sum_{(\ell, m, n) \in S_3} \chi_1(\lambda_\ell, \lambda_m, \lambda_n). \quad (5.11)$$

We obtain (5.10) by subtracting (5.10) from  $\lambda_1 \cdot (5.11)$ .

Sufficiency: We obtain (5.10) i.e. the affinity invariance of  $\mathbf{\Pi}$  by adding the equations  $0 = \chi_1(\lambda_2, \lambda_1, \lambda_3) + \chi_1(\lambda_3, \lambda_1, \lambda_2) - \lambda_1 \sum_{(\ell, m, n) \in S_3} \chi_1(\lambda_\ell, \lambda_m, \lambda_n)$  and (5.11) multiplied with  $\lambda_1$ .  $\square$

**Corollary. 5.12.** *A tuple  $\mathbf{\Pi} = [\Psi_0, \Psi_1, \chi_0, \chi_1]$  with range shift property and such that  $\chi_1 = \chi_0 \varphi$  for some  $\varphi \in \mathcal{C}^1(\text{dom}(\chi_0))$  is affinity invariant if for every  $(t_1, t_2, t_3) \in \Delta_3$  we have*

$$\begin{aligned} 1 &= \varphi(t_1, t_2, t_3) + \varphi(t_2, t_1, t_3), \\ \chi_0(t_3, t_1, t_2) \varphi(t_3, t_1, t_2) + \chi_0(t_2, t_1, t_3) \varphi(t_2, t_1, t_3) &= \\ &= t_1 \left[ \chi_0(t_1, t_2, t_3) + \chi_0(t_2, t_3, t_1) + \chi_0(t_2, t_3, t_1) \right]. \end{aligned} \quad 5.13$$

*Example. 5.14.* The tuple  $\mathbf{\Pi} [\Phi, \Phi, \chi_0, \chi_0 \varphi]$  is affinity invariant with

$$\Phi(t) = t^3(10 - 15t + 6t^2), \quad \chi_0(t_1, t_2, t_3) = 30t_1^2 t_2^2 t_3, \quad \varphi(t_1, t_2, t_3) = t_2 + \frac{1}{2} t_3.$$

*Proof.* The range shift property of  $\mathbf{\Pi}$  is established in Example 4.10. The identities (5.13) follow with straightforward calculation.

## References

- [1] M. BERGER 1987, *Geometry I, Springer-Verlag, New York.*
- [2] J. CAO, Z. CHEN, X. WEI AND Y.J. ZHANG, A finite element framework based on bivariate simplex splines on triagle configurations, *Comput. Methods Appl. Mech. Eng.* **357**, 112598.
- [3] S. HAHMANN AND G.-P. BONNEAU, Triangular  $G^1$  interpolation by 4-splitting domain triangles, *Comput. Aided Geom. Des.* **17**, 731–757.
- [4] I.V. SERGIENKO, O.M. LYTUVYN, O.O. LYTUVYN AND O.I. DENISOVA, Explicit formulas for interpolating splines of degree 5 on the triangle, *Cybernetics and Systems Analysis* **50**, no. 5, 670–678.

- [5] L.L. STACHÓ, A Simple Affine-Invariant Spline Interpolation over Triangular Meshes, *Mathematics* **10**, no. 5, 776.
- [6] L.L. STACHÓ, Locally Generated Polynomial C1-Splines over Triangular Meshes, *Miskolc Mathematical Notes* **23**, no. 2, 897–911.
- [7] H. WHITNEY, Analytic extensions of differentiable functions, *Transactions of the Amer. Math. Soc.* **36**, no. 1, 63–89.
- 8
- [8] M. ZLÁMAL AND Ā. ZENIŠEK, Mathematical aspects of the FEM, Technical Physical and Mathematical Principles of the FEM, *Praha, Czech Republic*, 15–39.

DEPARTMENT OF NUMERICAL AND APPLIED MATHEMATICS  
BOLYAI INSTITUTE  
UNIVERSITY OF SZEGED  
H-6720 SZEGED, ARADI VÉRTANÚK TERE 1  
HUNGARY  
*Current address:* Department of Numerical and Applied Mathematics  
Bolyai Institute  
University of Szeged  
H-6720 Szeged, Aradi Vértanúk tere 1  
Hungary

*E-mail:* [stacho@math.u-szeged.hu](mailto:stacho@math.u-szeged.hu)  
*URL:* <http://www.math.u-szeged/~stacho>