



Existence and uniqueness of damped solutions of singular IVPs with ϕ -Laplacian

Jana Burkotová¹, Irena Rachůnková¹, Martin Rohleder^{✉1} and
Jakub Stryja²

¹Department of Mathematics, Faculty of Science, Palacký University Olomouc
17. listopadu 12, 771 46 Olomouc, Czech Republic

²Department of Mathematics and Descriptive Geometry, VŠB - Technical University Ostrava
17. listopadu 15, 708 33 Ostrava, Czech Republic

Received 23 June 2016, appeared 25 December 2016

Communicated by Ivan Kiguradze

Abstract. We study analytical properties of a singular nonlinear ordinary differential equation with a ϕ -Laplacian. In particular we investigate solutions of the initial value problem

$$(p(t)\phi(u'(t)))' + p(t)f(\phi(u(t))) = 0, \quad u(0) = u_0 \in [L_0, L], \quad u'(0) = 0$$

on the half-line $[0, \infty)$. Here, f is a continuous function with three zeros $\phi(L_0) < 0 < \phi(L)$, function p is positive on $(0, \infty)$ and the problem is singular in the sense that $p(0) = 0$ and $1/p(t)$ may not be integrable on $[0, 1]$. The main goal of the paper is to prove the existence of damped solutions defined as solutions u satisfying $\sup\{u(t), t \in [0, \infty)\} < L$. Moreover, we study the uniqueness of damped solutions. Since the standard approach based on the Lipschitz property is not applicable here in general, the problem is more challenging. We also discuss the uniqueness of other types of solutions.

Keywords: second order ODE, time singularity, existence and uniqueness, ϕ -Laplacian, damped solution, half-line.

2010 Mathematics Subject Classification: 34A12, 34D05, 34C11, 34C37.

1 Introduction

We study the equation

$$(p(t)\phi(u'(t)))' + p(t)f(\phi(u(t))) = 0, \quad t \in (0, \infty) \tag{1.1}$$

with the initial conditions

$$u(0) = u_0, \quad u'(0) = 0, \quad u_0 \in [L_0, L], \tag{1.2}$$

[✉]Corresponding author. Email: martin.rohleder@upol.cz

where

$$\phi \in C^1(\mathbb{R}), \quad \phi'(x) > 0 \text{ for } x \in (\mathbb{R} \setminus \{0\}), \quad (1.3)$$

$$\phi(\mathbb{R}) = \mathbb{R}, \quad \phi(0) = 0, \quad (1.4)$$

$$L_0 < 0 < L, \quad f(\phi(L_0)) = f(0) = f(\phi(L)) = 0, \quad (1.5)$$

$$f \in C[\phi(L_0), \phi(L)], \quad xf(x) > 0 \text{ for } x \in ((\phi(L_0), \phi(L)) \setminus \{0\}), \quad (1.6)$$

$$p \in C[0, \infty) \cap C^1(0, \infty), \quad p'(t) > 0 \text{ for } t \in (0, \infty), \quad p(0) = 0. \quad (1.7)$$

A model example of (1.1), (1.2) is a problem with the α -Laplacian described below.

Example 1.1. Consider $\phi(x) = |x|^\alpha \operatorname{sgn} x$, $x \in \mathbb{R}$, where $\alpha \geq 1$. Then $\phi'(x) = \alpha|x|^{\alpha-1}$ and conditions (1.3) and (1.4) are fulfilled. If we take $p(t) = t^\beta$, $t \in [0, \infty)$, where $\beta > 0$, then p fulfils (1.7). As an example of f satisfying conditions (1.5) and (1.6) we can take $f(x) = x(x - \phi(L_0))(\phi(L) - x)$, $x \in \mathbb{R}$.

A special case of equation (1.1), which has the form

$$\left(t^{n-1}u'(t)\right)' + t^{n-1}f(u(t)) = 0, \quad t \in (0, \infty),$$

arises in many areas. For example in the study of phase transition of Van der Waals fluids [10], in population genetics, where it serves as a model for the spatial distribution of the genetic composition of a population [9], in the homogeneous nucleation theory [1], in the relativistic cosmology for description of particles which can be treated as domains in the universe [16], or in the nonlinear field theory, in particular, when describing bubbles generated by scalar fields of the Higgs type in the Minkowski spaces [7]. The equation

$$\left(p(t)u'(t)\right)' + q(t)f(u(t)) = 0, \quad t \in (0, \infty)$$

without ϕ -Laplacian, was investigated for $p \equiv q$ in [19–24] and for $p \not\equiv q$ in [5, 6, 25, 26]. Other problems without ϕ -Laplacian close to (1.1), (1.2) can be found in [2–4, 12–14] and those with ϕ -Laplacian in [8, 11, 15, 17, 18].

Definition 1.2. A function $u \in C^1[0, \infty)$ with $\phi(u') \in C^1(0, \infty)$ which satisfies equation (1.1) for every $t \in (0, \infty)$ is called a *solution* of equation (1.1). If moreover u satisfies the initial conditions (1.2), then u is called a *solution* of problem (1.1), (1.2).

Definition 1.3. Consider a solution u of problem (1.1), (1.2) with $u_0 \in (L_0, L)$ and denote

$$u_{\sup} = \sup\{u(t) : t \in [0, \infty)\}.$$

If $u_{\sup} < L$, then u is called a *damped solution* of problem (1.1), (1.2).

If $u_{\sup} = L$, then u is called a *homoclinic solution* of problem (1.1), (1.2).

The homoclinic solution is called a *regular homoclinic solution*, if $u(t) < L$ for $t \in [0, \infty)$ and a *singular homoclinic solution*, if there exists $t_0 > 0$ such that $u(t_0) = L$.

If $u_{\sup} > L$, then u is called an *escape solution* of problem (1.1), (1.2).

Remark 1.4. Equation (1.1) has the constant solutions $u(t) \equiv L$, $u(t) \equiv 0$ and $u(t) \equiv L_0$.

Our goal in this paper is to prove new existence and uniqueness results for equation (1.1) with ϕ -Laplacian. The presence of ϕ -Laplacian in equation (1.1) brings difficulties in the study of the uniqueness. For example if $\phi(x) = |x|^\alpha \operatorname{sgn} x$ and $\alpha > 1$, then ϕ fulfils the Lipschitz

condition on \mathbb{R} . On the other hand, $\phi^{-1} = |x|^{\frac{1}{\alpha}} \operatorname{sgn} x$ and $(\phi^{-1})'(x) = \frac{1}{\alpha} |x|^{\frac{1}{\alpha}-1}$. Thus we get $\lim_{x \rightarrow 0} (\phi^{-1})'(x) = \infty$ and the function ϕ^{-1} does not fulfil the Lipschitz condition in the neighbourhood of zero. Since both ϕ and ϕ^{-1} must be present in the operator form of problem (1.1), (1.2), (compare with (4.2)), we cannot use the standard approach with a Lipschitz constant to prove the uniqueness near zero. Therefore we develop a different approach near zero and show conditions which guarantee the uniqueness of damped and regular homoclinic solutions of problem (1.1), (1.2) (Theorem 5.4) and the uniqueness of escape solutions (Theorem 6.5) of the auxiliary problem (2.1), (1.2) introduced in Section 2.

We also present conditions sufficient for the existence of solutions of problem (1.1), (1.2). The existence of damped solutions of problem (1.1), (1.2) is proved here (Theorem 5.1). The more complicated questions about the existence of escape and homoclinic solutions and about nonexistence of singular homoclinic solutions remain open and they will be studied in our next paper.

2 Properties of solutions of auxiliary equation (2.1)

In this section we introduce an auxiliary equation with a bounded nonlinearity and we describe properties of its solutions. By means of these results we proceed to a priori estimates of solutions, existence and continuous dependence of solutions on initial values in next sections. The auxiliary equation has the form

$$(p(t)\phi(u'(t)))' + p(t)\tilde{f}(\phi(u(t))) = 0, \quad t \in (0, \infty), \quad (2.1)$$

where

$$\tilde{f}(x) = \begin{cases} f(x) & \text{for } x \in [\phi(L_0), \phi(L)], \\ 0 & \text{for } x < \phi(L_0), x > \phi(L). \end{cases} \quad (2.2)$$

Properties of solutions of (2.1) are derived in the next lemmas.

Lemma 2.1. *Let (1.3)–(1.7) hold and let u be a solution of equation (2.1).*

- a) *Assume that there exists $a \geq 0$ such that $u(a) \in (0, L)$ and $u'(a) = 0$. Then $u'(t) < 0$ for $t \in (a, \theta]$, where θ is the first zero of u on (a, ∞) . If such θ does not exist, then $u'(t) < 0$ for $t \in (a, \infty)$.*
- b) *Assume that there exists $b \geq 0$ such that $u(b) \in (L_0, 0)$ and $u'(b) = 0$. Then $u'(t) > 0$ for $t \in (b, \theta]$, where θ is the first zero of u on (b, ∞) . If such θ does not exist, then $u'(t) > 0$ for $t \in (b, \infty)$.*

Proof.

- a) Let $a \geq 0$ be such that $u(a) \in (0, L)$ and $u'(a) = 0$. First, we assume that there exists $\theta > a$ satisfying $u(t) > 0$ on (a, θ) and $u(\theta) = 0$. Assume that there exists $\tau \in (a, \theta)$ such that $u'(\tau) \geq 0$, $u(t) \in [u(a), L)$ for $t \in (a, \tau]$. Integrate (2.1) from a to τ and obtain

$$p(\tau)\phi(u'(\tau)) = - \int_a^\tau p(s)\tilde{f}(\phi(u(s))) ds < 0.$$

Hence, by (1.3) and (1.7), $u'(\tau) < 0$, a contradiction. Therefore $u' < 0$ on (a, θ) . Furthermore, integrating (2.1) over (a, θ) , we get

$$p(\theta)\phi(u'(\theta)) = - \int_a^\theta p(s)\tilde{f}(\phi(u(s))) ds < 0.$$

Thus, by (1.3) and (1.7), $u'(\theta) < 0$ and we have $u' < 0$ on $(a, \theta]$. If u is positive on $[a, \infty)$, we obtain as before $u' < 0$ on (a, ∞) .

b) We argue similarly as in a). □

Lemma 2.2. *Let (1.3)–(1.7) hold and let u be a solution of equation (2.1). Assume that there exists $a \geq 0$ such that $u(a) = L$ and $u'(a) = 0$.*

a) *Let $\theta > a$ be the first zero of u on (a, ∞) . Then there exists $a_1 \in [a, \theta]$ such that*

$$u(a_1) = L, \quad u'(a_1) = 0, \quad 0 \leq u(t) < L, \quad u'(t) < 0, \quad t \in (a_1, \theta].$$

b) *Let $u > 0$ on $[a, \infty)$ and $u \not\equiv L$ on $[a, \infty)$. Then there exists $a_1 \in [a, \infty)$ such that*

$$u(a_1) = L, \quad u'(a_1) = 0, \quad 0 < u(t) < L, \quad u'(t) < 0, \quad t \in (a_1, \infty).$$

In the both cases $u(t) = L$ for $t \in [a, a_1]$.

Proof.

a) Assume that there exists $t^* > a$ such that $u(t^*) > L$. Then we can find $\tau \in [a, t^*)$ satisfying

$$u(t) > L, \quad t \in (\tau, t^*], \quad u(\tau) = L. \quad (2.3)$$

Hence $u'(\tau) \geq 0$. Integrating (2.1) over $[\tau, t^*]$, we get, by (2.2),

$$p(t^*)\phi(u'(t^*)) = p(\tau)\phi(u'(\tau)) \geq 0,$$

which yields $u'(t^*) \geq 0$. Therefore $u > L$ on $[t^*, \infty)$ and u cannot have the zero θ , a contradiction. We have proved $0 < u \leq L$ on $[a, \theta]$ and deduce from (2.1) $(p(t)\phi(u'(t)))' \leq 0$ for $t \in [a, \theta]$. Consequently $u'(t) \leq 0$ and u is nonincreasing on $[a, \theta]$. Hence there exists $a_1 = [a, \theta]$ such that

$$u(a_1) = L, \quad u'(a_1) = 0, \quad 0 < u(t) < L, \quad t \in (a_1, \theta).$$

Since u is monotonous on $[a, a_1]$ then $u \equiv L$ on $[a, a_1]$. Now, we can argue as in the proof of Lemma 2.1 a) with a_1 instead of a .

b) Assume as in a) that there exists $t^* > a$ such that $u(t^*) > L$. Then we can find $\tau \in [a, t^*)$ satisfying (2.3). Hence $u'(\tau) \geq 0$. Integrate (2.1) over $[\tau, t]$, where $t \in (\tau, t^*]$. We get, by (2.2),

$$p(t)\phi(u'(t)) = p(\tau)\phi(u'(\tau)), \quad t \in (\tau, t^*].$$

If $u'(\tau) = 0$, then $u'(t) = 0$ for $t \in (\tau, t^*]$, which contradicts $u(\tau) = L$, $u(t^*) > L$. Therefore $u'(\tau) > 0$. Let $\xi \in [0, \tau)$ be the minimal number fulfilling $0 < u(t) < L$, $u'(t) > 0$, $t \in (\xi, \tau)$. Since $u(\xi) < L$, $u'(\xi) \geq 0$, we obtain $\xi > a$. Integrating (2.1) over $[a, \xi]$, we derive $u'(\xi) < 0$, a contradiction. We have proved that $0 < u \leq L$ on $[a, \infty)$, and that u is nonincreasing on (a, ∞) . If $u \not\equiv L$ on $[a, \infty)$, we can find $a_1 \geq a$ such that the assertion b) holds using the arguments from the proof of Lemma 2.1 a). Moreover, $u \equiv L$ on $[a, a_1]$. □

In order to derive further important properties of solutions of (2.1) we need to assume

$$\exists \bar{B} \in (L_0, 0) : \tilde{F}(\bar{B}) = \tilde{F}(L), \quad \text{where } \tilde{F}(x) = \int_0^x \tilde{f}(\phi(s)) \, ds, \quad x \in \mathbb{R} \quad (2.4)$$

and

$$\limsup_{t \rightarrow \infty} \frac{p'(t)}{p(t)} < \infty. \quad (2.5)$$

Remark 2.3. According to (2.4), we have $\tilde{F} \in C^1(\mathbb{R})$, $\tilde{F}(0) = 0$, \tilde{F} is positive and increasing on $[0, L]$ and positive and decreasing on $[L_0, 0]$.

Example 2.4. If p, ϕ and f are from Example 1.1 and in addition $L < |L_0|$, then conditions (2.4) and (2.5) are satisfied.

Remark 2.5. From (1.3) and (1.4), we get

$$x\phi(x) > 0 \quad \text{for } x \in (\mathbb{R} \setminus \{0\}), \quad (2.6)$$

and there exists an inverse function ϕ^{-1} , which is continuous and increasing on \mathbb{R} . By (1.7), the function p is positive and increasing on $(0, \infty)$.

Lemma 2.6. Assume that (1.3)–(1.7), (2.4) and (2.5) hold. Let u be a solution of equation (2.1) and let there exist $b \geq 0$ and $\theta > b$ such that

$$u(b) \in [\bar{B}, 0), \quad u'(b) = 0, \quad u(\theta) = 0, \quad u(t) < 0, \quad t \in [b, \theta). \quad (2.7)$$

Then there exists $a \in (\theta, \infty)$ such that

$$u'(a) = 0, \quad u'(t) > 0, \quad t \in (b, a), \quad u(a) \in (0, L).$$

Proof. Let u be a solution of equation (2.1) satisfying (2.7). Then

$$\phi'(u'(t))u''(t) + \frac{p'(t)}{p(t)}\phi(u'(t)) + \tilde{f}(\phi(u(t))) = 0, \quad t \in (0, \infty). \quad (2.8)$$

By Lemma 2.1 b) and by (2.7), we have $u'(t) > 0$ for $t \in (b, \theta]$.

Step 1. We assume that $a > \theta$ satisfying $u'(a) = 0$ does not exist. Then we get

$$u'(t) > 0, \quad t \in (b, \infty), \quad (2.9)$$

and hence u is increasing on (b, ∞) . Since $u(\theta) = 0$, the inequality

$$u(t) > 0, \quad t \in (\theta, \infty) \quad (2.10)$$

holds. Let $(\theta, A) \subset (\theta, \infty)$ be a maximal interval with the property

$$u(t) < L, \quad t \in (\theta, A). \quad (2.11)$$

Using (1.3), (1.5), (1.6) and (2.6) we obtain $\tilde{f}(\phi(u(t))) > 0$ for $t \in (\theta, A)$. Consequently, equation (2.8) yields

$$u''(t) < 0, \quad t \in (\theta, A), \quad (2.12)$$

and thus u' is decreasing on (θ, A) .

- (i) Let $A < \infty$. Then (2.11) implies $u(A) = L$. Multiplying (2.8) by u' and integrating from b to A we get

$$\int_b^A \phi'(u'(s)) u'(s) u''(s) ds + \int_b^A \frac{p'(s)}{p(s)} \phi(u'(s)) u'(s) ds + \int_b^A \tilde{f}(\phi(u(s))) u'(s) ds = 0.$$

After substitutions we derive

$$\int_{u'(b)}^{u'(A)} x \phi'(x) dx + \int_b^A \frac{p'(s)}{p(s)} \phi(u'(s)) u'(s) ds + \int_{u(b)}^{u(A)} \tilde{f}(\phi(y)) dy = 0. \quad (2.13)$$

Due to (2.7) and (2.9) $u'(b) = 0$ and $u'(A) > 0$. Therefore conditions (1.7) and (2.6) imply

$$\int_{u'(b)}^{u'(A)} x \phi'(x) dx > 0, \quad \int_b^A \frac{p'(s)}{p(s)} \phi(u'(s)) u'(s) ds > 0.$$

Using this we derive from (2.13)

$$\int_{u(b)}^{u(A)} \tilde{f}(\phi(y)) dy = \int_{u(b)}^L \tilde{f}(\phi(y)) dy < 0,$$

and hence $\tilde{F}(L) - \tilde{F}(u(b)) < 0$. By (2.4), (2.7) and Remark 2.3, we obtain

$$\tilde{F}(L) < \tilde{F}(u(b)) \leq \tilde{F}(\bar{B}) = \tilde{F}(L),$$

which is a contradiction.

- (ii) Now we assume that $A = \infty$. Inequalities (2.10) and (2.11) give

$$0 < u(t) < L, \quad t \in (\theta, \infty).$$

By (2.9) u is increasing on (θ, ∞) and

$$\lim_{t \rightarrow \infty} u(t) = \ell,$$

where $\ell \in (0, L]$. By (2.9) and (2.12) u' is decreasing and positive on (θ, ∞) and $\lim_{t \rightarrow \infty} u'(t) \geq 0$. Since ℓ is finite, we have

$$\lim_{t \rightarrow \infty} u'(t) = 0. \quad (2.14)$$

Let $\ell = L$. Similarly as before we derive

$$\int_{u'(b)}^{u'(t)} x \phi'(x) dx + \int_b^t \frac{p'(s)}{p(s)} \phi(u'(s)) u'(s) ds + \int_{u(b)}^{u(t)} \tilde{f}(\phi(y)) dy = 0, \quad t \in (b, \infty).$$

Since the first integral is positive, we have

$$\int_{u(b)}^{u(t)} \tilde{f}(\phi(y)) dy < - \int_b^t \frac{p'(s)}{p(s)} \phi(u'(s)) u'(s) ds, \quad t \in (b, \infty).$$

This yields

$$\lim_{t \rightarrow \infty} (\tilde{F}(u(t)) - \tilde{F}(u(b))) \leq - \int_b^\infty \frac{p'(s)}{p(s)} \phi(u'(s)) u'(s) ds < 0.$$

Using Remark 2.3 and conditions (2.4) and (2.7) we deduce

$$\tilde{F}(L) < \tilde{F}(u(b)) \leq \tilde{F}(\bar{B}) = \tilde{F}(L),$$

which is a contradiction.

Let $\ell \in (0, L)$. For $t \rightarrow \infty$ in (2.8) we get, by (1.4) and (2.5),

$$\phi'(0) \lim_{t \rightarrow \infty} u''(t) = -\tilde{f}(\phi(\ell)). \quad (2.15)$$

Since $-\tilde{f}(\phi(\ell)) \in (-\infty, 0)$, the inequality $\lim_{t \rightarrow \infty} u''(t) < 0$ holds, contrary to (2.14).

We have proved that there exists $a > \theta$ such that $u'(a) = 0$.

Step 2. Let $u' > 0$ on $[\theta, a)$. Then $u(a) > 0$. It remains to prove that $u(a) < L$. Multiplying (2.8) by u' and integrating from b to a we get similarly as before

$$\int_{u(b)}^{u(a)} \tilde{f}(\phi(y)) \, dy < 0, \quad t \in (b, a),$$

and

$$\tilde{F}(u(a)) < \tilde{F}(u(b)) \leq \tilde{F}(\bar{B}) = \tilde{F}(L).$$

By Remark 2.3, the inequality $u(a) < L$ holds. \square

Lemma 2.7. *Assume that (1.3)–(1.7), (2.4) and (2.5) hold. Let u be a solution of equation (2.1) and let there exist $a \geq 0$ and $\theta > a$ such that*

$$u(a) \in (0, L), \quad u'(a) = 0, \quad u(\theta) = 0, \quad u(t) > 0, \quad t \in [a, \theta). \quad (2.16)$$

Then there exists $b \in (\theta, \infty)$ such that

$$u'(b) = 0, \quad u'(t) < 0, \quad t \in (a, b), \quad u(b) \in (\bar{B}, 0).$$

Proof. We argue similarly as in the proof of Lemma 2.6. Let u be a solution of equation (2.1) satisfying (2.16). By Lemmas 2.1 a) and 2.2 a) and (2.16), we have $u'(t) < 0$, for $t \in (a, \theta]$.

Step 1. We assume that $b > \theta$ satisfying $u'(b) = 0$ does not exist. Then we get

$$u(t) < 0, \quad t \in (\theta, \infty), \quad u'(t) < 0, \quad t \in (a, \infty), \quad (2.17)$$

and hence u is decreasing on (a, ∞) . Let $(\theta, A) \subset (\theta, \infty)$ be the maximal interval with the property

$$u(t) > \bar{B}, \quad t \in (\theta, A). \quad (2.18)$$

Then

$$u''(t) > 0, \quad t \in (\theta, A) \quad (2.19)$$

and thus u' is increasing on (θ, A) .

(i) Let $A < \infty$. Then (2.18) implies $u(A) = \bar{B}$. Similarly as in the proof of Lemma 2.6 Step 1 part (i) we get the contradiction

$$\tilde{F}(\bar{B}) < \tilde{F}(u(a)) \leq \tilde{F}(L) = \tilde{F}(\bar{B}).$$

(ii) Now we assume that $A = \infty$. By (2.17) and (2.18), u is decreasing on (θ, ∞) and $\lim_{t \rightarrow \infty} u(t) = \ell \in [\bar{B}, 0)$. Due to (2.17) and (2.19) u' is increasing and negative on (θ, ∞) and $\lim_{t \rightarrow \infty} u'(t) \leq 0$. Since ℓ is finite, we have $\lim_{t \rightarrow \infty} u'(t) = 0$.

Similarly as in the proof of Lemma 2.6 Step 1 part (ii) we obtain a contradiction both for $\ell = \bar{B}$ and for $\ell \in (\bar{B}, 0)$.

We have shown that there exists $b > \theta$ such that $u'(b) = 0$.

Step 2. Let $u' < 0$ on $[\theta, b)$. Then $u(b) < 0$ and we proceed similarly as in the proof of Lemma 2.6 Step 2 and get $\tilde{F}(u(b)) < \tilde{F}(\bar{B})$. By Remark 2.3, the inequality $\bar{B} < u(b)$ holds. \square

Lemma 2.8. *Assume that (1.3)–(1.7) and (2.5) hold. Let u be a solution of equation (2.1) and let there exists $b \geq 0$ such that*

$$u(b) \in (L_0, 0), \quad u'(b) = 0, \quad u(t) < 0, \quad t \in [b, \infty).$$

Then

$$\lim_{t \rightarrow \infty} u(t) = 0, \quad \lim_{t \rightarrow \infty} u'(t) = 0.$$

Proof. By Lemma 2.1 b), $u'(t) > 0$ for $t \in (b, \infty)$. Hence u is increasing on (b, ∞) ,

$$L_0 < u(t) < 0, \quad t \in (b, \infty) \tag{2.20}$$

and

$$\lim_{t \rightarrow \infty} u(t) =: \ell \in (u(b), 0].$$

Multiplying equation (2.8) by u' and integrating it from b to t , we obtain

$$\psi_1(t) + \psi_2(t) + \psi_3(t) = 0, \quad t \in (b, \infty), \tag{2.21}$$

where

$$\psi_1(t) = \int_{u'(b)}^{u'(t)} x \phi'(x) dx, \quad \psi_2(t) = \int_b^t \frac{p'(s)}{p(s)} \phi(u'(s)) u'(s) ds, \quad \psi_3(t) = \int_{u(b)}^{u(t)} \tilde{f}(\phi(x)) dx.$$

We have $\psi_3(t) = \tilde{F}(u(t)) - \tilde{F}(u(b))$, where \tilde{F} is defined by (2.4). Since $\tilde{F}(x)$ is decreasing for $x \in (L_0, 0)$ and u is increasing on (b, ∞) , $\tilde{F}(u(t))$ is decreasing for $t \in (b, \infty)$ due to (2.20) and $\lim_{t \rightarrow \infty} \tilde{F}(u(t)) = \tilde{F}(\ell)$. Therefore

$$\lim_{t \rightarrow \infty} \psi_3(t) =: Q_3 \in (-\tilde{F}(L_0), 0).$$

The positivity of ψ_1 on (b, ∞) yields the inequality $\psi_2(t) < -\psi_3(t)$ for $t \in (b, \infty)$. Since ψ_2 is continuous, increasing and positive on (b, ∞) ,

$$\lim_{t \rightarrow \infty} \psi_2(t) =: Q_2 \in (0, -Q_3].$$

Consequently (2.21) gives

$$\lim_{t \rightarrow \infty} \psi_1(t) =: Q_1 \in [0, \tilde{F}(L_0)).$$

Therefore

$$\lim_{t \rightarrow \infty} \Phi(u'(t)) = Q_1, \quad \text{where } \Phi(z) := \int_0^z x \phi'(x) dx, \quad z > 0.$$

Φ is positive, continuous and increasing on $(0, \infty)$ and so its inverse Φ^{-1} is positive, continuous and increasing, as well. Thus

$$\lim_{t \rightarrow \infty} \Phi^{-1}(\Phi(u'(t))) = \lim_{t \rightarrow \infty} u'(t) = \Phi^{-1}(Q_1) \geq 0.$$

According to (2.20),

$$\lim_{t \rightarrow \infty} u'(t) = 0.$$

Finally, assume that $\ell \in (u(b), 0)$. Letting $t \rightarrow \infty$ in (2.8), we get, by (1.4), (2.5), that (2.15) holds. Since $-\tilde{f}(\phi(\ell)) \in (0, \infty)$, we get $\lim_{t \rightarrow \infty} u''(t) > 0$, contrary to $\lim_{t \rightarrow \infty} u'(t) = 0$. Therefore $\ell = 0$. \square

Lemma 2.9. *Assume that (1.3)–(1.7) and (2.5) hold. Let u be a solution of equation (2.1) and let there exists $a \geq 0$ such that*

$$u(a) \in (0, L], \quad u'(a) = 0, \quad u(t) > 0, \quad t \in [a, \infty).$$

Then either

$$u(t) = L, \quad t \in [a, \infty) \tag{2.22}$$

or

$$\lim_{t \rightarrow \infty} u(t) = 0, \quad \lim_{t \rightarrow \infty} u'(t) = 0.$$

Proof. Step 1. Let $u(a) \in (0, L)$. We continue analogously as in proof of Lemma 2.8. By Lemma 2.1 a), $u'(t) < 0$ for $t \in (a, \infty)$. Hence

$$0 < u(t) < L, \quad t \in (a, \infty) \tag{2.23}$$

and

$$\lim_{t \rightarrow \infty} u(t) =: \ell \in [0, u(a)).$$

Multiplying equation (2.8) by u' and integrating it from a to t , we obtain (2.21) with b replaced by a . By Remark 2.3, $\tilde{F}(x)$ is increasing for $x \in (0, L)$ and since u is decreasing on (a, ∞) , we get $\tilde{F}(u(t))$ is decreasing for $t \in (a, \infty)$ due to (2.23). Consequently $\lim_{t \rightarrow \infty} \tilde{F}(u(t)) = \tilde{F}(\ell)$. Let ψ_1, ψ_2 and ψ_3 be defined as in the proof of Lemma 2.8, where b is replaced by a . Then

$$\lim_{t \rightarrow \infty} \psi_3(t) = \lim_{t \rightarrow \infty} \tilde{F}(u(t)) - \tilde{F}(u(a)) =: Q_3 \in (-\tilde{F}(L), 0).$$

The positivity of ψ_1 on (a, ∞) yields the inequality $\psi_2(t) < -\psi_3(t)$ for $t \in (a, \infty)$. Since ψ_2 is continuous, increasing and positive on (a, ∞) , we get

$$\lim_{t \rightarrow \infty} \psi_2(t) =: Q_2 \in (0, -Q_3] \quad \text{and} \quad \lim_{t \rightarrow \infty} \psi_1(t) =: Q_1 \in [0, \tilde{F}(L)).$$

Therefore

$$\lim_{t \rightarrow \infty} \Phi(u'(t)) = Q_1, \quad \text{where} \quad \Phi(z) := \int_0^z x \phi'(x) dx, \quad z < 0.$$

Φ is positive, continuous and decreasing on $(-\infty, 0)$ and so its inverse Φ^{-1} is positive, continuous and decreasing, as well. Thus

$$\lim_{t \rightarrow \infty} \Phi^{-1}(\Phi(u'(t))) = \lim_{t \rightarrow \infty} u'(t) = \Phi^{-1}(Q_1) \geq 0.$$

According to (2.23), we have $\lim_{t \rightarrow \infty} u'(t) = 0$. Similarly as in the proof of Lemma 2.8 we derive a contradiction for $\ell \in (0, u(a))$ and get $\ell = 0$.

Step 2. Let $u(a) = L$. Assume that u does not fulfil (2.22). By Lemma 2.2 b) there exists $a_1 \geq a$ such that $0 < u(t) < L$, $u'(t) < 0$, $t \in (a_1, \infty)$, and we can use the arguments from Step 1 to prove the last assertion. \square

3 A priori estimates

In order to prove the existence and uniqueness of solutions of the auxiliary problem (2.1), (1.2) and of the original problem (1.1), (1.2), a priori estimates derived in this section are needed.

Lemma 3.1. *Assume that (1.3)–(1.7), (2.4) and (2.5) hold. Let u be a solution of problem (2.1), (1.2) with $u_0 \in (L_0, \bar{B})$. Let there exist $\theta > 0$, $a > \theta$ such that*

$$u(\theta) = 0, \quad u(t) < 0, \quad t \in [0, \theta), \quad u'(a) = 0, \quad u'(t) > 0, \quad t \in (\theta, a). \quad (3.1)$$

Then

$$u(a) \in (0, L], \quad u'(t) > 0, \quad t \in (0, a). \quad (3.2)$$

Proof. From Lemma 2.1 b) and (3.1), we have $u' > 0$ on $(0, a)$. Therefore, $u(a) > 0$. Now, assume that $u(a) > L$. Hence, there exists $a_0 \in (\theta, a)$ such that $u(t) > L$ on $(a_0, a]$. Integrating equation (2.1) over (a_0, a) and using (2.2), we get

$$p(a_0)\phi(u'(a_0)) - p(a)\phi(u'(a)) = \int_{a_0}^a p(s)\tilde{f}(\phi(u(s))) ds = 0,$$

and so $p(a_0)\phi(u'(a_0)) = 0$. Thus $u'(a_0) = 0$, contrary to $u' > 0$ on $(0, a)$. We have proved $u(a) \leq L$. \square

Lemma 3.2. *Let assumptions (1.3)–(1.7), (2.4) and (2.5) hold. Let u be a solution of problem (2.1), (1.2) with $u_0 \in (L_0, 0) \cup (0, L)$. Then*

$$u_0 \in [\bar{B}, 0) \cup (0, L) \quad \Rightarrow \quad \bar{B} < u(t) < L, \quad t \in (0, \infty), \quad (3.3)$$

$$u_0 \in (L_0, \bar{B}) \quad \Rightarrow \quad u_0 < u(t), \quad t \in (0, \infty). \quad (3.4)$$

Proof. Let $u(0) = u_0 \in (0, L)$. If $u > 0$ on $(0, \infty)$, then, by Lemma 2.1 a), $u' < 0$ on $(0, \infty)$ and (3.3) holds. Assume that there exists $\theta_1 > 0$ such that $u(\theta_1) = 0$, $u(t) > 0$ for $t \in [0, \theta_1)$. According to Lemma 2.7,

$$\exists b \in (\theta_1, \infty) : u'(b) = 0, \quad u'(t) < 0, \quad t \in (0, b), \quad u(b) = (\bar{B}, 0).$$

If $u < 0$ on (b, ∞) , then, by Lemma 2.1 b), u is increasing on (b, ∞) and (3.3) is valid. Assume that there exists $\theta_2 > b$ such that $u(\theta_2) = 0$, $u(t) < 0$ for $t \in [b, \theta_2)$. Due to Lemma 2.6,

$$\exists a \in (\theta_2, \infty) : u'(a) = 0, \quad u'(t) > 0, \quad t \in (b, a), \quad u(a) = (0, L).$$

Now we use the previous arguments replacing 0 by a .

Let $u(0) = u_0 \in [\bar{B}, 0)$. We have the same situation as before, where b is replaced by 0. So we argue similarly.

Let $u(0) = u_0 \in (L_0, \bar{B})$. If $u < 0$ on $(0, \infty)$, then, by Lemma 2.1 b), $u' > 0$ on $(0, \infty)$ and (3.4) is valid. Assume that there exists $\theta_1 > 0$ such that $u(\theta_1) = 0$, $u(t) < 0$ for $t \in [0, \theta_1)$. By Lemma 2.1 b), $u' > 0$ on $(0, \theta_1]$. If $u' > 0$ on (θ_1, ∞) , then (3.4) holds. Assume that there exists $a > \theta_1$ such that $u'(a) = 0$, $u'(t) > 0$ for $t \in (\theta_1, a)$. According to Lemma 3.1, (3.2) holds. If $u > 0$ on $[a, \infty)$, (3.4) is valid. Let there exists $\theta_2 > a$ such that $u(\theta_2) = 0$, $u > 0$ on $[a, \theta_2)$. We can apply Lemma 2.7 and argue as before. \square

Remark 3.3. According to (2.2), (3.3), (3.4) and Definition 1.3, u is a damped or a homoclinic solution of the auxiliary problem (2.1), (1.2) if and only if u is a damped or a homoclinic solution of the original problem (1.1), (1.2).

Note that the auxiliary nonlinearity is bounded due to (2.2). Therefore there exists $\tilde{M} > 0$ such that

$$|\tilde{f}(x)| \leq \tilde{M}, \quad x \in \mathbb{R}. \quad (3.5)$$

Lemma 3.4. *Assume (1.3)–(1.7). Let u be a solution of problem (2.1), (1.2) with $u_0 \in [L_0, L]$. The inequality*

$$\int_0^\beta \frac{p'(t)}{p(t)} |\phi(u'(t))| dt \leq \tilde{M}(\beta - \varphi(\beta)) \quad (3.6)$$

is valid for every $\beta > 0$. If moreover (2.4) and (2.5) hold, then there exists $\tilde{c} > 0$ such that

$$|u'(t)| \leq \tilde{c}, \quad t \in [0, \infty), \quad (3.7)$$

for every solution u of (2.1), (1.2) with $u_0 \in (L_0, 0) \cup (0, L)$.

Proof. Step 1. Let u be solution of (2.1), (1.2) with $u_0 \in [L_0, L]$. Integrating equation (2.1) over $(0, t)$, $t > 0$, and using (3.5), we have

$$|\phi(u'(t))| = \left| -\frac{1}{p(t)} \int_0^t p(\tau) \tilde{f}(\phi(u(\tau))) d\tau \right| \leq \frac{\tilde{M}}{p(t)} \int_0^t p(\tau) d\tau$$

and

$$\frac{p'(t)}{p(t)} |\phi(u'(t))| \leq \tilde{M} \frac{p'(t)}{p^2(t)} \int_0^t p(\tau) d\tau.$$

Choose a $\beta > 0$. Integrating this inequality by parts from 0 to β , we get (3.6).

Step 2. Assume moreover that (2.4) and (2.5) hold. Denote

$$\Psi_1(z) := \int_0^z x\phi'(x) dx; \quad \Psi_2(z) := \int_0^z x\phi'(-x) dx; \quad z \in [0, \infty).$$

Clearly, Ψ_1, Ψ_2 are positive, continuous and increasing on $(0, \infty)$. Put

$$\tilde{c} = \max \left\{ \Psi_1^{-1}(\tilde{F}(L_0)), \Psi_2^{-1}(\tilde{F}(L)) \right\}, \quad (3.8)$$

where \tilde{F} is defined in (2.4).

Let $u(0) = u_0 \in (L_0, 0)$, $u'(0) = 0$ and let u be a solution of equation (2.1). Then (2.8) holds.

- (i) Assume that $u < 0$ on $[0, \infty)$. By Lemma 2.1 b) $u' > 0$ on $(0, \infty)$, and by Lemma 2.8 $\lim_{t \rightarrow \infty} u'(t) = 0$. Therefore there exists $\xi \in (0, \infty)$ such that

$$\max_{t \in [0, \infty)} |u'(t)| = u'(\xi) > 0, \quad u(\xi) \in (u_0, 0). \quad (3.9)$$

Multiplying (2.8) by u' and integrating over $[0, \xi]$ we get

$$\int_{u'(0)}^{u'(\xi)} x\phi'(x) dx + \int_0^\xi \frac{p'(t)}{p(t)} \phi(u'(t)) u'(t) dt + \int_{u(0)}^{u(\xi)} \tilde{f}(\phi(x)) dx = 0. \quad (3.10)$$

Since the second integral in (3.10) is positive, (3.9) and (3.10) yield

$$\Psi_1(u'(\xi)) < \tilde{F}(u_0) - \tilde{F}(u(\xi)) < \tilde{F}(u_0) < \tilde{F}(L_0).$$

Therefore

$$0 < u'(\xi) < \Psi_1^{-1}(\tilde{F}(L_0)). \quad (3.11)$$

Due to (3.8) and (3.9) estimate (3.7) is proved.

- (ii) Assume that $\theta \in (0, \infty)$ is such that $u < 0$ on $[0, \theta)$, $u(\theta) = 0$. Then by Lemma 2.1 b), $u' > 0$ on $(0, \theta]$. Let $a > \theta$ be such that $u' > 0$ on (θ, a) , $u'(a) = 0$. On interval (θ, a) we have $u > 0$, $u' > 0$ and by (1.3), (1.6), (1.7), (2.6) and (2.8) we get $u'' < 0$ on $[\theta, a)$. Therefore u' is decreasing on $[\theta, a)$ and there exists $\zeta \in (0, \theta)$ such that

$$\max_{t \in [0, a)} |u'(t)| = u'(\zeta) > 0, \quad u(\zeta) \in (u_0, 0). \quad (3.12)$$

Analogously as in part (i) we get (3.11) and if $a = \infty$ then estimate (3.7) is proved.

- (iii) Assume that $a < \infty$ in (3.12). We have $u'(a) = 0$ and by Lemma 2.6 and Lemma 3.1 we deduce that $u(a) \in (0, L]$. Let $u > 0$ on $[a, \infty)$. Then Lemma 2.9 gives $\lim_{t \rightarrow \infty} u'(t) = 0$ and hence there exists $\eta \in (a, \infty)$ such that

$$\max_{t \in [a, \infty)} |u'(t)| = -u'(\eta) > 0, \quad u(\eta) \in (0, u(a)). \quad (3.13)$$

Multiplying (2.8) by u' and integrating over $[a, \eta]$ we get

$$\int_{u'(a)}^{|u'(\eta)|} x\phi'(-x) dx + \int_a^\eta \frac{p'(t)}{p(t)} \phi(u'(t)) u'(t) dt + \int_{u(a)}^{u(\eta)} \tilde{f}(\phi(x)) dx = 0.$$

Since the second integral in (3.10) is positive, (3.9) and (3.10) yield

$$\Psi_2(|u'(\eta)|) < \tilde{F}(u(a)) - \tilde{F}(u(\eta)) < \tilde{F}(L).$$

Then

$$0 < |u'(\eta)| < \Psi_2^{-1}(\tilde{F}(L)). \quad (3.14)$$

Using (3.11), (3.12), (3.13) and (3.14) we obtain (3.7) due to (3.8).

- (iv) Assume, that there exists $\chi \in (a, \infty)$ which is the next zero of u . Summarized, we have $u(a) \in (0, L]$, $u'(a) = 0$, $u(\chi) = 0$, $u > 0$ on $[a, \chi)$. By Lemma 2.7 there exists $b \in (\chi, \infty)$ such that $u'(b) = 0$, $u' < 0$ on (a, b) , $u(b) \in (\bar{B}, 0)$ and by (2.8) we have $u'' > 0$ on $[\chi, b)$. Consequently there exists $\eta \in (a, \chi)$ such that

$$\max_{t \in [a, b]} |u'(t)| = -u'(\eta) > 0, \quad u(\eta) \in (0, u(a)).$$

Similarly as in part (iii) we get (3.14) and (3.7).

- (v) Since $u(b) < 0$ we continue repeating the argument of parts (i)–(iii) with b on place of 0 and the arguments of part (iv) writing \tilde{b} instead of b . After finite or infinite number of steps we obtain (3.7).

If $u_0 \in (0, L)$, we can argue similarly. □

4 Existence and continuous dependence of solutions on initial values

This section is devoted to the existence of solutions of the auxiliary problem (2.1), (1.2) which is proved in Theorem 4.1 by means of the Schauder fixed point theorem. Moreover, the question about continuous dependence of solutions on initial values is discussed in Theorems 4.3, 4.6, 4.8. Theorem 4.3 provides also a uniqueness result for special kinds of ϕ -Laplacians.

For the following investigation, we introduce a function φ

$$\varphi(t) := \frac{1}{p(t)} \int_0^t p(s) ds, \quad t \in (0, \infty), \quad \varphi(0) = 0.$$

This function is continuous on $[0, \infty)$ and satisfies

$$0 < \varphi(t) \leq t, \quad t \in (0, \infty), \quad \lim_{t \rightarrow 0^+} \varphi(t) = 0. \quad (4.1)$$

Theorem 4.1 (Existence of solutions of problem (2.1), (1.2)). *Assume (1.3)–(1.7). Then, for each $u_0 \in [L_0, L]$, there exists a solution u of problem (2.1), (1.2).*

Proof. Clearly, for $u_0 = L_0$, $u_0 = 0$ and $u_0 = L$ there exists a solution by Remark 1.4. Assume that $u_0 \in (L_0, 0) \cup (0, L)$. Integrating equation (2.1), we get the equivalent form of problem (2.1), (1.2)

$$u(t) = u_0 + \int_0^t \phi^{-1} \left(-\frac{1}{p(s)} \int_0^s p(\tau) \tilde{f}(\phi(u(\tau))) d\tau \right) ds, \quad t \in [0, \infty). \quad (4.2)$$

Choose a $\beta > 0$, consider the Banach space $C[0, \beta]$ with the maximum norm and define an operator $\mathcal{F}: C[0, \beta] \rightarrow C[0, \beta]$,

$$(\mathcal{F}u)(t) = u_0 + \int_0^t \phi^{-1} \left(-\frac{1}{p(s)} \int_0^s p(\tau) \tilde{f}(\phi(u(\tau))) d\tau \right) ds.$$

Put $\Lambda = \max\{|L_0|, L\}$ and consider the ball $\mathcal{B}(0, R) = \{u \in C[0, \beta] : \|u\|_{C[0, \beta]} \leq R\}$, where $R = \Lambda + \beta\phi^{-1}(\tilde{M}\beta)$ and \tilde{M} is from (3.5). Since ϕ is increasing on \mathbb{R} , ϕ^{-1} is also increasing on \mathbb{R} and, by (4.1), $\phi^{-1}(\tilde{M}\varphi(t)) \leq \phi^{-1}(\tilde{M}\beta)$, $t \in [0, \beta]$. The norm of $\mathcal{F}u$ can be estimated as follows

$$\begin{aligned} \|\mathcal{F}u\|_{C[0, \beta]} &= \max_{t \in [0, \beta]} \left| u_0 + \int_0^t \phi^{-1} \left(-\frac{1}{p(s)} \int_0^s p(\tau) \tilde{f}(\phi(u(\tau))) d\tau \right) ds \right| \\ &\leq \Lambda + \int_0^t \left| \phi^{-1}(\tilde{M}\varphi(s)) \right| ds \leq \Lambda + \int_0^t \phi^{-1}(\tilde{M}\beta) ds \leq \Lambda + \beta\phi^{-1}(\tilde{M}\beta) = R, \end{aligned}$$

which yields that \mathcal{F} maps $\mathcal{B}(0, R)$ on itself.

Let us prove that \mathcal{F} is compact on $\mathcal{B}(0, R)$. Choose a sequence $\{u_n\} \subset C[0, \beta]$ such that $\lim_{n \rightarrow \infty} \|u_n - u\|_{C[0, \beta]} = 0$. We have

$$\begin{aligned} (\mathcal{F}u_n)(t) - (\mathcal{F}u)(t) &= \int_0^t \left(\phi^{-1} \left(-\frac{1}{p(s)} \int_0^s p(\tau) \tilde{f}(\phi(u_n(\tau))) d\tau \right) \right. \\ &\quad \left. - \phi^{-1} \left(-\frac{1}{p(s)} \int_0^s p(\tau) \tilde{f}(\phi(u(\tau))) d\tau \right) \right) ds. \end{aligned}$$

Since $\tilde{f}(\phi)$ is continuous on $[0, \beta]$, we get

$$\lim_{n \rightarrow \infty} \|\tilde{f}(\phi(u_n)) - \tilde{f}(\phi(u))\|_{C[0, \beta]} = 0.$$

Put

$$\begin{aligned} A_n(t) &= -\frac{1}{p(t)} \int_0^t p(\tau) \tilde{f}(\phi(u_n(\tau))) d\tau, \\ A(t) &= -\frac{1}{p(t)} \int_0^t p(\tau) \tilde{f}(\phi(u(\tau))) d\tau, \quad t \in (0, \beta], \quad A_n(0) = A(0) = 0, \quad n \in \mathbb{N}. \end{aligned}$$

Then, for a fixed $n \in \mathbb{N}$,

$$|A_n(t) - A(t)| = \left| \frac{1}{p(t)} \int_0^t p(\tau) (\tilde{f}(\phi(u(\tau))) - \tilde{f}(\phi(u_n(\tau)))) d\tau \right|, \quad t \in (0, \beta]$$

and, by (4.1) and (3.5), $\lim_{t \rightarrow 0^+} |A_n(t) - A(t)| = 0$. Therefore $A_n - A \in C[0, \beta]$ and

$$\|A_n - A\|_{C[0, \beta]} \leq \|\tilde{f}(\phi(u_n)) - \tilde{f}(\phi(u))\|_{C[0, \beta]} \beta, \quad n \in \mathbb{N}.$$

This implies that $\lim_{n \rightarrow \infty} \|A_n - A\|_{C[0, \beta]} = 0$. Using the continuity of ϕ^{-1} on \mathbb{R} , we have

$$\lim_{n \rightarrow \infty} \left\| \phi^{-1}(A_n) - \phi^{-1}(A) \right\|_{C[0, \beta]} = 0.$$

Therefore

$$\begin{aligned} \lim_{n \rightarrow \infty} \|\mathcal{F}u_n - \mathcal{F}u\|_{C[0, \beta]} &= \lim_{n \rightarrow \infty} \left\| \int_0^t (\phi^{-1}(A_n(s)) - \phi^{-1}(A(s))) ds \right\|_{C[0, \beta]} \\ &\leq \beta \lim_{n \rightarrow \infty} \left\| \phi^{-1}(A_n) - \phi^{-1}(A) \right\|_{C[0, \beta]} = 0, \end{aligned}$$

that is the operator \mathcal{F} is continuous.

Choose an arbitrary $\varepsilon > 0$ and put $\delta = \frac{\varepsilon}{\phi^{-1}(\tilde{M}\beta)}$. Then, for $t_1, t_2 \in [0, \beta]$ and $u \in \mathcal{B}(0, R)$,

$$\begin{aligned} |t_1 - t_2| < \delta &\Rightarrow |(\mathcal{F}u)(t_1) - (\mathcal{F}u)(t_2)| = \left| \int_{t_2}^{t_1} \phi^{-1} \left(-\frac{1}{p(s)} \int_0^s p(\tau) \tilde{f}(\phi(u(\tau))) d\tau \right) ds \right| \\ &\leq \left| \int_{t_2}^{t_1} \phi^{-1}(\tilde{M}\varphi(s)) ds \right| \leq \left| \int_{t_2}^{t_1} \phi^{-1}(\tilde{M}\beta) ds \right| = \phi^{-1}(\tilde{M}\beta) |t_1 - t_2| < \phi^{-1}(\tilde{M}\beta) \delta = \varepsilon. \end{aligned}$$

Hence, functions in $\mathcal{F}(\mathcal{B}(0, R))$ are equicontinuous, and, by the Arzelà–Ascoli theorem, the set $\mathcal{F}(\mathcal{B}(0, R))$ is relatively compact. Consequently, the operator \mathcal{F} is compact on $\mathcal{B}(0, R)$.

The Schauder fixed point theorem yields a fixed point u^* of \mathcal{F} in $\mathcal{B}(0, R)$. Therefore,

$$u^*(t) = u_0 + \int_0^t \phi^{-1} \left(-\frac{1}{p(s)} \int_0^s p(\tau) \tilde{f}(\phi(u^*(\tau))) d\tau \right) ds.$$

Hence, $u^*(0) = u_0$,

$$(p(t)\phi((u^*)'(t)))' = -p(t)\tilde{f}(\phi(u^*(t))), \quad t \in [0, \beta].$$

Further,

$$|(u^*)'(t)| = \left| \phi^{-1} \left(-\frac{1}{p(t)} \int_0^t p(s) \tilde{f}(\phi(u^*(s))) ds \right) \right| \leq \phi^{-1}(\tilde{M}\varphi(t)), \quad t \in [0, \beta].$$

Thus, by (4.1), $\lim_{t \rightarrow 0^+} \phi^{-1}(\tilde{M}\varphi(t)) = \phi^{-1}(0) = 0$ and therefore $\lim_{t \rightarrow 0^+} (u^*)'(t) = 0 = (u^*)'(0)$. According to (2.2), $\tilde{f}(\phi(u^*(t)))$ is bounded on $[0, \infty)$ and hence u^* can be extended to interval $[0, \infty)$ as a solution of equation (2.1). This classical extension result follows from more general Theorem 11.5 in [13]. \square

Example 4.2. Consider $\phi: \mathbb{R} \rightarrow \mathbb{R}$ given by one of the next formulas

$$\phi(x) = |x|^\alpha \operatorname{sgn} x, \quad \alpha > 1, \quad (4.3)$$

$$\phi(x) = (x^4 + 2x^2) \operatorname{sgn} x, \quad (4.4)$$

$$\phi(x) = \sinh x = \frac{e^x - e^{-x}}{2}, \quad (4.5)$$

$$\phi(x) = \arg \sinh x = \ln(x + \sqrt{x^2 + 1}), \quad (4.6)$$

$$\phi(x) = \ln(|x| + 1) \operatorname{sgn} x, \quad (4.7)$$

$$\phi(x) = (|x| + 1)^\alpha - 1) \operatorname{sgn} x, \quad \alpha \in (0, 1). \quad (4.8)$$

Assume that $\phi(L) < -\phi(L_0)$ and put

$$p(t) = t^\beta, \quad t \in [0, \infty), \quad \beta > 0,$$

$$f(x) = k|x|^\gamma \operatorname{sgn} x(x - \phi(L_0))(\phi(L) - x), \quad x \in [\phi(L_0), \phi(L)], \quad \gamma > 0, \quad k > 0.$$

Then the functions p , ϕ and f fulfil all assumptions of Theorem 4.1. In particular $\phi \in \operatorname{Lip}_{\text{loc}}(\mathbb{R})$ for each ϕ given by (4.3)–(4.8). Therefore the auxiliary problem (2.1), (1.2) has a solution for every $u_0 \in [L_0, L]$.

Further we examine the uniqueness of solutions of the auxiliary problem (2.1), (1.2). Our arguments are based on a continuous dependence on initial values expressed in Theorem 4.3, Theorem 4.6 and Theorem 4.8. Assumption (1.3) implies that $\phi \in \operatorname{Lip}_{\text{loc}}(\mathbb{R})$. This need not be true for ϕ^{-1} as we have shown in Introduction for $\phi(x) = |x|^\alpha \operatorname{sgn} x$, $\alpha > 1$. The special case when both ϕ and ϕ^{-1} are locally Lipschitz continuous is discussed in the next theorem.

Theorem 4.3 (Uniqueness and continuous dependence on initial values I). *Assume (1.3)–(1.7) and*

$$f \in \operatorname{Lip}[\phi(L_0), \phi(L)], \quad (4.9)$$

$$\phi^{-1} \in \operatorname{Lip}_{\text{loc}}(\mathbb{R}). \quad (4.10)$$

Let u_i be a solution of problem (2.1), (1.2) with $u_0 = B_i \in [L_0, L]$, $i = 1, 2$. Then, for each $\beta > 0$, there exists $K > 0$ such that

$$\|u_1 - u_2\|_{C^1[0, \beta]} \leq K|B_1 - B_2|. \quad (4.11)$$

Furthermore, any solution of problem (2.1), (1.2) with $u_0 \in [L_0, L]$ is unique on $[0, \infty)$.

Proof. Let $i \in 1, 2$ and let u_i be a solution of problem (2.1), (1.2) with $u_0 = B_i$. By integrating (2.1) over $[0, t]$, we obtain

$$\phi(u_i'(t)) = A_i(t), \quad u_i(t) = B_i + \int_0^t \phi^{-1}(A_i(s)) \, ds, \quad t \in [0, \infty), \quad (4.12)$$

where

$$A_i(s) = -\frac{1}{p(s)} \int_0^s p(\tau) \tilde{f}(\phi(u_i(\tau))) \, d\tau, \quad s \in [0, \infty).$$

Choose $\beta > 0$. Since $u_i, \phi(u_i') \in C[0, \beta]$, there exist $m, M \in \mathbb{R}$ such that

$$m \leq u_i(t) \leq M, \quad m \leq \phi(u_i'(t)) \leq M, \quad t \in [0, \beta], \quad i = 1, 2.$$

According to (1.3), (4.9) and (4.10) there exist positive constants $\Lambda_f, \Lambda_\phi, \Lambda_{\phi^{-1}}$ satisfying

$$\begin{aligned} |f(x_1) - f(x_2)| &\leq \Lambda_f |x_1 - x_2|, \quad x_1, x_2 \in [\phi(L_0), \phi(L)], \\ |\phi(x_1) - \phi(x_2)| &\leq \Lambda_\phi |x_1 - x_2|, \quad x_1, x_2 \in [m, M], \\ |\phi^{-1}(x_1) - \phi^{-1}(x_2)| &\leq \Lambda_{\phi^{-1}} |x_1 - x_2|, \quad x_1, x_2 \in [m, M]. \end{aligned}$$

Denote $\rho(t) := \max\{|u_1(s) - u_2(s)| : s \in [0, t]\}$, $t \in [0, \beta]$. Then, by (4.1),

$$\begin{aligned} |A_1(s) - A_2(s)| &\leq \frac{1}{p(s)} \int_0^s p(\tau) |\tilde{f}(\phi(u_1(\tau))) - \tilde{f}(\phi(u_2(\tau)))| \, d\tau \\ &\leq \Lambda_f \Lambda_\phi \frac{1}{p(s)} \int_0^s p(\tau) |u_1(\tau) - u_2(\tau)| \, d\tau \leq \Lambda_f \Lambda_\phi \rho(s) \beta, \end{aligned}$$

and by virtue of (4.12)

$$\begin{aligned} \rho(t) &\leq |B_1 - B_2| + \int_0^t |\phi^{-1}(A_1(s)) - \phi^{-1}(A_2(s))| \, ds \leq |B_1 - B_2| + \Lambda_{\phi^{-1}} \int_0^t |A_1(s) - A_2(s)| \, ds \\ &\leq |B_1 - B_2| + \Lambda_f \Lambda_\phi \Lambda_{\phi^{-1}} \beta \int_0^t \rho(s) \, ds, \quad t \in [0, \beta]. \end{aligned}$$

The Gronwall lemma yields

$$\rho(t) \leq |B_1 - B_2| e^{L\beta^2}, \quad t \in [0, \beta], \quad (4.13)$$

where $L := \Lambda_f \Lambda_\phi \Lambda_{\phi^{-1}}$. Similarly, from (4.12) it follows

$$|u'_1(t) - u'_2(t)| \leq |\phi^{-1}(A_1(t)) - \phi^{-1}(A_2(t))| \leq \Lambda_{\phi^{-1}} |A_1(t) - A_2(t)| \leq L\rho(t)\beta, \quad t \in [0, \beta].$$

Applying (4.13), we get

$$\max\{|u'_1(t) - u'_2(t)| : t \in [0, \beta]\} \leq |B_1 - B_2| L\beta e^{L\beta^2}.$$

Consequently,

$$\|u_1 - u_2\|_{C^1[0, \beta]} \leq |B_1 - B_2| (1 + L\beta) e^{L\beta^2},$$

that is (4.11) holds for

$$K := (1 + L\beta) e^{L\beta^2}.$$

Clearly, if $B_1 = B_2$, we have $u_1 = u_2$ on each $[0, \beta] \subset \mathbb{R}$ and the uniqueness for problem (2.1), (1.2) on $[0, \infty)$ follows. \square

Remark 4.4. If also (2.4) and (2.5) are fulfilled, we can use (3.7) and get universal estimates for $\phi(u'_i)$ and u_i . This is the case that K in (4.11) does not depend on a choice of u_1, u_2 .

Example 4.5. In order to apply Theorem 4.3 we need both ϕ and ϕ^{-1} from $\text{Lip}_{\text{loc}}(\mathbb{R})$. Let us check the functions ϕ in Example 4.2 from this point of view:

$$\begin{aligned} \phi(x) &= |x|^\alpha \operatorname{sgn} x, \quad \alpha > 1 && \Rightarrow \phi^{-1}(x) = |x|^{\frac{1}{\alpha}} \operatorname{sgn} x && \notin \text{Lip}_{\text{loc}}(\mathbb{R}), \\ \phi(x) &= (x^4 + 2x^2) \operatorname{sgn} x, && \Rightarrow \phi^{-1}(x) = \sqrt{\sqrt{|x| + 1} - 1} && \notin \text{Lip}_{\text{loc}}(\mathbb{R}), \\ \phi(x) &= \sinh x = \frac{e^x - e^{-x}}{2}, && \Rightarrow \phi^{-1}(x) = \operatorname{arg} \sinh x && \in \text{Lip}_{\text{loc}}(\mathbb{R}), \\ \phi(x) &= \operatorname{arg} \sinh x = \ln(x + \sqrt{x^2 + 1}) && \Rightarrow \phi^{-1}(x) = \sinh x && \in \text{Lip}_{\text{loc}}(\mathbb{R}), \\ \phi(x) &= \ln(|x| + 1) \operatorname{sgn} x && \Rightarrow \phi^{-1}(x) = (e^{|x|} - 1) \operatorname{sgn} x && \in \text{Lip}_{\text{loc}}(\mathbb{R}), \\ \phi(x) &= ((|x| + 1)^\alpha - 1) \operatorname{sgn} x, \quad \alpha \in (0, 1) && \Rightarrow \phi^{-1}(x) = ((|x| + 1)^{\frac{1}{\alpha}} - 1) \operatorname{sgn} x && \in \text{Lip}_{\text{loc}}(\mathbb{R}). \end{aligned}$$

Consider p, f from Example 4.2 with $\gamma \geq 1$ and ϕ given by one of the formulas (4.5)–(4.8). Then all assumptions of Theorem 4.3 are fulfilled and problem (2.1), (1.2) has a unique solution for $u_0 \in [L_0, L]$. Note that if $\gamma \in (0, 1)$, then f is not Lipschitz continuous on a neighbourhood of zero, that is (4.9) is not valid. Similarly, in the case that ϕ is given by (4.3) or (4.4), then ϕ^{-1} is not Lipschitz continuous on a neighbourhood of zero and hence (4.10) falls.

In the next two theorems we show assumptions under which solutions of problem (2.1), (1.2) continuously depend on their initial values in the case that ϕ^{-1} is not locally Lipschitz continuous.

Theorem 4.6 (Continuous dependence on initial values II). *Assume (1.3)–(1.7), (2.4), (2.5), (4.9) and*

$$\limsup_{x \rightarrow 0^-} \left(-x \left(\phi^{-1} \right)' (x) \right) < \infty, \quad \phi' \text{ is nonincreasing on } (-\infty, 0). \quad (4.14)$$

Let B_1, B_2 satisfy

$$B_1 \in (2\varepsilon, L - 2\varepsilon), \quad |B_1 - B_2| < \varepsilon$$

for some $\varepsilon > 0$. Let u_i be a solution of problem (2.1), (1.2) with $u_0 = B_i$, $i = 1, 2$. Then for each $\beta > 0$ where

$$u_i' < 0 \quad \text{on } (0, \beta], \quad i = 1, 2,$$

there exists $K \in (0, \infty)$ such that

$$\|u_1 - u_2\|_{C^1[0, \beta]} \leq K|B_1 - B_2|.$$

Proof. Let u_i be a solution of problem (2.1), (1.2) with $u_0 = B_i$, $i = 1, 2$. Then by integrating (2.1) over $[0, t]$, we obtain

$$\phi(u_i'(t)) = -\frac{1}{p(t)} \int_0^t p(s) \tilde{f}(\phi(u_i(s))) \, ds =: A_i(t), \quad t \in [0, \infty) \quad (4.15)$$

$$u_i(t) = B_i + \int_0^t \phi^{-1}(A_i(s)) \, ds, \quad t \in [0, \infty), \quad i = 1, 2.$$

Therefore

$$|u_1(t) - u_2(t)| \leq |B_1 - B_2| + \int_0^t \left| \phi^{-1}(A_1(s)) - \phi^{-1}(A_2(s)) \right| \, ds, \quad t \in [0, \infty). \quad (4.16)$$

In order to reach the required estimate, we restrict our consideration on a small interval $[0, \delta]$ for a suitably chosen $\delta > 0$ in Step 1. Then we prolongate the result on $[0, \beta]$ in Step 2.

Step 1. Assumptions (1.3)–(1.6), (4.9), (4.14) yield the existence of positive constants $\Lambda_f, \Lambda_\phi, K_1, K_2$ such that

$$\begin{aligned} |f(y_1) - f(y_2)| &\leq \Lambda_f |y_1 - y_2|, \quad y_1, y_2 \in [\phi(L_0), \phi(L)], \\ |\phi(x_1) - \phi(x_2)| &\leq \Lambda_\phi |x_1 - x_2|, \quad x_1, x_2 \in [L_0, L], \\ K_1 &= \min \{ f(\phi(x)) : x \in [B_1 - 2\varepsilon, B_1 + 2\varepsilon] \}, \end{aligned} \quad (4.17)$$

$$0 < -x \left(\phi^{-1} \right)' (x) \leq K_2, \quad x \in [-1, 0). \quad (4.18)$$

By Lemma 3.4, there exists $\tilde{c} > 0$ such that $|u_i'| \leq \tilde{c}$ on $[0, \infty)$, $i = 1, 2$. Let us choose δ such that

$$0 < \delta \leq \min \left\{ \frac{\varepsilon}{\tilde{c}}, \frac{1}{K_1}, \frac{K_1}{2K_2\Lambda_f\Lambda_\phi} \right\}. \quad (4.19)$$

Then we get

$$|B_1 - u_1(t)| = |u_1(0) - u_1(t)| \leq \tilde{c}\delta \leq \varepsilon,$$

which yields $u_1(t) \in [B_1 - \varepsilon, B_1 + \varepsilon]$ for $t \in [0, \delta]$. Moreover,

$$|B_1 - u_2(t)| \leq |B_1 - B_2| + |u_2(0) - u_2(t)| \leq \varepsilon + \delta\tilde{c} \leq 2\varepsilon,$$

thus $u_2(t) \in [B_1 - 2\varepsilon, B_1 + 2\varepsilon]$ holds for $t \in [0, \delta]$. Consequently, $\tilde{f}(\phi(u_i)(t)) \geq K_1$, for $t \in [0, \delta]$, $i = 1, 2$. Therefore

$$A_i(s) = - \int_0^s \frac{p(\tau)}{p(s)} \tilde{f}(\phi(u_i(\tau))) \, d\tau \leq -K_1 \int_0^s \frac{p(\tau)}{p(s)} \, d\tau, \quad s \in [0, \delta],$$

$$|A_1(s) - A_2(s)| \leq \int_0^s \frac{p(\tau)}{p(s)} |\tilde{f}(\phi(u_1(\tau))) - \tilde{f}(\phi(u_2(\tau)))| \, d\tau \leq \Lambda_f \Lambda_\phi \|u_1 - u_2\|_{C[0, \delta]} \int_0^s \frac{p(\tau)}{p(s)} \, d\tau.$$

Let $s \in (0, \delta]$ be fixed. By the mean value theorem there exists $A^*(s)$ between $A_1(s)$ and $A_2(s)$ such that

$$\left| \phi^{-1}(A_1(s)) - \phi^{-1}(A_2(s)) \right| \leq \left(\phi^{-1} \right)' (A^*(s)) |A_1(s) - A_2(s)|.$$

Since $(\phi^{-1})'$ is a nondecreasing function on $(-\infty, 0)$, we get

$$\begin{aligned} \left| \phi^{-1}(A_1(s)) - \phi^{-1}(A_2(s)) \right| &\leq \left(\phi^{-1} \right)' \left(-K_1 \int_0^s \frac{p(\tau)}{p(s)} \, d\tau \right) |A_1(s) - A_2(s)| \\ &\leq \left(\phi^{-1} \right)' \left(-K_1 \int_0^s \frac{p(\tau)}{p(s)} \, d\tau \right) \frac{\Lambda_f \Lambda_\phi \|u_1 - u_2\|_{C[0, \delta]}}{K_1} K_1 \int_0^s \frac{p(\tau)}{p(s)} \, d\tau. \end{aligned}$$

Using the monotonicity of p and (4.19), we have

$$0 < K_1 \int_0^s \frac{p(\tau)}{p(s)} \, d\tau \leq K_1 \delta \leq 1,$$

and hence, by (4.18), we get

$$\left| \phi^{-1}(A_1(s)) - \phi^{-1}(A_2(s)) \right| \leq \frac{K_2}{K_1} \Lambda_f \Lambda_\phi \|u_1 - u_2\|_{C[0, \delta]}.$$

Consequently, by (4.19), we derive from (4.16) for $t \in [0, \delta]$

$$\begin{aligned} |u_1(t) - u_2(t)| &\leq |B_1 - B_2| + \int_0^t \frac{K_2}{K_1} \Lambda_f \Lambda_\phi \|u_1 - u_2\|_{C[0, \delta]} \, ds \\ &\leq |B_1 - B_2| + \delta \frac{K_2}{K_1} \Lambda_f \Lambda_\phi \|u_1 - u_2\|_{C[0, \delta]} \leq |B_1 - B_2| + \frac{1}{2} \|u_1 - u_2\|_{C[0, \delta]}. \end{aligned}$$

This yields

$$\|u_1 - u_2\|_{C[0, \delta]} \leq 2|B_1 - B_2|. \quad (4.20)$$

Furthermore, by (4.15),

$$|u_1'(t) - u_2'(t)| = \left| \phi^{-1}(A_1(t)) - \phi^{-1}(A_2(t)) \right| \leq \frac{K_2}{K_1} \Lambda_f \Lambda_\phi \|u_1 - u_2\|_{C[0, \delta]}, \quad t \in [0, \delta].$$

Hence

$$\|u_1' - u_2'\|_{C[0, \delta]} \leq K_3 |B_1 - B_2|, \quad (4.21)$$

where $K_3 := 2\frac{K_2}{K_1}\Lambda_f\Lambda_\phi$. Finally,

$$\|u_1 - u_2\|_{C^1[0,\delta]} \leq K_{S1}|B_1 - B_2|, \quad \text{where } K_{S1} := K_3 + 2.$$

Step 2. In this step, we extend the continuous dependence on initial values from $[0, \delta]$ to $[0, \beta]$, where $u'_i(t) < 0$ for $t \in (0, \beta]$, $i = 1, 2$. To this aim, choose $i \in \{1, 2\}$ and denote

$$v_i = \max\{u'_i(t) : t \in [\delta, \beta]\} < 0, \quad m_1 = \max\{v_1, v_2\}, \quad m = \min\{-\tilde{c}, L_0\}.$$

Moreover, (1.3) yields the existence of positive Lipschitz constants $\Lambda_m, \Lambda_{\phi^{-1}}$ such that

$$\begin{aligned} |\phi(x_1) - \phi(x_2)| &\leq \Lambda_m|x_1 - x_2|, \quad x_1, x_2 \in [m, L], \\ |\phi^{-1}(y_1) - \phi^{-1}(y_2)| &\leq \Lambda_{\phi^{-1}}|y_1 - y_2|, \quad y_1, y_2 \in [\phi(-\tilde{c}), \phi(m_1)]. \end{aligned}$$

By integrating (2.1) over $[\delta, t]$, $t \in [\delta, \beta]$, we get

$$\phi(u'_i(t)) = \frac{p(\delta)}{p(t)}\phi(u'_i(\delta)) - \frac{1}{p(t)} \int_\delta^t p(s)\tilde{f}(\phi(u_i(s))) ds.$$

Let us denote

$$\begin{aligned} \tilde{A}_i(t) &:= - \int_\delta^t \frac{p(s)}{p(t)}\tilde{f}(\phi(u_i(s))) ds, \\ x_i(t) &:= \frac{p(\delta)}{p(t)}\phi(u'_i(\delta)) + \tilde{A}_i(t) = \phi(u'_i(t)), \quad t \in [\delta, \beta]. \end{aligned}$$

Then

$$u'_i(t) = \phi^{-1}(x_i(t)), \quad t \in [\delta, \beta]. \quad (4.22)$$

Since $-\tilde{c} \leq u'_i(t) \leq m_1$, then $x_i(t) \in [\phi(-\tilde{c}), \phi(m_1)]$, for $t \in [\delta, \beta]$. Integrating (4.22) from δ to t , $t \in [\delta, \beta]$, we get

$$u_i(t) = u_i(\delta) + \int_\delta^t \phi^{-1}(x_i(s)) ds.$$

By (4.20) we obtain for $t \in [\delta, \beta]$

$$\begin{aligned} |u_1(t) - u_2(t)| &\leq |u_1(\delta) - u_2(\delta)| + \int_\delta^t |\phi^{-1}(x_1(s)) - \phi^{-1}(x_2(s))| ds \\ &\leq 2|B_1 - B_2| + \Lambda_{\phi^{-1}} \int_\delta^t |x_1(s) - x_2(s)| ds. \end{aligned}$$

Further by (1.7), (4.21) we obtain for $s \in [\delta, \beta]$

$$\begin{aligned} |x_1(s) - x_2(s)| &\leq \frac{p(\delta)}{p(s)}|\phi(u'_1(\delta)) - \phi(u'_2(\delta))| + |\tilde{A}_1(s) - \tilde{A}_2(s)| \\ &\leq \Lambda_m|u'_1(\delta) - u'_2(\delta)| + \int_\delta^s |\tilde{f}(\phi(u_1(\tau))) - \tilde{f}(\phi(u_2(\tau)))| d\tau \\ &\leq \Lambda_m K_3|B_1 - B_2| + \Lambda_f \Lambda_m \int_\delta^s |u_1(\tau) - u_2(\tau)| d\tau. \end{aligned}$$

Therefore,

$$\begin{aligned} |u_1(t) - u_2(t)| &\leq 2|B_1 - B_2| + \Lambda_{\phi^{-1}}\Lambda_m K_3 \beta |B_1 - B_2| + \Lambda_{\phi^{-1}}\Lambda_f \Lambda_m \int_\delta^t \int_\delta^s |u_1(\tau) - u_2(\tau)| d\tau ds \\ &\leq K_4|B_1 - B_2| + K_5 \int_\delta^t |u_1(\tau) - u_2(\tau)| d\tau, \quad t \in [\delta, \beta], \end{aligned}$$

where $K_4 = 2 + \Lambda_{\phi^{-1}}\Lambda_m K_3\beta$, $K_5 = \Lambda_{\phi^{-1}}\Lambda_f\Lambda_m\beta$. Next we set for $t \in (\delta, \beta]$

$$\rho(t) = \max\{|u_1(s) - u_2(s)| : s \in [\delta, t]\}.$$

Then

$$\rho(t) \leq K_4|B_1 - B_2| + K_5 \int_{\delta}^t \rho(\tau) d\tau.$$

The Gronwall lemma yields that

$$\begin{aligned} \rho(t) &\leq K_4|B_1 - B_2|e^{K_5\beta}, \quad t \in [\delta, \beta] \\ \|u_2 - u_1\|_{C[\delta, \beta]} &\leq K_6|B_1 - B_2|, \quad \text{where } K_6 = K_4e^{K_5\beta}. \end{aligned}$$

By (4.22),

$$\begin{aligned} |u_1'(t) - u_2'(t)| &\leq \left| \phi^{-1}(x_1(t)) - \phi^{-1}(x_2(t)) \right| \leq \Lambda_{\phi^{-1}}|x_1(t) - x_2(t)| \\ &\leq \Lambda_{\phi^{-1}}\Lambda_m K_3|B_1 - B_2| + \Lambda_{\phi^{-1}}\Lambda_f\Lambda_m\beta \|u_1 - u_2\|_{C[\delta, \beta]} \leq K_7|B_1 - B_2|, \end{aligned}$$

where $K_7 = \Lambda_{\phi^{-1}}\Lambda_m K_3 + \Lambda_{\phi^{-1}}\Lambda_f\Lambda_m\beta K_6$. Hence

$$\begin{aligned} \|u_1' - u_2'\|_{C[\delta, \beta]} &\leq K_7|B_1 - B_2|, \\ \|u_1 - u_2\|_{C^1[\delta, \beta]} &\leq K_{S2}|B_1 - B_2| \end{aligned}$$

with $K_{S2} = K_6 + K_7$. Finally, there exists $K = K_{S1} + K_{S2}$ such that

$$\|u_1 - u_2\|_{C^1[0, \beta]} \leq K|B_1 - B_2|.$$

This completes the proof. \square

Remark 4.7. The approach developed in the proof of Theorem 4.6 cannot be used for $B_1 = L$ because then the positive constant K_1 in (4.17) which is crucial in the proof does not exist.

Theorem 4.8 (Continuous dependence on initial values III). *Assume (1.3)–(1.7), (2.4), (2.5), (4.9) and*

$$\limsup_{x \rightarrow 0^+} \left(x \left(\phi^{-1} \right)'(x) \right) < \infty, \quad \phi' \text{ is nondecreasing on } (0, \infty). \quad (4.23)$$

Let B_1, B_2 satisfy

$$B_1 \in (L_0 + 2\varepsilon, -2\varepsilon), \quad |B_1 - B_2| < \varepsilon$$

for some $\varepsilon > 0$. Let u_i be a solution of problem (2.1), (1.2) with $u_0 = B_i$, $i = 1, 2$. Then for each $\beta > 0$ where

$$u_i' > 0 \quad \text{on } (0, \beta], \quad i = 1, 2,$$

there exists $K \in (0, \infty)$ such that

$$\|u_1 - u_2\|_{C^1[0, \beta]} \leq K|B_1 - B_2|.$$

Proof. We proceed similarly as in the proof of Theorem 4.6. In Step 1 we replace $f(\phi(x))$ by $|f(\phi(x))|$ in (4.17) and the interval $[-1, 0]$ by $(0, 1]$ in (4.18). Then we derive the inequalities

$$\begin{aligned} -\tilde{f}(\phi(u_i)(t)) &= |\tilde{f}(\phi(u_i)(t))| \geq K_1, \quad t \in [0, \delta], \quad i = 1, 2, \\ A_i(s) &= - \int_0^s \frac{p(\tau)}{p(s)} \tilde{f}(\phi(u_i(\tau))) d\tau \geq K_1 \int_0^s \frac{p(\tau)}{p(s)} d\tau, \quad s \in [0, \delta]. \end{aligned}$$

Since $(\phi^{-1})'$ is nonincreasing on $(0, \infty)$, we get

$$\begin{aligned} \left| \phi^{-1}(A_1(s)) - \phi^{-1}(A_2(s)) \right| &\leq (\phi^{-1})' \left(K_1 \int_0^s \frac{p(\tau)}{p(s)} d\tau \right) |A_1(s) - A_2(s)| \\ &\leq (\phi^{-1})' \left(K_1 \int_0^s \frac{p(\tau)}{p(s)} d\tau \right) \frac{\Lambda_f \Lambda_\phi \|u_1 - u_2\|_{C[0,\delta]}}{K_1} K_1 \int_0^s \frac{p(\tau)}{p(s)} d\tau, \end{aligned}$$

and follow Step 1 in the proof of Theorem 4.6. In Step 2 having $u'_i(t) > 0$ for $t \in (0, \beta]$, $i = 1, 2$, we denote

$$v_i = \min\{u'_i(t) : t \in [\delta, \beta]\} > 0, \quad m_0 = \min\{v_1, v_2\}, \quad M = \max\{\tilde{c}, L\}.$$

By (1.3) there exists positive Lipschitz constants $\Lambda_m, \Lambda_{\phi^{-1}}$ such that

$$\begin{aligned} |\phi(x_1) - \phi(x_2)| &\leq \Lambda_m |x_1 - x_2|, \quad x_1, x_2 \in [L_0, M], \\ |\phi^{-1}(y_1) - \phi^{-1}(y_2)| &\leq \Lambda_{\phi^{-1}} |y_1 - y_2|, \quad y_1, y_2 \in [\phi(m_0), \phi(\tilde{c})]. \end{aligned}$$

We derive (4.22) and since $m_0 \leq u'_i(t) \leq \tilde{c}$, we get $x_i(t) \in [\phi(m_0), \phi(\tilde{c})]$, for $t \in [\delta, \beta]$, $i = 1, 2$. Further we argue as in the proof of Theorem 4.6. \square

5 Existence and uniqueness of damped solutions of problem (1.1), (1.2)

Main results of the present paper are formulated in this section. The existence of damped solutions is proved in Theorem 5.1 and the uniqueness is derived in Theorem 5.4. Both results hold not only for the auxiliary problem (2.1), (1.2), but above all for the original problem (1.1), (1.2). Due to Remark 3.3, immediately from Theorem 4.1 and Lemma 3.2, we obtain the existence result.

Theorem 5.1 (Existence of damped solutions of problem (1.1), (1.2)). *Assume (1.3)–(1.7), (2.5) and*

$$\exists \bar{B} \in (L_0, 0) : \int_0^{\bar{B}} f(\phi(z)) dz = \int_0^L f(\phi(z)) dz.$$

Then, for each $u_0 \in [\bar{B}, L)$, problem (1.1), (1.2) has a solution. Every solution of problem (1.1), (1.2) with $u_0 \in [\bar{B}, L)$ is damped.

Example 5.2. Problem (1.1), (1.2) with p, f and ϕ from Example 4.2 has for each $u_0 \in [\bar{B}, L)$ a damped solution.

Remark 5.3. By Theorem 5.1, we can get homoclinic or escape solutions only if $u_0 \in (L_0, \bar{B})$.

If $\phi^{-1} \notin \text{Lip}_{\text{loc}}(\mathbb{R})$, we derive results about the uniqueness by means of Theorems 4.6 and 4.8. Since the next uniqueness result concerns damped solutions, it can be formulated directly for the original problem (1.1), (1.2) due to Remark 3.3.

Theorem 5.4 (Uniqueness of damped solutions). *Assume (1.3)–(1.7), (2.4), (2.5), (4.9), (4.14) and (4.23). Let u be a damped solution of problem (1.1), (1.2) with $u_0 \in (L_0, 0) \cup (0, L)$. Then u is a unique solution of this problem.*

Proof. Assume that u is a damped solution of the auxiliary problem (2.1), (1.2) and that there exists another solution v of problem (2.1), (1.2). Definition 1.3 yields

$$u(t) < L, \quad t \in [0, \infty). \quad (5.1)$$

By Lemma 3.2, we have

$$L_0 < u(t), \quad L_0 < v(t), \quad t \in (0, \infty]. \quad (5.2)$$

Step 1. Let $u_0 \in (L_0, 0)$.

- (i) According to Lemma 2.1 b), there exists $\beta > 0$ such that $u'(t) > 0, v'(t) > 0$ for $t \in (0, \beta]$. Put

$$a = \sup\{\beta > 0 : u'(t) > 0, v'(t) > 0, t \in (0, \beta]\},$$

$$\rho(t) = u(t) - v(t), \quad t \in [0, \infty).$$

Since $u' > 0, v' > 0$ on $(0, a)$ and $B_1 := u_0 = v(0) =: B_2$, Theorem 4.8 yields

$$\rho(t) = 0, \quad t \in [0, a). \quad (5.3)$$

If $a = \infty$, then

$$u(t) = v(t), \quad t \in [0, \infty). \quad (5.4)$$

Consequently, by (5.1) and (5.2), u is a unique solution of problem (1.1), (1.2).

Let $a < \infty$. Since $u, v \in C^1[0, \infty)$, we get, by (5.3),

$$\lim_{t \rightarrow a^-} \rho(t) = \rho(a) = u(a) - v(a) = 0, \quad \lim_{t \rightarrow a^-} \rho'(t) = \rho'(a) = u'(a) - v'(a) = 0. \quad (5.5)$$

Therefore $u'(a) = v'(a)$.

- (ii) According to the definition of number a , we have $u'(a) = v'(a) = 0$. By (5.1) and Lemma 2.6 or Lemma 3.1, $u(a) = v(a) \in (0, L)$. Due to Lemma 2.1 a), there exists $\gamma > a$ such that $u'(t) < 0, v'(t) < 0$, for $t \in (a, \gamma]$. Put

$$b = \sup\{\gamma > a : u'(t) < 0, v'(t) < 0, t \in (a, \gamma]\}.$$

Since $u' < 0, v' < 0$ on (a, b) and $u(a) = v(a) \in (0, L)$, by Theorem 4.6 (working with $a, \gamma, u(a)$ and $v(a)$ instead of $0, \beta, B_1$ and B_2 respectively), we get

$$\rho(t) = 0, \quad t \in [a, b). \quad (5.6)$$

If $b = \infty$, then (5.4) holds and, by (5.1), (5.2), u is a unique solution of problem (1.1), (1.2).

Let $b < \infty$. Since $u, v \in C^1[0, \infty)$, (5.6) yields

$$\lim_{t \rightarrow b^-} \rho(t) = \rho(b) = u(b) - v(b) = 0, \quad \lim_{t \rightarrow b^-} \rho'(t) = \rho'(b) = u'(b) - v'(b) = 0.$$

Hence $u'(b) = v'(b)$ and, due to the definition of b , $u'(b) = v'(b) = 0$. Lemma 2.7 implies $u(b) = v(b) \in (\bar{B}, 0)$. Repeating the arguments in parts (i) and (ii), we get that u is a unique solution of problem (1.1), (1.2).

Step 2. Let $u_0 \in (0, L)$. We have the same situation as in part (ii) of Step 1, where a is replaced by 0, and so we argue similarly. \square

6 Uniqueness of regular homoclinic and escape solutions

In this section we discuss homoclinic and escape solutions and hence, by Remark 5.3, we take $u_0 \in (L_0, \bar{B})$. Results concerning homoclinic solutions can be formulated directly for the original problem (1.1), (1.2) due to Remark 3.3.

Theorem 6.1 (Nonexistence of singular homoclinic solutions). *Assume (1.3)–(1.7), (4.9) and (4.10). Then each homoclinic solution of problem (1.1), (1.2) with $u_0 \in (L_0, \bar{B})$ is regular.*

Proof. Let u be a singular homoclinic solution of problem (2.1), (1.2) with $u_0 \in (L_0, \bar{B})$. Then, by Definition 1.3, there exists $t_0 > 0$ such that

$$u(t_0) = L, \quad u'(t_0) = 0, \quad (6.1)$$

and

$$u(t) < L, \quad t \in [0, t_0]. \quad (6.2)$$

Using the substitution $s = t_0 - t$, $q(s) = p(t)$, $v(s) = u(t)$ for $t \in [\frac{t_0}{2}, t_0]$, we transform the terminal value problem (2.1), (6.1) on $[\frac{t_0}{2}, t_0]$ to the initial value problem

$$-(q(s)\phi(-v'(s)))' + q(s)\tilde{f}(\phi(v(s))) = 0, \quad s \in \left[0, \frac{t_0}{2}\right], \quad v(0) = L, \quad v'(0) = 0.$$

By Theorem 4.3, the only possible function satisfying this problem is the constant function $v(s) = L$ for $s \in [0, \frac{t_0}{2}]$. Therefore $u(t) = L$ for $t \in [\frac{t_0}{2}, t_0]$, which contradicts (6.2). \square

Theorem 6.1 discusses the case where $\phi^{-1} \in \text{Lip}_{\text{loc}}(\mathbb{R})$. Now we will study the case where condition (4.10) falls, that is $\phi^{-1} \notin \text{Lip}_{\text{loc}}(\mathbb{R})$. Then both regular and singular homoclinic solutions may exist and, according to Remark 4.7, we are able to prove the uniqueness just for regular ones.

Lemma 6.2 (Regular homoclinic solution is increasing). *Assume (1.3)–(1.7), (2.4), (2.5). Let u be a regular homoclinic solution of problem (1.1), (1.2) with $u_0 \in (L_0, \bar{B})$. Then*

$$\lim_{t \rightarrow \infty} u(t) = L, \quad u'(t) > 0, \quad t \in (0, \infty). \quad (6.3)$$

Moreover,

$$\lim_{t \rightarrow \infty} u'(t) = 0. \quad (6.4)$$

Proof. Let u be a regular homoclinic solution of problem (2.1), (1.2) with $u_0 \in (L_0, \bar{B})$. Thus, by Definition 1.3, $u_{\text{sup}} = L$.

Step 1. By Lemma 2.1 b) there exists $\theta_0 > 0$ such that $u(\theta_0) = 0$, $u(t) < 0$ for $t \in (0, \theta_0)$ and $u'(t) > 0$ for $t \in (0, \theta_0]$. Assume on contrary with (6.3), that $a_1 > \theta_0$ is the first zero of u' . Since u is a regular homoclinic solution, $u(a_1)$ belongs to $(0, L)$. If $u > 0$ on $[a_1, \infty)$, then by Lemma 2.1 a), u is decreasing which contradicts $u_{\text{sup}} = L$. Therefore, there exists $\theta_1 > a_1$ such that $u(\theta_1) = 0$, $u'(t) < 0$ for $t \in (a_1, \theta_1]$. Hence we have

$$u(a_1) \in (0, L), \quad u'(a_1) = 0, \quad u'(t) > 0, \quad t \in (0, a_1). \quad (6.5)$$

By Lemma 2.7 there exists $b_1 > \theta_1$ such that

$$u(b_1) \in (\bar{B}, 0), \quad u'(b_1) = 0, \quad u'(t) < 0, \quad t \in [\theta_1, b_1].$$

Since, $u_{\text{sup}} = L$, there exists $\theta_2 > b_1$, such that $u(\theta_2) = 0$, $u'(t) > 0$ for $t \in (b_1, \theta_2]$. By Lemma 2.6 there exists $a_2 > \theta_2$ such that

$$u(a_2) \in (0, L), \quad u'(a_2) = 0, \quad u'(t) > 0, \quad t \in (b_1, a_2).$$

Repeating this procedure, we obtain a sequence of zeros $\{\theta_n\}_{n=0}^{\infty}$ of u and a sequence of local maxima $\{u(a_n)\}_{n=1}^{\infty}$ of u . Now, we prove that the sequence $\{u(a_n)\}_{n=1}^{\infty}$ is nonincreasing. Choose $n \in \mathbb{N}$. Multiplying equation (1.1) by u'/p , then integrating from a_n to a_{n+1} we obtain

$$\int_{a_n}^{a_{n+1}} \phi'(u'(t))u''(t)u'(t) dt + \int_{a_n}^{a_{n+1}} \frac{p'(t)}{p(t)}\phi(u'(t))u'(t) dt + \int_{a_n}^{a_{n+1}} \tilde{f}(\phi(u(t)))u'(t) dt = 0.$$

The first integral is equal zero since $u'(a_n) = u'(a_{n+1}) = 0$. The second integral is nonnegative due to (1.7) and (2.6). Therefore,

$$0 \geq \int_{a_n}^{a_{n+1}} \tilde{f}(\phi(u(t)))u'(t) dt = \int_{u(a_n)}^{u(a_{n+1})} \tilde{f}(\phi(y)) dy = \tilde{F}(u(a_{n+1})) - \tilde{F}(u(a_n)).$$

Since \tilde{F} is increasing function, we get $u(a_n) \geq u(a_{n+1})$. The sequence $\{u(a_n)\}_{n=1}^{\infty}$ is nonincreasing, because n is chosen arbitrarily. Thus $u_{\text{sup}} < L$, which cannot be fulfilled because u is a homoclinic solution. This contradiction yields that

$$u'(t) > 0, \quad t \in (0, \infty).$$

Since $u_{\text{sup}} = L$, then $\lim_{t \rightarrow \infty} u(t) = L$.

Step 2. Since, $u > 0$ on (θ_0, ∞) , we have $f(\phi(u)) > 0$ on (θ_0, ∞) . From (1.1) we obtain that

$$0 > (p(t)\phi(u'(t)))' = p'(t)\phi(u'(t)) + p(t)(\phi(u'(t)))', \quad t \in (\theta_0, \infty).$$

Since p, p', u' and $\phi(u')$ are positive on $(0, \infty)$, we get that $\phi(u')$ is decreasing on (θ_0, ∞) . On the other hand ϕ is an increasing function. Therefore u' is a decreasing function on (θ_0, ∞) . Since $u' > 0$ on $(0, \infty)$, there exists a nonnegative limit

$$\lim_{t \rightarrow \infty} u'(t) =: K \geq 0.$$

If $K > 0$, then

$$K(t - \theta_0) \leq \int_{\theta_0}^t u'(s) ds = u(t) - u(\theta_0) = u(t).$$

As t tends to infinity the limit yields,

$$L = \lim_{t \rightarrow \infty} u(t) \geq \lim_{t \rightarrow \infty} K(t - \theta_0) = \infty,$$

a contradiction. Therefore (6.4) holds. \square

Since assumptions (1.6) are imposed to f on the interval $[\phi(L_0), \phi(L)]$ and we have no information about a behaviour of f out of this interval, we formulate results concerning escape solutions for the auxiliary problem (2.1), (1.2).

Lemma 6.3 (Escape solution is increasing). *Assume that (1.3)–(1.7), (2.4) and (2.5) hold. Let u be an escape solution of problem (2.1), (1.2) with $u_0 \in (L_0, \bar{B})$. Then*

$$u'(t) > 0, \quad t \in (0, \infty).$$

Proof. Let u be an escape solution of problem (2.1), (1.2) with $u_0 \in (L_0, \bar{B})$. Thus, by Definition 1.3, $u_{\text{sup}} > L$. Then there exists a point $c \in (0, \infty)$ such that $u(c) = L$, $u'(c) \geq 0$ and $u(t) < L$ for $t \in [0, c)$. First we exclude the case $u'(c) = 0$. Lemma 2.2 yields that if $u'(c) = 0$ then either u has a zero point $u(\theta) = 0$, $u(t) \leq L$, $t \in [c, \theta]$ or u is positive and nonincreasing on $[c, \infty)$. The later case is in contradiction with u being an escape solution. Therefore, such zero point $\theta > c$ must exist. Applying Lemma 2.1 a), b) and Lemma 2.7 and repeating the arguments as in Step 1 in the proof of Lemma 6.2, we get that u has a nonincreasing sequence $\{u(a_n)\}_{n=1}^{\infty}$ of its local maxima. Thus $u(t) \leq L$ for $t \geq 0$ on contrary that u is an escape solution. Therefore $u'(c) > 0$.

Let $c_1 > c$ be such that $u'(c_1) = 0$ and $u(t) > L$, $u'(t) > 0$ for $t \in (c, c_1)$. Integrating (2.1) over $[c, c_1]$ we get, due to (1.3), (1.4), (1.7) and (2.2),

$$\phi(u'(c_1)) = \frac{p(c)\phi(u'(c))}{p(c_1)} > 0,$$

contrary to $u'(c_1) = 0$. We have proved $u'(t) > 0$ for $t > c$.

Further, we prove that $u'(t) > 0$ for $t \in (0, \theta_0]$. Since $u_0 \in (L_0, 0)$, Lemma 2.1 b) yields that there exists $\theta_0 > 0$ such that $u(\theta_0) = 0$, $u(t) < 0$ for $t \in (0, \theta_0)$, $u'(t) > 0$ for $t \in (0, \theta_0]$.

It remains to prove that $u'(t) > 0$ for $t \in (\theta_0, c)$. Assume on the contrary that there exists $a_1 \in (\theta_0, c)$ such that (6.5) holds. We derive a contradiction as in Step 1 in the proof of Lemma 6.2. To summarize, $u'(t) > 0$ for $t > 0$. \square

We are ready to prove uniqueness results for regular homoclinic and escape solutions.

Theorem 6.4 (Uniqueness of regular homoclinic solutions). *Assume (1.3)–(1.7), (2.4), (2.5), (4.9), (4.14) and (4.23). Let u be a regular homoclinic solution of problem (1.1), (1.2) with $u_0 \in (L_0, \bar{B})$. Then u is a unique solution of this problem.*

Proof. Let u be a regular homoclinic solution of problem (2.1), (1.2). According to Lemma 6.2, $u' > 0$ on $(0, \infty)$. Consider that v is another solution of problem (2.1), (1.2). Assume that there exists $c \in (0, \infty)$ such that $v'(c) = 0$. By Lemma 2.1 b), there exists $\theta > 0$ such that $v(\theta) = 0$, $v'(t) > 0$ for $t \in (0, \theta]$. Therefore $c > \theta$ and there exists $a \in (\theta, c]$ such that $v'(a) = 0$, $v'(t) > 0$ for $t \in (0, a)$. Put

$$\rho(t) = u(t) - v(t), \quad t \in [0, \infty).$$

Since $u' > 0$, $v' > 0$ on $(0, a)$, Theorem 4.8, where $u_0 = B_1 = B_2$, gives

$$\rho(t) = 0, \quad \rho'(t) = 0, \quad t \in [0, a). \quad (6.6)$$

Since $u, v \in C^1[0, \infty)$, we get that (5.5) holds. Thus $u'(a) = v'(a)$. According to the definition of number a , we have $u'(a) = v'(a) = 0$, which contradicts the inequality $u' > 0$ on $(0, \infty)$. Therefore $a = \infty$ and, by (6.6), u is a unique solution of problem (2.1), (1.2). \square

Theorem 6.5 (Uniqueness of escape solutions). *Assume (1.3)–(1.7), (2.4), (2.5), (4.9) and (4.23). Let u be an escape solution of problem (2.1), (1.2) with $u_0 \in (L_0, \bar{B})$. Then u is a unique solution of this problem.*

Proof. Let u be an escape solution of problem (2.1), (1.2). By Lemma 6.3, $u' > 0$ on $(0, \infty)$ and we can argue as in the proof of Theorem 6.4. \square

Example 6.6. Consider p, f from Example 4.2 with $\gamma \geq 1$ and ϕ given by (4.3). Then

$$\begin{aligned} \phi^{-1}(x) &= |x|^{\frac{1}{\alpha}} \operatorname{sgn} x, & (\phi^{-1})'(x) &= \frac{1}{\alpha} |x|^{\frac{1}{\alpha}-1}, & \lim_{x \rightarrow 0} x (\phi^{-1})'(x) &= \frac{1}{\alpha} \lim_{x \rightarrow 0} x |x|^{\frac{1}{\alpha}-1} = 0 \in \mathbb{R}, \\ \phi'(x) &= \alpha |x|^{\alpha-1}, & \phi''(x) &= \frac{\alpha(\alpha-1)|x|^{\alpha-1}}{x} \begin{cases} \leq 0 & \text{for } x < 0, \\ \geq 0 & \text{for } x > 0. \end{cases} \end{aligned}$$

Hence ϕ' is nonincreasing on $(-\infty, 0)$, nondecreasing on $(0, \infty)$ and conditions (4.14) and (4.23) hold. If ϕ is given by (4.4), then

$$\begin{aligned} \phi^{-1}(x) &= \sqrt{\sqrt{|x|+1}-1}, & (\phi^{-1})'(x) &= \frac{\operatorname{sgn} x}{4\sqrt{\sqrt{|x|+1}-1}\sqrt{|x|+1}}, \\ \lim_{x \rightarrow 0} x (\phi^{-1})'(x) &= \lim_{x \rightarrow 0} \frac{|x|}{4\sqrt{\sqrt{|x|+1}-1}\sqrt{|x|+1}} = 0 \in \mathbb{R}, \\ \phi'(x) &= 4(x^3+x)\operatorname{sgn} x, & \phi''(x) &= 4(3x^2+1)\operatorname{sgn} x \begin{cases} < 0 & \text{for } x < 0, \\ > 0 & \text{for } x > 0. \end{cases} \end{aligned}$$

Therefore ϕ' is decreasing on $(-\infty, 0)$ and increasing on $(0, \infty)$. Function ϕ satisfies conditions (1.3), (1.4), (4.14) and (4.23). In both cases all assumptions of Theorem 4.1, Theorem 5.4 and Theorem 6.5 are fulfilled. Therefore problem (2.1), (1.2) has for $u_0 \in [L_0, L]$ a solution u . If $u_0 \in (L_0, 0) \cup (0, L)$ and $u < L$ on $[0, \infty)$, then u is a solution of the original problem (1.1), (1.2) and it is a unique solution of this problem. If $u_0 \in (L_0, \bar{B})$ and u is an escape solution of problem (2.1), (1.2), then u is a unique solution of this problem.

Remark 6.7. Theorem 6.1 does not cover equations having a ϕ -Laplacian in the form (4.3) or (4.4) because such ϕ -Laplacian does not fulfil the condition $\phi^{-1} \in \operatorname{Lip}_{\text{loc}}(\mathbb{R})$. Therefore to find conditions which guarantee that singular homoclinic solutions do not exist while $\phi^{-1} \notin \operatorname{Lip}_{\text{loc}}(\mathbb{R})$ is an open problem and we plan to solve it in our next paper where we also will discuss the existence and asymptotic properties of regular homoclinic and escape solutions.

Acknowledgement

The first three authors gratefully acknowledge support received from the grant No. 14-06958S of the Grant Agency of the Czech Republic.

References

- [1] F. F. ABRAHAM, *Homogeneous nucleation theory*, Acad. Press, New York 1974.
- [2] R. P. AGARWAL, D. O'REGAN, *Infinite interval problems for differential, difference and integral equations*, Kluwer, Dordrecht 2001. [MR1845855](#)
- [3] H. BERESTYCKI, P. L. LIONS, L. A. PELETIER, An ODE approach to the existence of positive solutions for semilinear problems in R^n , *Indiana Math. Univ. J.* **30**(1981), 141–157. [MR0600039](#)

- [4] D. BONHEURE, J. M. GOMES, L. SANCHEZ, Positive solutions of a second-order singular ordinary differential equation, *Nonlinear Analysis* **61**(2005), 1383–1399. [MR2135816](#)
- [5] J. BURKOTOVÁ, M. HUBNER, I. RACHŮNKOVÁ, E. B. WEINMÜLLER, Asymptotic properties of Kneser solutions to nonlinear second order ODEs with regularly varying coefficients, *J. Appl. Math. Comp.* **274**(2015), 65–82. [MR3433115](#)
- [6] J. BURKOTOVÁ, M. ROHLER, J. STRYJA, On the existence and properties of three types of solutions of singular IVPs, *Electron. J. Qual. Theory Differ. Equ.* **2015**, No. 29, 1–25. [MR3353200](#); [url](#)
- [7] G. H. DERRICK, Comments on nonlinear wave equations as models for elementary particles, *J. Math. Physics* **5**(1965), 1252–1254. [MR0174304](#)
- [8] Z. DOŠLÁ, M. MARINI, S. MATUCCI, A boundary value problem on a half-line for differential equations with indefinite weight, *Commun. Appl. Anal.* **15**(2011), 341–352. [MR2867356](#)
- [9] P. C. FIFE, *Mathematical aspects of reacting and diffusing systems*, Lecture Notes in Biomathematics, Vol. 28, Springer-Verlag, Berlin–New York, 1979. [MR0527914](#)
- [10] H. GOUIN, G. ROTOLI, An analytical approximation of density profile and surface tension of microscopic bubbles for Van der Waals fluids, *Mech. Research Commun.* **24**(1997), 255–260. [url](#)
- [11] J. JAROŠ, T. KUSANO, J.V. MANOJLOVIĆ, Asymptotic analysis of positive solutions of generalized Emden–Fowler differential equations in the framework of regular variation, *Cent. Eur. J. Math.* **11**(2013), No. 12, 2215–2233. [MR3111718](#)
- [12] I. T. KIGURADZE, *Some singular boundary value problems for ordinary differential equations*, ITU, Tbilisi, 1975. [MR0499402](#)
- [13] I. T. KIGURADZE, T. A. CHANTURIA, *Asymptotic properties of solutions of nonautonomous ordinary differential equations*, Kluwer Academic, Dordrecht, 1993. [MR1220223](#)
- [14] I. T. KIGURADZE, B. SHEKHTER, *Singular boundary value problems for ordinary differential equations of the second order*, Viniti, Moscow, 1987.
- [15] T. KUSANO, J.V. MANOJLOVIĆ, J. MILOŠEVIĆ, Intermediate solutions of second order quasi-linear ordinary differential equations in the framework of regular variation, *Appl. Math. Comput.* **219**(2013), 8178–8191. [MR3037526](#)
- [16] A. P. LINDE, *Particle physics and inflationary cosmology*, Harwood Academic, Chur, Switzerland, 1990.
- [17] M. MARINI, S. MATUCCI, A boundary value problem on the half-line for superlinear differential equations with changing sign weight, *Rend. Istit. Mat. Univ. Trieste* **44**(2012), 117–132. [MR3019556](#)
- [18] S. MATUCCI, P. ŘEHÁK, Asymptotics of decreasing solutions of coupled p -Laplacian systems in the framework of regular variation, *Ann. Mat. Pura Appl.* **193**(2014), 837–858. [MR3210171](#)

- [19] I. RACHŮNKOVÁ, L. RACHŮNEK, Asymptotic formula for oscillatory solutions of some singular nonlinear differential equation, *Abstr. Appl. Anal.* **2011**, Art. ID 981401, 9 pp. [MR2802847](#)
- [20] I. RACHŮNKOVÁ, L. RACHŮNEK, J. TOMEČEK, Existence of oscillatory solutions of singular nonlinear differential equations, *Abstr. Appl. Anal.* **2011**, Art. ID 408525, 20 pp. [MR2795071](#)
- [21] I. RACHŮNKOVÁ, J. TOMEČEK, Homoclinic solutions of singular nonautonomous second order differential equations, *Bound. Value Probl.* **2009**, Art. ID 959636, 21 pp. [MR2552066](#)
- [22] I. RACHŮNKOVÁ, J. TOMEČEK, Bubble-type solutions of nonlinear singular problem, *Math. Comp. Modelling* **51**(2010), 658–669. [MR2594716](#)
- [23] I. RACHŮNKOVÁ, J. TOMEČEK, Strictly increasing solutions of a nonlinear singular differential equation arising in hydrodynamics, *Nonlinear Anal.* **72**(2010), 2114–2118. [MR2577608](#)
- [24] I. RACHŮNKOVÁ, J. TOMEČEK, J. STRYJA, Oscillatory solutions of singular equations arising in hydrodynamics, *Adv. Difference Equ.* **2010**, Art. ID 872160, 13 pp. [MR2652448](#)
- [25] M. ROHLER, On the existence of oscillatory solutions of the second order nonlinear ODE, *Acta Univ. Palack. Olomuc. Fac. Rerum Natur. Math.* **51**(2012), No. 2, 107–127. [MR3058877](#)
- [26] J. VAMPOLOVÁ, On existence and asymptotic properties of Kneser solutions to singular second order ODE, *Acta Univ. Palack. Olomuc. Fac. Rerum Natur. Math.* **52**(2013), No. 1, 135–152. [MR3202755](#)