



Stability index of linear random dynamical systems

Anna Cima¹, Armengol Gasull^{1, 2} and Víctor Mañosa ³

¹Departament de Matemàtiques, Facultat de Ciències, Universitat Autònoma de Barcelona,
08193 Cerdanyola del Vallès, Barcelona, Spain

²Centre de Recerca Matemàtica, Edifici Cc, Campus de Bellaterra,
08193 Cerdanyola del Vallès, Barcelona, Spain

³Departament de Matemàtiques, Institut de Matemàtiques de la UPC-BarcelonaTech (IMTech),
Universitat Politècnica de Catalunya, Colom 11, 08222 Terrassa, Spain

Received 28 February 2020, appeared 19 March 2021

Communicated by Mihály Pituk

Abstract. Given a homogeneous linear discrete or continuous dynamical system, its stability index is given by the dimension of the stable manifold of the zero solution. In particular, for the n dimensional case, the zero solution is globally asymptotically stable if and only if this stability index is n . Fixed n , let X be the random variable that assigns to each linear random dynamical system its stability index, and let p_k with $k = 0, 1, \dots, n$, denote the probabilities that $P(X = k)$. In this paper we obtain either the exact values p_k , or their estimations by combining the Monte Carlo method with a least square approach that uses some affine relations among the values $p_k, k = 0, 1, \dots, n$. The particular case of n -order homogeneous linear random differential or difference equations is also studied in detail.


Keywords: stability index, random differential equations, random difference equations, random dynamical systems.

2020 Mathematics Subject Classification: 37H10, 34F05, 39A25, 37C75.

1 Introduction

Nowadays it is unnecessary to emphasize the importance of ordinary differential equations and discrete dynamical systems to model real world phenomena, from physics to biology, from economics to sociology. These dynamical systems, a concept that includes both continuous and discrete models (and even dynamic equations in time-scales), can have undetermined coefficients that in the case of real applications must be adjusted to fixed values that serve to make good predictions: this is known as the identification process. Once these coefficients are fixed we obtain a deterministic model.

In recent years some authors have highlighted the utility of considering random rather than deterministic coefficients to incorporate effects due to errors in the identification process,

 Corresponding author. Email: victor.manosa@upc.edu

natural variability in some of the physical parameters, or as a method to treat and to incorporate uncertainties in the model, see [5,6,21] for examples coming from biological modeling and [11] for examples coming from mechanical systems.

In the same aim that inspires some works like [1,7,14], in this paper we focus on giving a statistical measure of the stability for both discrete and continuous linear dynamical systems,

$$\dot{\mathbf{x}} = A \mathbf{x} \quad \text{or} \quad \mathbf{x}_{k+1} = A \mathbf{x}_k, \quad (1.1)$$

where both $\mathbf{x}, \mathbf{x}_k \in \mathbb{R}^n$ and A is an $n \times n$ real matrix.

More concretely, in the continuous (resp. discrete) case we define the *stability index* of the origin, $s(A)$, as the number of eigenvalues, taking into account their multiplicities, of A with negative real part (resp. modulus smaller than 1). This index coincides with the dimension of the invariant stable manifold of the origin. Notice also that if $s(A) = n$ (resp. $s(A) = 0$) the origin is a global stable attractor (resp. a global unstable repeller).

In this work we study the probabilities p_k for a linear dynamic system (1.1) to have a given stability index k when the parameters of the matrix A are random variables with a given natural distribution. As we will see in Section 2, this distribution must be that all the elements of A are *independent and identically distributed* (i.i.d.) normal random variables with zero mean. We also will study the same question for linear n -th order differential equations and for linear difference equations.

We also remark that our results can be extrapolated to know a measure of the stability behaviour of critical or fixed points for general non-linear dynamical systems, because near them they can be written as

$$\dot{\mathbf{x}} = A \mathbf{x} + f(\mathbf{x}), \quad \text{or} \quad \mathbf{x}_{k+1} = A \mathbf{x}_k + f(\mathbf{x}_k),$$

with f being a non-linear term vanishing at zero. Moreover, while the situation where the origin is non-hyperbolic is negligible, in the complementary situation, the stability index of the linear part coincides with the dimension of the local stable manifold at the point.

In the continuous case, the key tool to know the stability index of a matrix is the Routh–Hurwitz criterion, see for instance [10, §15.715, p. 1076]. This approach allows to know the number of roots of a polynomial with negative real part in terms of algebraic inequalities among its coefficients. Similarly, its counterpart for the discrete case is called the Jury criterion. It is worth observing that in fact both are equivalent and it is possible to get one from the other by using a Möbius transformation that sends the left hand part of the complex plane into the complex ball of radius 1.

In all the cases, when we do not know how to compute analytically the true probabilities, we introduce a two step approach to obtain estimations of them:

- **Step 1:** We start using the celebrated Monte Carlo method. Recall that this computational algorithm relies on repeated random sampling and gives estimations of the true probabilities based on the law of large numbers and the law of iterated logarithm, see [2,3,13,18]. It is the case that using M samples this approach gives the true value with an absolute error of order $O(((\log \log M)/M)^{1/2})$, which practically behaves as $O(M^{-1/2})$, where O stands for the usual Landau notation. In all our simulations we will work with $M = 10^8$, so our first approaches to the desired probabilities will have an asymptotic absolute error of order 10^{-4} . More detailed explanations of the sharpness of our estimations for this value of M are given in Section 3.2 by using the Chebyshev inequality and the Central limit theorem.

We have used the default in-built pseudo-random number generator in the Statistics package of Maple in our simulations*. This procedure use the Mersenne Twister method with period $2^{19937} - 1$ to generate uniformly-distributed pseudo-random numbers, and then the Ziggurat method, which is a kind of rejection sampling algorithm, to obtain the normally-distributed pseudo-random numbers, see [16] and [17]. Observe that our sample size $M = 10^8$ is much smaller than the period of the pseudo-random number generator, which is greater than 10^{6001} .

- **Step 2:** Since the results of the plain Monte Carlo simulations do not satisfy certain linear constraints concerning the true probabilities, we propose to correct them by using a least squares approach. We take as final estimates of the true probabilities the least squares solution ([20, Def. 6.1]) of the inconsistent overdetermined system obtained when relative frequencies of the Monte Carlo simulation are forced to satisfy these linear constraints. See Section 3.2 for more details. We would like to remark that there are other options to improve plain Monte Carlo simulations like variance reduction and quasi-Monte Carlo methods [2, 13].

To have a flavour of the type of results that we will obtain we describe several consequences of some of our results for linear homogeneous differential or difference equations of order n with constant coefficients (see Sections 5 and 7). A first result is that in both cases the expected stability index is $n/2$. Moreover, let r_n denote the probability of the 0 solution to be a global stable attractor (stability index equals n) for them. Then, for differential equations, $r_n \leq 1/2^n$. Furthermore, $r_1 = 1/2$, $r_2 = 1/4$, $r_3 = 1/16$ and our two step approach gives that $r_4 \simeq 0.00925$, $r_5 \simeq 0.00071$, and that r_k is smaller than 10^{-4} for bigger k . In the case of difference equations we prove that $r_1 = 1/2$ and $r_2 = \frac{1}{\pi} \arctan(\sqrt{2}) \simeq 0.304$.

2 A suitable probability space

In our approach, the starting point is to determine which is the natural choice of the probability space and the distribution law of the coefficients of the linear dynamical system. Only after this step is fixed we can ask for the probabilities of some dynamical features or some phase portraits.

For completeness, we start with some previous considerations and with an example, already considered in the literature, see [1, 14, 23]. Consider the planar linear differential system:

$$\begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \quad (2.1)$$

where A, B, C, D are random variables, so we can set the sample space to be $\Omega = \mathbb{R}^4$. It is plausible to require that these real random variables are independent and identically distributed (i.i.d.) and continuous. Also, according to the *principle of indifference* (or principle of insufficient reason) [8], it would seem reasonable to impose that these variables were such that the random vector (A, B, C, D) had some kind of uniform distribution in \mathbb{R}^4 . But there is no uniform distribution for unbounded probability spaces. Nevertheless, there is a natural choice for the distribution of the variables A, B, C and D .

Indeed, it is well-known that the phase portrait of the above system does not vary if we multiply the right-hand side of both equations by a positive constant (which corresponds to a

*Concretely, we use the commands `RandomVariable(Normal(0,1))` and `Sample`.

proportional change in the time scale). This means that in the space of parameters, \mathbb{R}^4 , all the systems with parameters belonging to the same half-straight line passing through the origin are topologically equivalent and in particular have the same stability index. Hence, we can ask for a probability distribution density f of the coefficients such that the random vector

$$\left(\frac{A}{S}, \frac{B}{S}, \frac{C}{S}, \frac{D}{S} \right), \quad \text{with } S = \sqrt{A^2 + B^2 + C^2 + D^2}, \quad (2.2)$$

has a uniform distribution on the sphere $S^3 \subset \mathbb{R}^4$. This achieves our objective, since S^3 is a compact set.

The question is: which are the probability densities f that give rise to a uniform distribution of the vector (2.2) on the sphere? The answer is that, just assuming that f is continuous and positive, f must be the density of a normal random variable with zero mean. Moreover, this result is true for arbitrary dimension: see the next theorem. We remark that the converse result is well-known [15, 19].

Theorem 2.1. *Let X_1, X_2, \dots, X_n be i.i.d. one-dimensional random variables with a continuous positive density function f . The random vector*

$$\left(\frac{X_1}{S}, \frac{X_2}{S}, \dots, \frac{X_n}{S} \right), \quad \text{with } S = \left(\sum_{i=1}^n X_i^2 \right)^{1/2},$$

has a uniform distribution in $S^{n-1} \subset \mathbb{R}^n$ if and only if each X_i is a normal random variable with zero mean.

Curiously, in the case that we cannot assign uniform distributions, there is an extension of the indifference principle which suggests to use those distributions that maximize the entropy, i.e. the quantity $h(f) = -\int_{\Omega} f(x) \ln(f(x)) dx$ for any given density f . The one-dimensional random variables with continuous probability density function f on $\Omega = \mathbb{R}$ that maximize the entropy are again the Gaussian ones, [8, Thm 3.2].

Of course, if instead of properties concerning general dynamical systems one focuses on particular models in which the parameters have specific restrictions—due to physical or biological reasons—one must consider other type of distributions, see for instance [21].

Using Theorem 2.1, and going back to the initial motivating example, in order to study (2.1) we have to consider the probability space (Ω, \mathcal{F}, P) where $\Omega = \mathbb{R}^4$, \mathcal{F} is the σ -algebra generated by the open sets of \mathbb{R}^4 and $P : \mathcal{F} \rightarrow [0, 1]$ is the probability function with density $\frac{1}{4\pi^2} e^{-(a^2+b^2+c^2+d^2)/2}$, where for simplicity we take variance 1 in each marginal density function.

For instance, assume that we want to compute the probability α of system (2.1) to have exactly one eigenvalue with negative real part. Next, we observe that the probability of having one null eigenvalue is zero. This is because the event which characterizes this possibility is a subset of an event which is itself described by an algebraic equality between the random variables A, B, C, D . This subset has Lebesgue measure zero and therefore, by virtue of the fact that the joint distribution is continuous, the probability of this event, and therefore the event characterizing the null eigenvalue, must also be zero. Thus we have that α coincides with the probability of having a saddle (stability index 1) at the origin, i.e. $AD - BC < 0$. Then, the open set $\mathcal{U} := \{(a, b, c, d) \in \mathbb{R}^4 : ad - bc < 0\}$ belongs to \mathcal{F} and

$$\alpha = P(AD - BC < 0) = \frac{1}{4\pi^2} \int_{\mathcal{U}} e^{-\frac{a^2+b^2+c^2+d^2}{2}} da db dc dd,$$

which is 1/2 by symmetry, as we will see.

Proof of Theorem 2.1. Let (X_1, \dots, X_n) be the random vector associated with the random variables of the statement, with joint continuous density function $g(x_1, \dots, x_n)$. We claim that

$$g(x_1, \dots, x_n) = h(x_1^2 + \dots + x_n^2), \quad (2.3)$$

for some continuous function h .

Taking spherical coordinates, we consider the new random vector $(R, \Theta) \in \mathbb{R}^n$ where $R = (X_1^2 + \dots + X_n^2)^{1/2}$ and $\Theta = (\Theta_1, \dots, \Theta_{n-1})$. We have $X_1 = R \cos \Theta_1$, $X_2 = R \sin \Theta_1 \cos \Theta_2, \dots$, $X_{n-1} = R \sin \Theta_1 \sin \Theta_2 \cdots \sin \Theta_{n-2} \cos \Theta_{n-1}$ and $X_n = R \sin \Theta_1 \sin \Theta_2 \cdots \sin \Theta_{n-2} \sin \Theta_{n-1}$. By the change of variables theorem, the joint density function of (R, Θ) is

$$g_{R,\Theta}(r, \theta) = g(r \cos(\theta_1), \dots, r \sin(\theta_1) \cdots \sin(\theta_{n-1})) r^{n-1} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \cdots \sin(\theta_{n-2}) \cdot \chi$$

where $\theta = (\theta_1, \dots, \theta_{n-1})$, and

$$\chi := \chi_{[0,\infty)}(r) \cdot \chi_{[0,2\pi)}(\theta_{n-1}) \cdot \prod_{i=1}^{n-2} \chi_{[0,\pi)}(\theta_i),$$

where χ_A stands for the characteristic function of the set A .

The density function of (R, Θ) conditioned to R , $g_{\Theta|R}$, is

$$g_{\Theta|R}(r, \theta) := \frac{g_{R,\Theta}(r, \theta)}{g_R(r)},$$

where $g_R(r)$ is the marginal density of R :

$$g_R(r) := \int_0^\pi \cdots \int_0^\pi \int_0^{2\pi} g(r \cos(\theta_1), \dots, r \sin(\theta_1) \cdots \sin(\theta_{n-1})) d\mathcal{S},$$

where $d\mathcal{S} = r^{n-1} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \cdots \sin(\theta_{n-2}) d\theta_{n-1} \cdots d\theta_1$ is the n -dimensional surface element in spherical coordinates.

To prove the statement, we need to characterize which are the joint density functions $g(x_1, \dots, x_n)$ such that when we fix $R = r$, the probability on the $(n-1)$ -dimensional sphere of radius r , denoted by $\mathbb{S}^{n-1}(r)$, is uniformly distributed. In such a case the partial spherical segment $\Sigma_r = \{R = r, \theta_i \in [\alpha_i, \beta_i] \text{ for } i = 1, \dots, n-1\}$ must have probability $P(\Sigma_r) = \mathcal{S}(\Sigma_r) / \mathcal{S}(\mathbb{S}^{n-1}(r))$ where \mathcal{S} denotes the surface area. Set $\alpha = (\alpha_1, \dots, \alpha_{n-1})$ and $\beta = (\beta_1, \dots, \beta_{n-1})$. Notice that

$$\mathcal{S}(\Sigma_r) = \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} d\mathcal{S} = r^{n-1} A(\alpha, \beta)$$

where

$$A(\alpha, \beta) = \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \cdots \sin(\theta_{n-2}) d\theta_{n-1} \cdots d\theta_1$$

and $\mathcal{S}(\mathbb{S}^{n-1}(r)) = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} r^{n-1}$. Hence, on the one hand,

$$P(\Sigma_r) = \Gamma\left(\frac{n}{2}\right) \frac{A(\alpha, \beta)}{2\pi^{\frac{n}{2}}},$$

which does not depend on r . On the other hand,

$$P(\Sigma_r) = \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} g_{\Theta|R} d\theta$$

where $d\theta = d\theta_{n-1} \cdots d\theta_2 d\theta_1$. This implies that

$$\begin{aligned} & \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} \frac{g(r \cos(\theta_1), \dots, r \sin(\theta_1) \cdots \sin(\theta_{n-1})) r^{n-1} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \cdots \sin(\theta_{n-2}) \cdot \chi}{g_R(r)} d\theta \\ &= \frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}}} \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \cdots \sin(\theta_{n-2}) d\theta, \end{aligned}$$

for all $\alpha_i, \beta_i \in [0, \pi)$ for $i = 1, \dots, n-2$ with $\alpha_i < \beta_i$ and $\alpha_{n-2}, \beta_{n-2} \in [0, 2\pi)$ with $\alpha_{n-2} < \beta_{n-2}$. This last equality implies that almost everywhere

$$\frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}}} = \frac{g(r \cos(\theta_1), \dots, r \sin(\theta_1) \cdots \sin(\theta_{n-1})) r^{n-1}}{g_R(r)},$$

and therefore $g(r \cos(\theta_1), \dots, r \sin(\theta_1) \cdots \sin(\theta_{n-1}))$ is a function that only depends on r . In consequence, writing this fact in Cartesian coordinates, we get that almost everywhere $g(x_1, \dots, x_n) = h(x_1^2 + \cdots + x_n^2)$, for some continuous function h and the claim (2.3) follows.

Now we complete the proof. Since X_1, \dots, X_n are i.i.d. with positive density f , we know that $g(x_1, \dots, x_n) = f(x_1) \cdots f(x_n)$. So equation (2.3) can be expressed as

$$f(x_1) \cdots f(x_n) = h(x_1^2 + \cdots + x_n^2) \quad \text{for all } (x_1, \dots, x_n) \in \mathbb{R}^n$$

where h is a positive function. Taking $x_2 = \cdots = x_n = 0$ we have that $f(x_1) f(0)^{n-1} = h(x_1^2)$ and $h(0) = (f(0))^n > 0$. Thus,

$$f(x_1) \cdots f(x_n) = \frac{h(x_1^2)}{(f(0))^{n-1}} \cdots \frac{h(x_n^2)}{(f(0))^{n-1}} = \frac{h(x_1^2)}{(f(0))^n} \cdots \frac{h(x_n^2)}{(f(0))^n} (f(0))^n = h(x_1^2 + \cdots + x_n^2).$$

Hence, using that $h(0) = (f(0))^n > 0$,

$$\frac{h(x_1^2)}{h(0)} \cdots \frac{h(x_n^2)}{h(0)} = \frac{h(x_1^2 + \cdots + x_n^2)}{h(0)}.$$

Taking $H(\xi) := h(\xi)/h(0)$, and $u_i = x_i^2$, it holds that

$$H(u_1) \cdots H(u_n) = H(u_1 + \cdots + u_n) \quad \text{with } H(0) = 1. \quad (2.4)$$

Hence, $\varphi(u) = \log(H(u))$ is a continuous function that satisfies Cauchy's functional equation

$$\varphi(u_1) + \cdots + \varphi(u_n) = \varphi(u_1 + \cdots + u_n) \quad \text{with } \varphi(0) = 0.$$

It is well-known that all its continuous solutions are $\varphi(x) = ax$, for some $a \in \mathbb{R}$. Hence all continuous solutions of (2.4) are $H(x) = e^{ax}$.

As a consequence, $f(x) = b e^{ax^2}$ for some $(a, b) \in \mathbb{R}^2$. Since f is a density function, $a < 0$. Moreover, using $\int_{-\infty}^{\infty} b e^{ax^2} dx = b \sqrt{-\pi/a} = 1$, and setting $a = -1/(2\sigma^2)$, we get that

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{x^2}{2\sigma^2}},$$

so each variable X_i is a normal random variable $N(0, \sigma^2)$.

The converse part is straightforward and well-known [15, 19]. \square

Remark 2.2. The continuity condition for f in Theorem 2.1 is relevant since Equation (2.4) also admits non-continuous solutions that can be constructed, for instance, from non-continuous solutions of Cauchy's functional equation known for $n = 2$, see [12].

3 A preliminary result and methodology

We will investigate the probabilities of having a certain stability index for several linear dynamical systems with random coefficients. In particular we consider:

- (a) Differential systems $\dot{\mathbf{x}} = A \mathbf{x}$ where $\mathbf{x} \in \mathbb{R}^n$ and A is a real constant $n \times n$ matrix,
- (b) Homogeneous linear differential equation of order n with constant coefficients: $a_n x^{(n)} + a_{n-1} x^{(n-1)} + \dots + a_1 x' + a_0 x = 0$,
- (c) Linear discrete systems $b \mathbf{x}_{k+1} = A \mathbf{x}_k$ where $\mathbf{x}_k \in \mathbb{R}^n$, $b \in \mathbb{R}$; and A is a real constant $n \times n$ matrix,
- (d) Linear homogeneous difference equation of order n with constant coefficients $a_n x_{k+n} + a_{n-1} x_{k+n-1} + \dots + a_1 x_{k+1} + a_0 x_k = 0$.

Notice that in the four situations the behaviour of the dynamical systems does not change if we multiply all the involved constants by the same positive real number. This fact situates the four problems in the same context that the motivating example (2.1). Hence, following the results of Section 2, in all the cases, we may take the coefficients to be i.i.d. random normal variables with zero mean and variance 1.

Hence in every case we have a well-defined probability space (Ω, \mathcal{F}, P) , where $\Omega = \mathbb{R}^m$, with $m = n^2, n+1, n^2+1$ or $n+1$ according we are in case (a), (b), (c) or (d), respectively, \mathcal{F} is the σ -algebra generated by the open sets and for each $\mathcal{A} \in \mathcal{F}$,

$$P(\mathcal{A}) = \frac{1}{(\sqrt{2\pi})^m} \int_{\mathcal{A}} e^{-\|\mathbf{a}\|^2/2} d\mathbf{a}, \quad (3.1)$$

where $\mathbf{a} = (a_1, a_2, \dots, a_m)$, $\|\mathbf{a}\|^2 = \sum_{j=1}^m a_j^2$ and $d\mathbf{a} = da_1 da_2 \dots da_m$. For instance the matrices A appearing in case (a) and (c) are the so called *random matrices*.

The use of Routh–Hurwitz algorithm is a very useful tool to count the number of roots of a polynomial with negative real parts and it is implemented in many computer algebra systems. These conditions are given in terms of algebraic inequalities among the coefficients of the polynomials. Let us recall how to use it to count the number of roots with modulus less than one of a polynomial and, hence, to obtain the so called Jury conditions.

Given any polynomial $Q(\lambda) = q_n \lambda^n + q_{n-1} \lambda^{n-1} + \dots + q_1 \lambda + q_0$ with $q_j \in \mathbb{C}$, by using the conformal transformation $\lambda = \frac{z+1}{z-1}$, we get the associated polynomial

$$Q^*(z) = q_n (z+1)^n + q_{n-1} (z+1)^{n-1} (z-1) + \dots + q_0 (z-1)^n. \quad (3.2)$$

It is straightforward to observe that $\lambda_0 \in \mathbb{C}$ is a root of $Q(\lambda)$ such that $|\lambda_0| < 1$ if and only if $z_0 = (\lambda_0 + 1)/(\lambda_0 - 1)$ is a root of $Q^*(z)$ such that $\text{Re}(z_0) < 0$.

Hence, because Routh–Hurwitz and Jury conditions are semi-algebraic, in every case the random variable X that assigns to each dynamical system its stability index $k, 0 \leq k \leq n$, is measurable. Hence $\mathcal{A}_k := \{\mathbf{a} \in \mathbb{R}^m : X(\mathbf{a}) = k\} \in \mathcal{F}$ and its probability $p_k := P(\mathcal{A}_k)$ is well-defined. Observe also that the non-hyperbolic cases are totally negligible because in their characterization some algebraic equalities appear. In this paper we will either calculate or estimate in the four situations the values p_k for $k \leq 10$.

3.1 A preliminary result

In three of the above considered cases we will apply the following auxiliary result:

Lemma 3.1. *Let (Ω, \mathcal{F}, P) be a probability space and let $Y : \Omega \rightarrow \mathbb{R}$ be a discrete random variable with image $\text{Im}(Y) = \{0, 1, \dots, n\}$, and probability mass function $p_k = P(Y = k)$ such that $p_k = p_{n-k}$ for all $k = 0, \dots, n$. Then $E(Y) = \sum_{k=0}^n k p_k = n/2$. Moreover*

(a) *If n is odd then $2 \sum_{k=0}^{\frac{n-1}{2}} p_k = 1$. In particular, when $n = 1$, $p_0 = p_1 = \frac{1}{2}$.*

(b) *If n is even and $n \geq 2$ then $2 \sum_{k=0}^{\frac{n}{2}-1} p_k + p_{\frac{n}{2}} = 1$.*

*If, additionally, n is even** and $\sum_{i \text{ odd}} p_i = \sum_{i \text{ even}} p_i = \frac{1}{2}$, then*

(c) *If $\frac{n}{2}$ is even, then $2 \sum_{k=0, k \text{ even}}^{\frac{n}{2}-2} p_k + p_{\frac{n}{2}} = \frac{1}{2}$, and $2 \sum_{k=1, k \text{ odd}}^{\frac{n}{2}-1} p_k = \frac{1}{2}$. In particular, when $n = 4$, $p_1 = p_3 = \frac{1}{4}$, $p_2 = \frac{1}{2} - 2p_0$ and $p_4 = p_0$.*

(d) *If $\frac{n}{2}$ is odd, then $2 \sum_{k=0, k \text{ even}}^{\frac{n}{2}-1} p_k = \frac{1}{2}$, and $\sum_{k=1, k \text{ odd}}^{\frac{n}{2}-2} p_k + p_{\frac{n}{2}} = \frac{1}{2}$. In particular, when $n = 2$, $p_0 = p_2 = \frac{1}{4}$ and $p_1 = \frac{1}{2}$.*

Proof. We start proving that $E(Y) = n/2$. Assume for instance that n is odd. Since $p_k = p_{n-k}$, it holds that $k p_k + (n-k) p_{n-k} = n p_k$, for each $k \leq (n-1)/2$. Hence,

$$\begin{aligned} E(Y) &= n p_0 + n p_1 + \dots + n p_{\frac{n-1}{2}} = \frac{n}{2} (2 p_0 + 2 p_1 + \dots + 2 p_{\frac{n-1}{2}}) \\ &= \frac{n}{2} ((p_0 + p_n) + (p_1 + p_{n-1}) + \dots + (p_{\frac{n-1}{2}} + p_{\frac{n+1}{2}})) = \frac{n}{2}. \end{aligned}$$

When n is even the proof is similar.

The proof of all the four items is straightforward and we omit it. \square

3.2 Experimental methodology

In every case considered in the paper, when we can not give an exact value of the probabilities p_k we start estimating them by using the *Monte Carlo* method, see [18]. The estimates obtained (namely, the observed relative frequencies) are then improved via the *least squares* method, by using the linear constraints given in Corollaries 4.2, 5.2 and 7.4.

In every case we will use Monte Carlo method with $M = 10^8$ to obtain an estimation, say \tilde{p} , for a probability $p := P(\mathcal{A})$ like the one given in equality (3.1) for different measurable sets \mathcal{A} . Further details for each concrete situation are given in each of the following subsections.

In brief, recall that \tilde{p} is given by the proportion of samples that are in \mathcal{A} . For studying, for a given M , how close are p and \tilde{p} , let $B_j, j = 1, \dots, M$ be i.i.d. Bernoulli random variables, where each one of them that takes the value 1 with probability p and the value 0 with probability $1 - p$.

Define $P_M = \frac{1}{M} \sum_{j=1}^M B_j$. Then, the value obtained for the random variable P_M , \tilde{p} is the approximation of p given by the Monte Carlo method. Let us see, by using Chebyshev inequality or the Central limit theorem, that with very high probability, \tilde{p} is a good approximation of p .

Notice first that $E(P_M) = p$ and due to the independence of the B_j ,

$$\text{Var}(P_M) = \text{Var}\left(\frac{1}{M} \sum_{j=1}^M B_j\right) = \frac{1}{M^2} M \text{Var}(B_1) = \frac{p(1-p)}{M} \leq \frac{1}{4M}$$

**When n is odd the imposed equalities automatically hold.

because $p(1-p) \leq 1/4$. Recall also that for each $\varepsilon > 0$ and any random variable X , with $E(X^2) < \infty$, the Chebyshev inequality reads as

$$P(|X - E(X)| < \varepsilon) \geq 1 - \frac{\text{Var}(X)}{\varepsilon^2}.$$

Hence, applying the Chebyshev inequality to $X = P_M$ we get that

$$P(|P_M - p| < \varepsilon) \geq 1 - \frac{p(1-p)}{M\varepsilon^2} \geq 1 - \frac{1}{4M\varepsilon^2}.$$

Taking $M = 10^8$, as in our computations, denoting $\tilde{p} = P_{10^8}$, and considering $\varepsilon = 10^{-3}$ we get that the above probability gives the following conservative estimate of the reliability of the method

$$P(|\tilde{p} - p| < 10^{-3}) \geq 1 - \frac{1}{400} = \frac{399}{400} = 0.9975.$$

Let us see, by using the Central limit theorem, that the above probability seems to be much bigger. By this theorem we know that for M big enough, and $p(1-p)M$ also big enough, the distribution of the random variable

$$\frac{P_M - E(P_M)}{\sqrt{\text{Var}(P_M)}} = \frac{P_M - p}{\sqrt{\frac{p(1-p)}{M}}}$$

can be practically considered to be a random variable Z with distribution $N(0,1)$. In fact, in Statistics it is usually imposed that $p(1-p)M > 18$. Hence, unless p is very close to 0 or 1, the value $M = 10^8$ is big enough. Hence

$$\begin{aligned} P(|P_M - p| < \varepsilon) &= P\left(\frac{\sqrt{M}|P_M - p|}{\sqrt{p(1-p)}} < \frac{\varepsilon\sqrt{M}}{\sqrt{p(1-p)}}\right) \simeq P\left(|Z| < \frac{\varepsilon\sqrt{M}}{\sqrt{p(1-p)}}\right) \\ &= 2\Phi\left(\frac{\varepsilon\sqrt{M}}{\sqrt{p(1-p)}}\right) - 1 > 2\Phi(2\varepsilon\sqrt{M}) - 1, \end{aligned}$$

where Φ is the distribution function of a $N(0,1)$ random variable. Taking again $M = 10^8$ and either $\varepsilon = 10^{-3}$ or $\varepsilon = 2 \times 10^{-4}$ we get

$$P(|\tilde{p} - p| < 10^{-3}) \gtrsim 2\Phi(20) - 1 > 1 - 10^{-88}$$

or

$$P(|\tilde{p} - p| < 2 \times 10^{-4}) \gtrsim 2\Phi(4) - 1 > 0.99993.$$

In fact, for instance looking at the values p_k of Table 4.1 for $n = 2$ in Section 4, that can also be obtained analytically, we get that $|\tilde{p}_k - p_k| \leq 6 \times 10^{-5}$, for $k = 0, 1, 2$. So, the actual bound is smaller than the bounds obtained above.

Finally, to illustrate how the error decays when the sample size increases, we show the evolution of the errors in one case where the true probabilities are known. We consider the second order difference equation $A_2x_{k+2} + A_1x_{k+1} + A_0x_k = 0$ where A_i are i.i.d. random variables with $N(0,1)$ distribution. The stability index is given by the number of zeroes with modulus smaller than 1 of the characteristic polynomial $Q(\lambda) = A_2\lambda^2 + A_1\lambda + A_0$. Let X be the random variable that counts the number of roots with modulus smaller than 1 of $Q(\lambda)$, and $p_k = P(X = k)$ for $k = 0, 1, 2$. The true value of the probabilities p_k is obtained in Corollary 7.4.

Performing Monte Carlo simulations with $M = 10^m$ with $m = 2, \dots, 10$ we obtain the observed frequencies $\tilde{p}_2(m)$ shown in Table 3.1. These frequencies are the estimated probabilities for the origin to be asymptotically stable. Notice that in Proposition 7.3 and in Corollary 7.4 we prove that $p_0 = p_2 = \arctan(\sqrt{2})/\pi$ and, of course, $p_1 = 1 - p_0 - p_2 = 2 \arctan(1/\sqrt{2})/\pi$. For $M = 10^m$ we denote the absolute error $e_m = |\tilde{p}_2(m) - p_2|$:

$M = 10^2$	$M = 10^3$	$M = 10^4$
$\tilde{p}_2(2) = 0.37$	$\tilde{p}_2(3) = 0.319$	$\tilde{p}_2(4) = 0.3102$
$e_2 \approx 0.065913276015$	$e_3 \approx 0.014913276015$	$e_4 \approx 0.006113276015$
$M = 10^5$	$M = 10^6$	$M = 10^7$
$\tilde{p}_2(5) = 0.30416$	$\tilde{p}_2(6) = 0.303892$	$\tilde{p}_2(7) = 0.3041241$
$e_5 \approx 0.000073276015$	$e_6 \approx 0.000194723985$	$e_7 \approx 0.000037376015$
$M = 10^8$	$M = 10^9$	$M = 10^{10}$
$\tilde{p}_2(8) = 0.30406079$	$\tilde{p}_2(9) = 0.304076699$	$\tilde{p}_2(10) = 0.304079098$
$e_8 \approx 0.000025933985$	$e_9 \approx 0.000010024985$	$e_{10} \approx 0.000007625985$

Table 3.1: Observed frequency and absolute error of p_2 for second order difference equations, using that $p_2 = \arctan(\sqrt{2})/\pi \approx 0.304086723985$.

With the above results, the regression line of $Y = \log(e_m)$ versus $X = \log(M) = m \log(10)$ is $Y = -0.505X - 1.260$ with $R^2 = 0.893$. The slope is therefore $-0.505 \approx -1/2$ as was expected a priori since, in practice, the absolute error behaves as $O(M^{-1/2})$ as $M \rightarrow \infty$ (see the Step 2 in the Introduction).

A more detailed explanation of the second step, about the improvement of the Monte Carlo estimations using the least squares method, is as follows: the probabilities p_k satisfy some affine relations, like the ones in Lemma 3.1 or the ones in Proposition 7.3 below. Then, if we denote $\mathbf{p} = (p_0, \dots, p_n)^t \in \mathbb{R}^{n+1}$ it is possible to write $\mathbf{p} = \mathbf{M}\mathbf{q} + b$ where $\mathbf{q} \in \mathbb{R}^k$ with $k \leq n$ is a vector whose components are different elements of p_0, \dots, p_n ; $\mathbf{M} \in \mathcal{M}_{n \times k}(\mathbb{R})$; and $b \in \mathbb{R}^k$. Let $\tilde{\mathbf{p}} = (\tilde{p}_0, \dots, \tilde{p}_n)^t$ be the vector with the estimated probabilities obtained by the observed relative frequencies using the Monte Carlo method. Then, we can find the least squares solution [20, Def. 6.1] of the the system,

$$\tilde{\mathbf{p}} = \mathbf{M}\hat{\mathbf{q}} + b, \quad (3.3)$$

which is

$$\hat{\mathbf{q}} = (\mathbf{M}^t \cdot \mathbf{M})^{-1} \cdot \mathbf{M}^t \cdot (\tilde{\mathbf{p}} - b), \quad (3.4)$$

see [9, Sect. 5.7, p. 198] or [22, p. 200]. So we can find some improved estimations $\hat{\mathbf{p}}$, via

$$\hat{\mathbf{p}} = \mathbf{M}\hat{\mathbf{q}} + b. \quad (3.5)$$

Some detailed examples are given in Sections 4, 5 and 7.

4 Linear random differential systems

Consider linear differential systems $\dot{\mathbf{x}} = A\mathbf{x}$ where $\mathbf{x} \in \mathbb{R}^n$, $A \in \mathcal{M}_{n \times n}(\mathbb{R})$, where A is a random matrix whose entries are i.i.d. random variables with $N(0, 1)$ distribution. Let X be the random variable that counts the number of eigenvalues of A with negative real part, $s(A)$.

Proposition 4.1. *With the above notations, set $p_k = P(X = k)$. The following holds:*

(a) $\sum_{k=0}^n p_k = 1$.

(b) For all $k \in \{0, 1, \dots, n\}$, $p_k = p_{n-k}$.

(c) $\sum_{i \text{ odd}} p_i = \sum_{i \text{ even}} p_i = \frac{1}{2}$.

Proof. The assertion (a) is trivial. To prove (b) we observe that if a matrix A has k eigenvalues with negative real part, then $B = -A$ has $n - k$ eigenvalues with negative real part. Calling q_m the probability that B has m eigenvalues with negative real part, we get that $p_m = q_m$. This is so, because if $X \sim N(0, 1)$ then $-X \sim N(0, 1)$ and as a consequence the entries of A and B have the same distribution. Then, $q_k = p_{n-k}$ and the result follows.

To see (c) we claim that $s(A)$ is even if and only if the determinant of A is positive and, moreover, $P(\det(A) > 0) = 1/2$. From this claim we get the result because $\sum_{i \text{ even}} p_i$ is the probability of $s(A)$ being even. To prove the first part of the claim notice first that we can assume that $0 \neq \det(A) = \lambda_1 \lambda_2 \cdots \lambda_n$, where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the n eigenvalues of A . We write $\lambda_1 \lambda_2 \cdots \lambda_n = (\lambda_1 \lambda_2 \cdots \lambda_k)(\lambda_{k+1} \lambda_{k+2} \cdots \lambda_n)$ where $\lambda_1, \lambda_2, \dots, \lambda_k$ are all the real negative eigenvalues. Observe also that for complex eigenvalues $\lambda \bar{\lambda} > 0$. Hence $\lambda_{k+1} \lambda_{k+2} \cdots \lambda_n > 0$, $\text{sign}(\det(A)) = (-1)^k$ and the condition that $s(A)$ is even is characterized by $\det(A) > 0$. To prove that $P(\det(A) > 0) = 1/2$ note that if B is the matrix obtained by changing the sign of one column of A then $\det(A) \cdot \det(B) < 0$ and hence $P(\det(A) < 0) = P(\det(B) > 0)$. Furthermore, since the entries of A and B have the same distribution we have $P(\det(B) > 0) = P(\det(A) > 0)$, thus $P(\det(A) < 0) = P(\det(A) > 0) = 1/2$. \square

From the above proposition it easily follows:

Corollary 4.2. *Consider $\dot{\mathbf{x}} = A \mathbf{x}$, $\mathbf{x} \in \mathbb{R}^n$ with $A \in \mathcal{M}_{n \times n}(\mathbb{R})$ a random matrix with i.i.d. $N(0, 1)$ entries, let X be the random variable defined above and $p_k = P(X = k)$. Then the probabilities p_k satisfy all the consequences of Lemma 3.1. In particular $E(X) = n/2$.*

Now we reproduce some experiments to estimate the probabilities p_k for low dimensional cases. We apply the Monte Carlo method, that is, for each considered dimension n , we have generated 10^8 matrices $A \in \mathcal{M}_{n \times n}(\mathbb{R})$ whose entries are pseudo-random numbers simulating the realizations on n^2 independent random variables with $N(0, 1)$ distribution. For each matrix A we have computed the characteristic polynomial, and counted the number of eigenvalues with negative real part by using the Routh–Hurwitz zeros counter [10, §15.715, p. 1076]. We are aware that the stability of the calculation of the coefficients of the characteristic polynomial from the entries of a matrix is critical (see [22, pp. 378–379] and references therein); however we have only used this calculation for low dimensions, namely $n \leq 4$. For $n \geq 5$, and in order to decrease the computation time, we have directly computed numerically the eigenvalues of A and counted the number of them with negative real part.

For each considered dimension of the phase space n , and in order to take advantage of the relations stated in Corollary 4.2, we can refine the solutions using the least squares solutions of the inconsistent linear system associated with these relations when using the observed frequencies obtained by the Monte Carlo simulation.

We give details of one example. Set $n = 7$, for instance. By Corollary 4.2 we have $p_3 = p_4 = \frac{1}{2} - p_0 - p_1 - p_2$; $p_5 = p_2$; $p_6 = p_1$ and $p_7 = p_0$. So, using the notation introduced in

Section 3.2, we can write $\mathbf{p} = \mathbf{M}\mathbf{q} + b$, where $\mathbf{p}^t = (p_0, \dots, p_7)$;

$$\mathbf{M} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & -1 & -1 \\ -1 & -1 & -1 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}; \quad \mathbf{q} = \begin{pmatrix} p_0 \\ p_1 \\ p_2 \end{pmatrix}; \quad \text{and } b = \begin{pmatrix} 0 \\ 0 \\ 0 \\ \frac{1}{2} \\ \frac{1}{2} \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$

The observed relative frequencies in our Monte Carlo simulation are

$$\tilde{\mathbf{p}}^t = \left(\frac{31643}{50000000}, \frac{261137}{12500000}, \frac{7124967}{50000000}, \frac{1344047}{4000000}, \frac{33597117}{100000000}, \frac{14248187}{100000000}, \frac{1043913}{50000000}, \frac{63379}{100000000} \right).$$

By finding the least squares solution of the system (3.3) ([9, Sect. 5.7, p. 198] or [22, p. 200]), given by (3.5), we obtain

$$\hat{\mathbf{p}}^t = \left(\frac{25333}{40000000}, \frac{2088461}{100000000}, \frac{28498121}{200000000}, \frac{16799573}{50000000}, \frac{16799573}{50000000}, \frac{28498121}{200000000}, \frac{2088461}{100000000}, \frac{25333}{40000000} \right).$$

The other cases follow similarly.

We summarize the results of our experiments in the Table 4.1, where the observed relative frequencies and the estimates are presented only up to the fifth decimal (in the table, and in the whole paper, frequency stands for relative frequency) because as we already explained in the introduction, the predicted absolute error will be of order 10^{-4} . Observe that in the cases $n = 1, 2$ the true probabilities are known. We include the results of the Monte Carlo simulations for completeness, but it makes no sense to apply the least squares method.

Dimension	Observed frequency	Least squares	Relations (Corol. 4.2)
$n = 1$	$\tilde{p}_0 = 0.49996$ $\tilde{p}_1 = 0.50004$		$p_0 = 0.5$ $p_1 = 0.5$
$n = 2$	$\tilde{p}_0 = 0.24999$ $\tilde{p}_1 = 0.50006$ $\tilde{p}_2 = 0.24995$		$p_0 = 0.25$ $p_1 = 0.5$ $p_2 = 0.25$
$n = 3$	$\tilde{p}_0 = 0.10447$ $\tilde{p}_1 = 0.39542$ $\tilde{p}_2 = 0.39557$ $\tilde{p}_3 = 0.10454$	$\hat{p}_0 = 0.10450$ $\hat{p}_1 = 0.39550$ $\hat{p}_2 = 0.39550$ $\hat{p}_3 = 0.10450$	p_0 $p_1 = \frac{1}{2} - p_0$ $p_2 = \frac{1}{2} - p_0$ $p_3 = p_0$
$n = 4$	$\tilde{p}_0 = 0.03722$ $\tilde{p}_1 = 0.25009$ $\tilde{p}_2 = 0.42556$ $\tilde{p}_3 = 0.24998$ $\tilde{p}_4 = 0.03715$	$\hat{p}_0 = 0.03721$ $\hat{p}_1 = 0.25000$ $\hat{p}_2 = 0.42558$ $\hat{p}_3 = 0.25000$ $\hat{p}_4 = 0.03721$	p_0 $p_1 = \frac{1}{4}$ $p_2 = \frac{1}{2} - 2p_0$ $p_3 = \frac{1}{4}$ $p_4 = p_0$
$n = 5$	$\tilde{p}_0 = 0.01126$ $\tilde{p}_1 = 0.13028$ $\tilde{p}_2 = 0.35848$ $\tilde{p}_3 = 0.35852$ $\tilde{p}_4 = 0.13020$ $\tilde{p}_5 = 0.01126$	$\hat{p}_0 = 0.01126$ $\hat{p}_1 = 0.13024$ $\hat{p}_2 = 0.35850$ $\hat{p}_3 = 0.35850$ $\hat{p}_4 = 0.13024$ $\hat{p}_5 = 0.01126$	p_0 p_1 $p_2 = \frac{1}{2} - p_0 - p_1$ $p_3 = \frac{1}{2} - p_0 - p_1$ $p_4 = p_1$ $p_5 = p_0$

Dimension	Observed frequency	Least squares	Relations (Corol. 4.2)
$n = 6$	$\tilde{p}_0 = 0.00289$ $\tilde{p}_1 = 0.05675$ $\tilde{p}_2 = 0.24710$ $\tilde{p}_3 = 0.38642$ $\tilde{p}_4 = 0.24714$ $\tilde{p}_5 = 0.56810$ $\tilde{p}_6 = 0.00289$	$\hat{p}_0 = 0.00288$ $\hat{p}_1 = 0.05678$ $\hat{p}_2 = 0.24712$ $\hat{p}_3 = 0.38644$ $\hat{p}_4 = 0.24712$ $\hat{p}_5 = 0.05678$ $\hat{p}_6 = 0.00288$	p_0 p_1 $p_2 = \frac{1}{4} - p_0$ $p_3 = \frac{1}{2} - 2p_1$ $p_4 = \frac{1}{4} - p_0$ $p_5 = p_1$ $p_6 = p_0$
$n = 7$	$\tilde{p}_0 = 0.00063$ $\tilde{p}_1 = 0.02089$ $\tilde{p}_2 = 0.14250$ $\tilde{p}_3 = 0.33601$ $\tilde{p}_4 = 0.33597$ $\tilde{p}_5 = 0.14248$ $\tilde{p}_6 = 0.02088$ $\tilde{p}_7 = 0.00063$	$\hat{p}_0 = 0.00063$ $\hat{p}_1 = 0.02088$ $\hat{p}_2 = 0.14249$ $\hat{p}_3 = 0.33600$ $\hat{p}_4 = 0.33600$ $\hat{p}_5 = 0.14249$ $\hat{p}_6 = 0.02088$ $\hat{p}_7 = 0.00063$	p_0 p_1 p_2 $p_3 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_4 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_5 = p_2$ $p_6 = p_1$ $p_7 = p_0$
$n = 8$	$\tilde{p}_0 = 0.00012$ $\tilde{p}_1 = 0.00651$ $\tilde{p}_2 = 0.06948$ $\tilde{p}_3 = 0.24356$ $\tilde{p}_4 = 0.36080$ $\tilde{p}_5 = 0.24346$ $\tilde{p}_6 = 0.06946$ $\tilde{p}_7 = 0.00650$ $\tilde{p}_8 = 0.00012$	$\hat{p}_0 = 0.00012$ $\hat{p}_1 = 0.00650$ $\hat{p}_2 = 0.06948$ $\hat{p}_3 = 0.24350$ $\hat{p}_4 = 0.36080$ $\hat{p}_5 = 0.24350$ $\hat{p}_6 = 0.06948$ $\hat{p}_7 = 0.00650$ $\hat{p}_8 = 0.00012$	p_0 p_1 p_2 $p_3 = \frac{1}{4} - p_1$ $p_4 = \frac{1}{2} - 2p_0 - 2p_2$ $p_5 = \frac{1}{4} - p_1$ $p_6 = p_2$ $p_7 = p_1$ $p_8 = p_0$
$n = 9$	$\tilde{p}_0 = 0.00002$ $\tilde{p}_1 = 0.00171$ $\tilde{p}_2 = 0.02880$ $\tilde{p}_3 = 0.14952$ $\tilde{p}_4 = 0.31987$ $\tilde{p}_5 = 0.31999$ $\tilde{p}_6 = 0.14958$ $\tilde{p}_7 = 0.02878$ $\tilde{p}_8 = 0.00171$ $\tilde{p}_9 = 0.00002$	$\hat{p}_0 = 0.00002$ $\hat{p}_1 = 0.00171$ $\hat{p}_2 = 0.02879$ $\hat{p}_3 = 0.14955$ $\hat{p}_4 = 0.31993$ $\hat{p}_5 = 0.31993$ $\hat{p}_6 = 0.14955$ $\hat{p}_7 = 0.02879$ $\hat{p}_8 = 0.00171$ $\hat{p}_9 = 0.00002$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_5 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_6 = p_3$ $p_7 = p_2$ $p_8 = p_1$ $p_9 = p_0$
$n = 10$	$\tilde{p}_0 = 0$ $\tilde{p}_1 = 0.00038$ $\tilde{p}_2 = 0.01015$ $\tilde{p}_3 = 0.07850$ $\tilde{p}_4 = 0.23987$ $\tilde{p}_5 = 0.34224$ $\tilde{p}_6 = 0.23984$ $\tilde{p}_7 = 0.07849$ $\tilde{p}_8 = 0.01015$ $\tilde{p}_9 = 0.00038$ $\tilde{p}_{10} = 0$	$\hat{p}_0 = 0$ $\hat{p}_1 = 0.00038$ $\hat{p}_2 = 0.01015$ $\hat{p}_3 = 0.07850$ $\hat{p}_4 = 0.23985$ $\hat{p}_5 = 0.34224$ $\hat{p}_6 = 0.23985$ $\hat{p}_7 = 0.07850$ $\hat{p}_8 = 0.01015$ $\hat{p}_9 = 0.00038$ $\hat{p}_{10} = 0$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{4} - p_0 - p_2$ $p_5 = \frac{1}{2} - 2p_1 - 2p_3$ $p_6 = \frac{1}{4} - p_0 - p_2$ $p_7 = p_3$ $p_8 = p_2$ $p_9 = p_1$ $p_{10} = p_0$

Table 4.1: Linear stability indexes for linear random differential systems.

5 Linear random differential equations of order n

In this section we consider linear random homogeneous differential equations of order n

$$A_n x^{(n)} + A_{n-1} x^{(n-1)} + \cdots + A_2 x'' + A_1 x' + A_0 x = 0, \quad (5.1)$$

where $x = x(t)$, the derivatives are taken in respect to t , and A_j are again i.i.d. random variables with $N(0, 1)$ distribution.

To get the stability index for these differential equations we only need to know the probability distributions of the number of roots with negative real part of its associated random characteristic polynomial:

$$Q(\lambda) = A_n \lambda^n + A_{n-1} \lambda^{n-1} + \cdots + A_1 \lambda + A_0.$$

Let X be the random variable that counts the number of roots of $Q(\lambda)$ with negative real parts and define $p_k = P(X = k)$ for $k = 0, 1, \dots, n$.

Proposition 5.1. *Set $p_k = P(X = k)$, where X is the random variable defined above. Then*

- (a) $\sum_{k=0}^n p_k = 1$.
- (b) For all $k \in \{0, 1, \dots, n\}$, $p_k = p_{n-k}$.
- (c) $\sum_{i \text{ odd}} p_i = \sum_{i \text{ even}} p_i = \frac{1}{2}$.

Proof. The proof of (a) is trivial. To prove (b) consider equation (5.1) with its characteristic polynomial $Q(\lambda)$ and also the new differential equation

$$(-1)^n A_n x^{(n)} + (-1)^{n-1} A_{n-1} x^{(n-1)} + \cdots + A_2 x'' - A_1 x' + A_0 x = 0 \quad (5.2)$$

with its characteristic polynomial $Q^*(\lambda) = Q(-\lambda) = (-1)^n A_n \lambda^n + (-1)^{n-1} A_{n-1} \lambda^{n-1} + \cdots - A_1 \lambda + A_0$. Since $Q(\lambda) = 0$ if and only if $Q^*(-\lambda) = 0$ we get that $p_k = p_{n-k}^*$ where p_i^* the probability that $Q^*(\lambda)$ has i roots with negative real part. But also $p_k = p_k^*$ because the coefficients of the equations (5.1) and (5.2) have the same distributions. Hence, the result follows.

Similarly, as in the proof of (c) of Proposition 4.1, we observe that the polynomial $Q(\lambda)$ has an odd number of roots with negative real part if and only if $A_0 \cdot A_n < 0$, because we can neglect the case of having some roots with zero real part. Since the coefficients of (5.1) are symmetric independent random variables, the probability that $Q(\lambda)$ has an odd number of roots with negative real part is

$$P(\{A_0 > 0\} \cap \{A_n < 0\}) + P(\{A_0 < 0\} \cap \{A_n > 0\}) = \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} \times \frac{1}{2} = \frac{1}{2}. \quad \square$$

Corollary 5.2. *Consider the linear random homogeneous differential equation of order n (5.1), with all A_i being i.i.d. $N(0, 1)$ random variables, let X be defined above, and set $p_k = P(X = k)$. Then the probabilities p_k satisfy all the conclusions of Lemma 3.1. In particular $E(X) = n/2$.*

For each n , let r_n be the probability of the origin to be a global stable attractor (asymptotically stable equilibrium) for (5.1), that is $r_n = p_n$. By Proposition 5.1(b) this probability coincides with the probability of being a repeller because $p_n = p_0$. Our results in Proposition 5.4 seem to indicate that r_n decreases with n . Before proving this proposition we need a preliminary result.

Lemma 5.3. *Let U, V, S and T be i.i.d. random variables with standard normal distribution. Then $p^+ := P(U > 0; V > 0; S > 0; T > 0; UT - SV > 0) = 1/32$.*

Proof. Set $\mathcal{A}^\pm = \{U > 0; V > 0; S > 0; T > 0; \pm(UT - SV) > 0\}$, and $\mathcal{A}^0 = \{U > 0; V > 0; S > 0; T > 0; UT - SV = 0\}$. Denote by $p^\pm = P(\mathcal{A}^\pm)$ and $p^0 = P(\mathcal{A}^0)$. Then, since $p^0 = 0$ and $\mathcal{A}^- \cup \mathcal{A}^0 \cup \mathcal{A}^+ = \{U > 0; V > 0; S > 0; T > 0\}$ it holds that $p^+ + p^- = (1/2)^4 = 1/16$. To end the proof it suffices to show that $p^+ = p^-$.

Notice first that

$$\begin{aligned} \mathcal{A}^+ &= \{U > 0; V > 0; S > 0; T > 0; UT - SV > 0\} = \{V > 0; S > 0; T > 0; UT - SV > 0\}, \\ \mathcal{A}^- &= \{U > 0; V > 0; S > 0; T > 0; UT - SV < 0\} = \{U > 0; S > 0; T > 0; SV - UT > 0\}. \end{aligned}$$

This is so, because for instance in the definition of \mathcal{A}^+ , the last inequality can also be written as $U > SV/T > 0$ and from it we know that the condition $U > 0$ can be removed. Finally, interchanging U and V and S and T we get the same relations in the definitions of \mathcal{A}^+ and \mathcal{A}^- . Since all variables are independent $N(0, 1)$, both sets have the same probability and $p^+ = p^-$, as we wanted to prove. \square

Proposition 5.4. *With the above notations, $r_n \leq 1/2^n$, for all $n \geq 1$. Moreover, $r_1 = 1/2$, $r_2 = 1/4$, $r_3 = 1/16$ and $r_4 < r_3/2 = 1/32$.*

Proof of Proposition 5.4. Notice that r_n is the probability that the characteristic polynomial $Q(\lambda)$, associated with the random differential equation (5.1), is a Hurwitz stable polynomial; that is $r_n = P(\text{Every root of } Q(\lambda) \text{ belongs to } \mathfrak{R}^-)$, where $\mathfrak{R}^- = \{z \in \mathbb{C} \text{ such that } \text{Re}(z) < 0\}$. It is well-known that a necessary condition for a polynomial to have every root in \mathfrak{R}^- is that all its coefficients have the same sign. This is so because it holds for polynomials of degree 1 and 2, and this property is preserved when we multiply two polynomials satisfying it. Hence,

$$\{A_0, \dots, A_n \text{ such that all roots of } P(\lambda) \text{ are in } \mathfrak{R}^-\} \subset \left\{ \bigcap_{i=0}^n \{A_i < 0\} \right\} \cup \left\{ \bigcap_{i=0}^n \{A_i > 0\} \right\}. \quad (5.3)$$

Since the variables A_i are independent and symmetric

$$P\left(\bigcap_{i=0}^n \{A_i < 0\}\right) = P\left(\bigcap_{i=0}^n \{A_i > 0\}\right) = \frac{1}{2^{n+1}}.$$

As a consequence,

$$r_n \leq P\left(\bigcap_{i=0}^n \{A_i < 0\}\right) + P\left(\bigcap_{i=0}^n \{A_i > 0\}\right) = \frac{1}{2^n},$$

and the first statement follows.

The equalities $r_1 = 1/2$ and $r_2 = 1/4$ are a simple consequence that for $n = 1, 2$ the inclusion (5.3) is an equality.

Let us prove that $r_3 = p_3 = 1/16$. By using the Routh–Hurwitz criterion [10, §15.715, p. 1076], it can be seen that $a_3\lambda^3 + a_2\lambda^2 + a_1\lambda + a_0$ has every root in \mathfrak{R}^- if and only if all its coefficients have the same sign and moreover $a_1a_2 - a_0a_3 > 0$. Hence, $p_3 = p_3^- + p_3^+$, where $p_3^- := P(A_0 < 0; A_1 < 0; A_2 < 0; A_3 < 0; A_1A_2 - A_0A_3 > 0)$; and $p_3^+ :=$

$P(A_0 > 0; A_1 > 0; A_2 > 0; A_3 > 0; A_1A_2 - A_0A_3 > 0)$, with all the A_i being $N(0,1)$ distributed and independent. Due to their symmetry, the random variables A_i and $-A_i$, for $i = 0, \dots, 3$ have the same distribution and hence $p_3^+ = p_3^-$. Therefore $p_3 = 2p_3^+$. The result follows now by Lemma 5.3, which gives $p_3^+ = 1/32$.

Let us prove that $r_4 < r_3/2$. To compare both probabilities, here it will be more convenient to write the coefficients of the polynomials with subscripts with increasing ordering, that is $q_n(x) = a_0x^n + a_1x^{n-1} + \dots + a_{n-1}x + a_n$. With this notation, which also respects the traditional notation when writing the Hurwitz matrices, and when $a_0 > 0$, the Routh–Hurwitz conditions to have stability index n for $n = 3, 4$ are precisely that the principal minors of the following matrices

$$\begin{pmatrix} a_1 & a_3 & 0 \\ a_0 & a_2 & 0 \\ 0 & a_1 & a_3 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} a_1 & a_3 & 0 & 0 \\ a_0 & a_2 & a_4 & 0 \\ 0 & a_1 & a_3 & 0 \\ 0 & a_0 & a_2 & a_4 \end{pmatrix},$$

are positive, where the left-hand one corresponds to the case $n = 3$ and the other to the case when $n = 4$. Hence, these conditions when $a_0 > 0$ and for $n = 3$ are: $a_1 > 0, a_1a_2 - a_0a_3 > 0$ and $a_3 > 0$. Similarly, for $n = 4$ the conditions are $a_1 > 0, a_1a_2 - a_0a_3 > 0, a_3(a_1a_2 - a_0a_3) - a_4a_1^2 > 0$ and $a_4 > 0$.

Consider now, for $n = 3, 4$, the random polynomials $Q_n(x) = \tilde{A}_0x^n + \tilde{A}_1x^{n-1} + \dots + \tilde{A}_{n-1}x + \tilde{A}_n$, where $\tilde{A}_i \sim N(0,1)$ and are independent (notice that with this notation each coefficient \tilde{A}_i is the coefficient A_{n-i} of the characteristic polynomial). For simplicity we denote with the same name the coefficients of Q_3 and Q_4 although they are different random variables. As above, $r_3 = 2p_3^+$ and $r_4 = 2p_4^+$, where $p_k^+ = P(\mathcal{A}_k^+)$, with

$$\begin{aligned} \mathcal{A}_3^+ &= \{\tilde{A}_0 > 0; \tilde{A}_1 > 0; \tilde{A}_3 > 0; \tilde{A}_1\tilde{A}_2 - \tilde{A}_0\tilde{A}_3 > 0\}, \\ \mathcal{A}_4^+ &= \{\tilde{A}_0 > 0; \tilde{A}_1 > 0; \tilde{A}_3 > \tilde{A}_4\tilde{A}_1^2 / (\tilde{A}_1\tilde{A}_2 - \tilde{A}_0\tilde{A}_3); \tilde{A}_1\tilde{A}_2 - \tilde{A}_0\tilde{A}_3 > 0, \tilde{A}_4 > 0\}. \end{aligned}$$

Notice that if we define

$$\mathcal{B} = \{\tilde{A}_0 > 0; \tilde{A}_1 > 0; \tilde{A}_3 > 0; \tilde{A}_1\tilde{A}_2 - \tilde{A}_0\tilde{A}_3 > 0; \tilde{A}_4 > 0\}$$

it is clear that $P(\mathcal{B}) = p_3^+/2$ and, moreover $\mathcal{A}_4^+ \subset \mathcal{B}$, with the inclusion being strict. Since the joint density is positive and $\mathcal{B} \cap (\mathcal{A}_4^+)^c$ has positive Lebesgue measure, we have $P(\mathcal{B} \cap (\mathcal{A}_4^+)^c) > 0$. Thus $P(\mathcal{A}_4^+) < P(\mathcal{B})$, and hence $p_4^+ = P(\mathcal{A}_4^+) < P(\mathcal{B}) = p_3^+/2$, and $r_4 < r_3/2$, as we wanted to show. \square

Corollary 5.5. *Consider a linear random homogeneous differential equation of order $n = 3$ and the random variable X defined above. Then $p_0 = p_3 = 1/16$ and $p_1 = p_2 = 7/16$.*

Proof. By the above proposition, for $n = 3$, $p_0 = p_3 = r_3 = 1/16$. Hence, by Proposition 5.1, $p_1 = p_2 = 7/16$. \square

The computations in this case are similar to the ones of the previous section and the obtained results are summarized in Table 5.1. We only give some comments for the cases $n = 8$ and 10 , where we have encountered that the vectors $\hat{\mathbf{p}}$ have negative and very small entries. This has occurred because the observed frequencies obtained by the Monte Carlo approach corresponding to these probabilities are not accurate enough. For this reason, we

have made a new optimization step. As before, we use the least squares method to obtain a vector $\hat{\mathbf{p}}$. However, if negative entries appear in this vector (which is clearly objectionable), we force them to be zero and find a new least squares estimate, which still respects the original linear constraints.

We explain this process for the $n = 8$ order case; the $n = 10$ case follows analogously. The observed relative frequencies vector obtained by the Monte Carlo method is

$$\tilde{\mathbf{p}}^t = \left(\frac{1}{50000000}, \frac{6599}{50000000}, \frac{1159359}{50000000}, \frac{4996163}{20000000}, \frac{45377377}{100000000}, \frac{4995607}{20000000}, \frac{2318357}{100000000}, \frac{13497}{100000000}, \frac{1}{100000000} \right).$$

The relations stated in Corollary 5.2 are $p_3 = 1/4 - p_1$, $p_4 = 1/2 - 2p_0 - 2p_2$, $p_5 = p_3$, $p_6 = p_2$, $p_7 = p_1$, $p_8 = p_0$. By solving the system (3.3) with

$$\mathbf{M} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \\ -2 & 0 & -2 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}; \quad \mathbf{q} = \begin{pmatrix} p_0 \\ p_1 \\ p_2 \end{pmatrix}; \quad \text{and } b = \begin{pmatrix} 0 \\ 0 \\ 0 \\ \frac{1}{2} \\ \frac{1}{4} \\ \frac{1}{2} \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

we obtain

$$\hat{\mathbf{q}}^t = \left(-\frac{5779}{200000000}, \frac{13569}{80000000}, \frac{4631293}{200000000} \right).$$

Hence, by (3.5) we get

$$\hat{\mathbf{p}} = \left(\frac{-5779}{200000000}, \frac{13569}{80000000}, \frac{4631293}{200000000}, \frac{19986431}{80000000}, \frac{22687243}{50000000}, \frac{19986431}{80000000}, \frac{4631293}{200000000}, \frac{13569}{80000000}, \frac{-5779}{200000000} \right).$$

So we impose that $p_0 = p_8 = 0$. Thus we have $p_3 = p_5 = 1/4 - p_1$, $p_4 = 1/2 - 2p_0 - 2p_2 = 1/2 - 2p_2$, $p_6 = p_2$ and $p_7 = p_1$. We find the least squares solution of the system

$$\begin{pmatrix} \hat{p}_1 \\ \hat{p}_2 \\ \hat{p}_3 \\ \hat{p}_4 \\ \hat{p}_5 \\ \hat{p}_6 \\ \hat{p}_7 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ -1 & 0 \\ 0 & -2 \\ -1 & 0 \\ 0 & 1 \\ 1 & 0 \end{pmatrix} \cdot \begin{pmatrix} \hat{p}_1^* \\ \hat{p}_2^* \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ \frac{1}{4} \\ \frac{1}{2} \\ \frac{1}{4} \\ 0 \\ 0 \end{pmatrix}.$$

Using (3.4) and (3.5) we obtain

$$\hat{\mathbf{p}}^* = \left(0, \frac{13569}{80000000}, \frac{13882321}{600000000}, \frac{19986431}{80000000}, \frac{136117679}{300000000}, \frac{19986431}{80000000}, \frac{13882321}{600000000}, \frac{13569}{80000000}, 0 \right) \\ \simeq (0, 0.00017, 0.02314, 0.24983, 0.45373, 0.24983, 0.02314, 0.00017, 0).$$

Dimension	Observed frequency	Least squares	Relations (Corol. 5.2 and 5.5)
$n = 1$	$\tilde{p}_0 = 0.49997$ $\tilde{p}_1 = 0.50003$		$p_0 = 0.5$ $p_1 = 0.5$
$n = 2$	$\tilde{p}_0 = 0.24994$ $\tilde{p}_1 = 0.49999$ $\tilde{p}_2 = 0.25007$		$p_0 = 0.25$ $p_1 = 0.5$ $p_2 = 0.25$
$n = 3$	$\tilde{p}_0 = 0.06252$ $\tilde{p}_1 = 0.43743$ $\tilde{p}_2 = 0.43756$ $\tilde{p}_3 = 0.06249$		$p_0 = \frac{1}{16} = 0.0625$ $p_1 = \frac{7}{16} = 0.4375$ $p_2 = \frac{7}{16} = 0.4375$ $p_3 = \frac{1}{16} = 0.0625$
$n = 4$	$\tilde{p}_0 = 0.00928$ $\tilde{p}_1 = 0.24998$ $\tilde{p}_2 = 0.48152$ $\tilde{p}_3 = 0.24994$ $\tilde{p}_4 = 0.00929$	$\hat{p}_0 = 0.00925$ $\hat{p}_1 = 0.25$ $\hat{p}_2 = 0.48150$ $\hat{p}_3 = 0.25$ $\hat{p}_4 = 0.00925$	p_0 $p_1 = \frac{1}{4}$ $p_2 = \frac{1}{2} - 2p_0$ $p_3 = \frac{1}{4}$ $p_4 = p_0$
$n = 5$	$\tilde{p}_0 = 0.00071$ $\tilde{p}_1 = 0.08405$ $\tilde{p}_2 = 0.41526$ $\tilde{p}_3 = 0.41523$ $\tilde{p}_4 = 0.08404$ $\tilde{p}_5 = 0.00071$	$\hat{p}_0 = 0.00071$ $\hat{p}_1 = 0.08404$ $\hat{p}_2 = 0.41525$ $\hat{p}_3 = 0.41525$ $\hat{p}_4 = 0.08404$ $\hat{p}_5 = 0.00071$	p_0 p_1 $p_2 = \frac{1}{2} - p_0 - p_1$ $p_3 = \frac{1}{2} - p_0 - p_1$ $p_4 = p_1$ $p_5 = p_0$
$n = 6$	$\tilde{p}_0 = 0.00003$ $\tilde{p}_1 = 0.01723$ $\tilde{p}_2 = 0.24994$ $\tilde{p}_3 = 0.46562$ $\tilde{p}_4 = 0.24993$ $\tilde{p}_5 = 0.01723$ $\tilde{p}_6 = 0.00003$	$\hat{p}_0 = 0.00005$ $\hat{p}_1 = 0.01720$ $\hat{p}_2 = 0.24995$ $\hat{p}_3 = 0.46560$ $\hat{p}_4 = 0.24995$ $\hat{p}_5 = 0.01720$ $\hat{p}_6 = 0.00005$	p_0 p_1 $p_2 = \frac{1}{4} - p_0$ $p_3 = \frac{1}{2} - 2p_1$ $p_4 = \frac{1}{4} - p_0$ $p_5 = p_1$ $p_6 = p_0$
$n = 7$	$\tilde{p}_0 = 0$ $\tilde{p}_1 = 0.00200$ $\tilde{p}_2 = 0.09571$ $\tilde{p}_3 = 0.40224$ $\tilde{p}_4 = 0.40233$ $\tilde{p}_5 = 0.09573$ $\tilde{p}_6 = 0.00199$ $\tilde{p}_7 = 0$	$\hat{p}_0 = 0$ $\hat{p}_1 = 0.00200$ $\hat{p}_2 = 0.09572$ $\hat{p}_3 = 0.40228$ $\hat{p}_4 = 0.40228$ $\hat{p}_5 = 0.09572$ $\hat{p}_6 = 0.00200$ $\hat{p}_7 = 0$	p_0 p_1 p_2 $p_3 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_4 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_5 = p_2$ $p_6 = p_1$ $p_7 = p_0$
$n = 8$	$\tilde{p}_0 = 0$ $\tilde{p}_1 = 0.00013$ $\tilde{p}_2 = 0.02319$ $\tilde{p}_3 = 0.24981$ $\tilde{p}_4 = 0.45377$ $\tilde{p}_5 = 0.24978$ $\tilde{p}_6 = 0.02318$ $\tilde{p}_7 = 0.00013$ $\tilde{p}_8 = 0$	$\hat{p}_0^* = 0$ $\hat{p}_1^* = 0.00017$ $\hat{p}_2^* = 0.02314$ $\hat{p}_3^* = 0.24983$ $\hat{p}_4^* = 0.45372$ $\hat{p}_5^* = 0.24983$ $\hat{p}_6^* = 0.02314$ $\hat{p}_7^* = 0.00017$ $\hat{p}_8^* = 0$	p_0 p_1 p_2 $p_3 = \frac{1}{4} - p_1$ $p_4 = \frac{1}{2} - 2p_0 - 2p_2$ $p_5 = \frac{1}{4} - p_1$ $p_6 = p_2$ $p_7 = p_1$ $p_8 = p_0$

Dimension	Observed frequency	Least squares	Relations (Corol. 5.2 and 5.5)
$n = 9$	$\tilde{p}_0 = 0$ $\tilde{p}_1 = 0.00001$ $\tilde{p}_2 = 0.00336$ $\tilde{p}_3 = 0.10337$ $\tilde{p}_4 = 0.39328$ $\tilde{p}_5 = 0.39328$ $\tilde{p}_6 = 0.10332$ $\tilde{p}_7 = 0.00338$ $\tilde{p}_8 = 0$ $\tilde{p}_9 = 0$	$\hat{p}_0 = 0$ $\hat{p}_1 = 0.00005$ $\hat{p}_2 = 0.00337$ $\hat{p}_3 = 0.10335$ $\hat{p}_4 = 0.39328$ $\hat{p}_5 = 0.39328$ $\hat{p}_6 = 0.10335$ $\hat{p}_7 = 0.00337$ $\hat{p}_8 = 0.00005$ $\hat{p}_9 = 0$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_5 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_6 = p_3$ $p_7 = p_2$ $p_8 = p_1$ $p_9 = p_0$
$n = 10$	$\tilde{p}_0 = 0$ $\tilde{p}_1 = 0$ $\tilde{p}_2 = 0.00030$ $\tilde{p}_3 = 0.02784$ $\tilde{p}_4 = 0.24976$ $\tilde{p}_5 = 0.44421$ $\tilde{p}_6 = 0.24973$ $\tilde{p}_7 = 0.02787$ $\tilde{p}_8 = 0.00029$ $\tilde{p}_9 = 0$ $\tilde{p}_{10} = 0$	$\hat{p}_0^* = 0$ $\hat{p}_1^* = 0.00002$ $\hat{p}_2^* = 0.00028$ $\hat{p}_3^* = 0.02787$ $\hat{p}_4^* = 0.24972$ $\hat{p}_5^* = 0.44422$ $\hat{p}_6^* = 0.24972$ $\hat{p}_7^* = 0.02787$ $\hat{p}_8^* = 0.00028$ $\hat{p}_9^* = 0.00002$ $\hat{p}_{10}^* = 0$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{4} - p_0 - p_2$ $p_5 = \frac{1}{2} - 2p_1 - 2p_3$ $p_6 = \frac{1}{4} - p_0 - p_2$ $p_7 = p_3$ $p_8 = p_2$ $p_9 = p_1$ $p_{10} = p_0$

Table 5.1: Stability indexes for order n linear random homogeneous differential equations.

6 Linear random maps

In order to keep the approach of the preceding sections, we suggest to consider random linear discrete dynamical systems of the form

$$\mathcal{B} \mathbf{x}_{k+1} = A \mathbf{x}_k \text{ where } \mathbf{x} \in \mathbb{R}^n, \quad (6.1)$$

where \mathcal{B} and each of the n^2 entries of the random matrix A are i.i.d. $N(0,1)$ random variables. Observe that to ensure that the results are invariant under time-scaling, is necessary to add the term \mathcal{B} in the left-hand side of Equation (6.1). Then, given a linear discrete random system (6.1), its characteristic random polynomial associated with the matrix $\frac{1}{\mathcal{B}}A$ is

$$Q(\lambda) = Q_n \lambda^n + Q_{n-1} \lambda^{n-1} + \cdots + Q_1 \lambda + Q_0$$

where each random variable Q_j is a polynomial in the variables $1/\mathcal{B}, A_{1,1}, \dots, A_{n,n}$ which has a complicated distribution function. We denote by X the random variables that assigns to each Q its number of roots with modulus smaller than 1, that is, the stability index of the matrix $\frac{1}{\mathcal{B}}A$. Also p_k denotes the probabilities that X takes the value k .

As we will see in the examples, in this case the condition $p_k = p_{n-k}$ is no longer satisfied. Among other reasons it happens that the entries of A^{-1} have complicated distributions. Since we do not know other relations on the probabilities p_k apart from the trivial one $\sum_{k=0}^n p_k = 1$, and this is directly fulfilled by the observed relative frequencies, in this case we do not perform the least squares refinement.

The case $n = 1$ is the only one that we have been able to solve analytically. Notice that in this situation the only solution of $Q(\lambda) = 0$ is $\lambda = A/B$, with A and B independent and $N(0, 1)$. Hence $p_0 = P(|A/B| > 1)$ and $p_1 = P(|A/B| < 1) = P(|B/A| > 1)$. Since A/B and B/A have the same distribution it holds that $p_0 = p_1 = 1/2$. The results obtained for $n \leq 10$ are shown in Table 6.1.

Dimension	Observed frequency	Dimension	Observed frequency
$n = 1$	$\tilde{p}_0 = 0.49994$ $\tilde{p}_1 = 0.50006$	$n = 8$	$\tilde{p}_0 = 0.44937$ $\tilde{p}_1 = 0.24381$ $\tilde{p}_2 = 0.13702$ $\tilde{p}_3 = 0.07388$ $\tilde{p}_4 = 0.04207$ $\tilde{p}_5 = 0.02394$ $\tilde{p}_6 = 0.01526$ $\tilde{p}_7 = 0.00768$ $\tilde{p}_8 = 0.00698$
$n = 2$	$\tilde{p}_0 = 0.46348$ $\tilde{p}_1 = 0.27705$ $\tilde{p}_2 = 0.25947$	$n = 9$	$\tilde{p}_0 = 0.44941$ $\tilde{p}_1 = 0.24374$ $\tilde{p}_2 = 0.13680$ $\tilde{p}_3 = 0.07371$ $\tilde{p}_4 = 0.04139$ $\tilde{p}_5 = 0.02400$ $\tilde{p}_6 = 0.01374$ $\tilde{p}_7 = 0.00889$ $\tilde{p}_8 = 0.00434$ $\tilde{p}_9 = 0.00397$
$n = 3$	$\tilde{p}_0 = 0.45261$ $\tilde{p}_1 = 0.25828$ $\tilde{p}_2 = 0.15351$ $\tilde{p}_3 = 0.13560$	$n = 10$	$\tilde{p}_0 = 0.44934$ $\tilde{p}_1 = 0.24371$ $\tilde{p}_2 = 0.13687$ $\tilde{p}_3 = 0.07358$ $\tilde{p}_4 = 0.04129$ $\tilde{p}_5 = 0.02348$ $\tilde{p}_6 = 0.01388$ $\tilde{p}_7 = 0.00792$ $\tilde{p}_8 = 0.00520$ $\tilde{p}_9 = 0.00247$ $\tilde{p}_{10} = 0.00226$
$n = 4$	$\tilde{p}_0 = 0.45040$ $\tilde{p}_1 = 0.24732$ $\tilde{p}_2 = 0.14799$ $\tilde{p}_3 = 0.08127$ $\tilde{p}_4 = 0.07302$		
$n = 5$	$\tilde{p}_0 = 0.44957$ $\tilde{p}_1 = 0.24536$ $\tilde{p}_2 = 0.13956$ $\tilde{p}_3 = 0.08116$ $\tilde{p}_4 = 0.04443$ $\tilde{p}_5 = 0.03992$		
$n = 6$	$\tilde{p}_0 = 0.44944$ $\tilde{p}_1 = 0.24419$ $\tilde{p}_2 = 0.13838$ $\tilde{p}_3 = 0.07536$ $\tilde{p}_4 = 0.04606$ $\tilde{p}_5 = 0.02449$ $\tilde{p}_6 = 0.02209$		
$n = 7$	$\tilde{p}_0 = 0.44937$ $\tilde{p}_1 = 0.24394$ $\tilde{p}_2 = 0.13723$ $\tilde{p}_3 = 0.07480$ $\tilde{p}_4 = 0.04226$ $\tilde{p}_5 = 0.02636$ $\tilde{p}_6 = 0.01367$ $\tilde{p}_7 = 0.01236$		

Table 6.1: Stability indexes for linear random maps.

As in the other models, for each dimension $n \leq 10$, we generate 10^8 discrete systems of the form (6.1). For each matrix $\frac{1}{B}A$ we have computed the characteristic polynomial Q and its

associated polynomial Q^* (see Equation (3.2)) and have counted the number of roots of this last polynomial by using the Routh–Hurwitz zero counter. For $n \geq 5$ and in order to decrease the computation time we have directly numerically computed the eigenvalues of the matrix and counted the number of them with modulus less than one.

7 Linear random difference equations of order n

Finally we consider difference equations of order n of type

$$A_n x_{k+n} + A_{n-1} x_{k+n-1} + \cdots + A_1 x_{k+1} + A_0 x_k = 0,$$

where all the coefficients are i.i.d. random variables with $N(0,1)$ distribution. In this situation, the stability index is given by the number of zeros with modulus smaller than 1 of the random characteristic polynomial $Q(\lambda) = A_n \lambda^n + A_{n-1} \lambda^{n-1} + \cdots + A_1 \lambda + A_0$. As in the preceding sections let X be the random variable that counts the number of roots of $Q(\lambda)$ with modulus smaller than 1 and set $p_k = P(X = k)$ for $k = 0, 1, \dots, n$.

Before proving some relations among the probabilities p_k , we give two preliminary lemmas. Let $\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-u^2} du$ be the error function. The following result is stated in [4]. We prove it for the sake of completeness.

Lemma 7.1. For $\alpha > 0$ and $\beta \in \mathbb{R}$,

$$F(\alpha, \beta) := \int_0^\infty e^{-\alpha^2 x^2} \operatorname{erf}(\beta x) dx = \frac{\arctan(\beta/\alpha)}{\alpha \sqrt{\pi}}.$$

Proof. Fixed $\alpha > 0$, the function that defines F is absolutely integrable because $|\operatorname{erf}(x)| \leq 1$. Moreover its partial derivative with respect to β is also absolutely integrable. Hence $\lim_{\beta \rightarrow 0} F(\alpha, \beta) = F(\alpha, 0) = 0$ and

$$\begin{aligned} \frac{\partial F(\alpha, \beta)}{\partial \beta} &= \int_0^\infty \frac{\partial}{\partial \beta} \left(e^{-\alpha^2 x^2} \operatorname{erf}(\beta x) \right) dx = \frac{2}{\sqrt{\pi}} \int_0^\infty x e^{-\alpha^2 x^2} e^{-\beta^2 x^2} dx \\ &= \frac{2}{\sqrt{\pi}} \int_0^\infty x e^{-(\alpha^2 + \beta^2) x^2} dx = \frac{1}{(\alpha^2 + \beta^2) \sqrt{\pi}}. \end{aligned}$$

Therefore

$$F(\alpha, \beta) = F(\alpha, 0) + \int_0^\beta \frac{\partial F(\alpha, t)}{\partial t} dt = \int_0^\beta \frac{1}{(\alpha^2 + t^2) \sqrt{\pi}} dt = \frac{\arctan(\beta/\alpha)}{\alpha \sqrt{\pi}},$$

as we wanted to prove. \square

The next result is a consequence of the previous lemma.

Lemma 7.2. Let $U \sim N(0, \sigma^2)$ and $V \sim N(0, \rho^2)$ be independent normal random variables. Then $P(U^2 - V^2 > 0) = \frac{2}{\pi} \arctan(\sigma/\rho)$.

Proof. The joint density function of the random vector (U, V) is $f_\sigma(u) f_\rho(v)$, where $f_s(u) = e^{-u^2/(2s^2)} / (\sqrt{2\pi}s)$. Observe that the points $(u, v) \in \mathbb{R}^2$ such that $u^2 - v^2 > 0$ is the region where $-|u| < v < |u|$, hence by symmetry,

$$\begin{aligned} P(U^2 - V^2 > 0) &= 4 \int_0^\infty f_\sigma(u) \int_0^u f_\rho(v) dv du = \frac{4}{2\pi\sigma\rho} \int_0^\infty e^{-u^2/(2\sigma^2)} \int_0^u e^{-v^2/(2\rho^2)} dv du \\ &= \frac{2}{\pi\sigma\rho} \int_0^\infty e^{-u^2/(2\sigma^2)} \operatorname{erf}\left(\frac{u}{\sqrt{2}\rho}\right) \sqrt{\frac{\pi}{2}} \rho du = \frac{2}{\pi} \arctan\left(\frac{\sigma}{\rho}\right), \end{aligned}$$

where in the last equality we have used Lemma 7.1. \square

Notice that with the notation of the above lemma, $P(U^2 - V^2 > 0) + P(U^2 - V^2 < 0) = 1$. Hence

$$P(U^2 - V^2 < 0) = 1 - \frac{2}{\pi} \arctan\left(\frac{\sigma}{\rho}\right) = \frac{2}{\pi} \arctan\left(\frac{\rho}{\sigma}\right), \quad (7.1)$$

where we have used the fact that $\arctan(x) + \arctan(1/x) = \pi/2$ or, simply, the same lemma interchanging U and V . Observe also that when $\sigma = \rho$, $P(U^2 - V^2 > 0) = P(U^2 - V^2 < 0) = 1/2$, a result that, in fact, is a straightforward consequence that in this situation $U^2 - V^2$ and $V^2 - U^2$ have the same distribution.

Proposition 7.3. *With the above notation:*

- (a) $\sum_{k=0}^n p_k = 1$.
- (b) For all $k \in \{0, 1, \dots, n\}$, $p_k = p_{n-k}$.
- (c) When n is odd, $\sum_{i \text{ even}} p_i = \sum_{i \text{ odd}} p_i = \frac{1}{2}$.
- (d) When $n = 2k$ is even,

$$\sum_{i \text{ even}} p_i = \frac{2}{\pi} \arctan\left(\sqrt{\frac{k+1}{k}}\right) \quad \text{and} \quad \sum_{i \text{ odd}} p_i = \frac{2}{\pi} \arctan\left(\sqrt{\frac{k}{k+1}}\right). \quad (7.2)$$

Proof. The first assertion is obvious. To see the second one we compare the difference equation $a_n x_{k+n} + a_{n-1} x_{k+n-1} + \dots + a_1 x_{k+1} + a_0 x_k = 0$, with $a_i \in \mathbb{R}$, $i = 0, 1, \dots, n$, with characteristic polynomial $Q(\lambda) = a_n \lambda^n + a_{n-1} \lambda^{n-1} + \dots + a_2 \lambda^2 + a_1 \lambda + a_0$ with the difference equation $a_n x_k + a_{n-1} x_{k+1} + \dots + a_0 x_{k+n} = 0$ with characteristic polynomial $Q^*(\lambda) = a_n + a_{n-1} \lambda + \dots + a_1 \lambda^{n-1} + a_0 \lambda^n$. Notice that if $Q(\lambda)$ has m non-zero roots with modulus smaller than 1 and $n - m$ with modulus bigger than 1, then the converse follows for $Q^*(\lambda)$ because $Q(\lambda) = 0$ if and only if $Q^*(\frac{1}{\lambda}) = 0$. From this result applied to the corresponding random polynomials we get that $p_k = p_{n-k}$, because both have identically distributed coefficients. So we have proved statement (b). To prove items (c) and (d) recall first that it was proved in item (c) of Proposition 5.1 that a polynomial $Q(\lambda) = a_n \lambda^n + a_{n-1} \lambda^{n-1} + \dots + a_2 \lambda^2 + a_1 \lambda + a_0$, without roots with zero real part, has an even number of roots with negative real part if and only if $a_n a_0 > 0$. By using the polynomial

$$\begin{aligned} Q^*(z) &= a_n (z+1)^n + a_{n-1} (z+1)^{n-1} (z-1) + \dots + a_0 (z-1)^n \\ &= (a_n + a_{n-1} + \dots + a_1 + a_0) z^n + \dots + (a_n - a_{n-1} + a_{n-2} - \dots + (-1)^n a_0), \end{aligned}$$

introduced in Section 3 (Equation (3.2)) we get that $Q(\lambda)$, without roots of modulus 1, has an even number ($2m$) of roots with modulus smaller than 1 if and only if $Q^*(z)$ has exactly $2m$ roots with negative real part and this happens if and only if $(a_n + a_{n-1} + \dots + a_1 + a_0) \cdot (a_n - a_{n-1} + a_{n-2} - \dots + (-1)^n a_0) > 0$. Hence, considering the corresponding random polynomials, we have that

$$\begin{aligned} \sum_{i \text{ even}} p_i &= P((A_n + A_{n-1} + \dots + A_0) \cdot (A_n - A_{n-1} + \dots + (-1)^n A_0) > 0) \\ &= P(U^2 - V^2 > 0), \end{aligned}$$

where $U = A_n + A_{n-2} + A_{n-4} + \dots$ and $V = A_{n-1} + A_{n-3} + A_{n-5} + \dots$ and the sums end either at A_0 or A_1 according the parity of n . Since $A_j \sim N(0, 1)$ and all A_j are independent

we get that when $n = 2k$ (resp. $n = 2k - 1$) then $U \sim N(0, k + 1)$ (resp. $U \sim N(0, k)$) and $V \sim N(0, k)$ and U and V are independent. Hence, by using Lemma 7.2, we obtain that when $n = 2k - 1$, $P(U^2 - V^2 > 0) = 1/2$ and that when $n = 2k$,

$$\sum_{i \text{ even}} p_i = P(U^2 - V^2 > 0) = \frac{2}{\pi} \arctan \left(\sqrt{\frac{k+1}{k}} \right).$$

The sum of all p_i when i is odd can be obtained from the above formula, see also (7.1). \square

Corollary 7.4.

(i) Consider the linear random homogeneous difference equation of order n , let X be the random variable defined above and $p_k = P(X = k)$. Then the probabilities p_k satisfy all the conclusions of Lemma 3.1. In particular $E(X) = n/2$.

(ii) Moreover the new affine relations given in Equations (7.2) hold. In particular, for $n = 2$, $p_0 = p_2 = \frac{1}{\pi} \arctan(\sqrt{2})$ and $p_1 = \frac{2}{\pi} \arctan(1/\sqrt{2})$; and for $n = 4$, $p_1 = p_3 = \frac{1}{\pi} \arctan(\sqrt{2/3})$.

In this case, and for the situations where we have not been able to obtain the exact probabilities we have done similar computations than in the previous section, first with the Monte Carlo method, generating for each order $n = 0, \dots, 10$, 10^8 random vectors $(A_0, \dots, A_n) \in \mathbb{R}^{n+1}$ whose components are pseudo-random numbers with $N(0, 1)$ distribution. Then, by using the relations in Proposition 7.3 and Corollary 7.4 we have performed a least squares refinement.

For instance for $n = 4$, by Corollary 7.4 we have $p_1 = p_3 = \arctan(\sqrt{2/3})/\pi \simeq 0.217953$; $p_2 = 2 \arctan(\sqrt{3/2})/\pi - 2p_0$ and $p_4 = p_0$. Hence, we fix the values $\hat{p}_1 = p_1$ and $\hat{p}_3 = p_3$ and system (3.3) can be written in the form

$$\mathbf{M}\hat{\mathbf{q}} + \mathbf{b} = \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \cdot (\hat{p}_0) + \begin{pmatrix} 0 \\ \frac{2}{\pi} \arctan\left(\sqrt{\frac{3}{2}}\right) \\ 0 \end{pmatrix} = \begin{pmatrix} \tilde{p}_0 \\ \tilde{p}_2 \\ \tilde{p}_4 \end{pmatrix}.$$

Hence we can easily find the least squares solution of the above incompatible linear system:

$$(1, -2, 1) \cdot \left[\begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \cdot (\hat{p}_0) + \begin{pmatrix} 0 \\ \frac{2}{\pi} \arctan\left(\sqrt{\frac{3}{2}}\right) \\ 0 \end{pmatrix} \right] = (1, -2, 1) \cdot \begin{pmatrix} \tilde{p}_0 \\ \tilde{p}_2 \\ \tilde{p}_4 \end{pmatrix},$$

and thus we get the result in Equation (3.4): $\hat{p}_0 = \frac{1}{6} (\tilde{p}_0 - 2\tilde{p}_2 + \tilde{p}_4) + \frac{2}{3\pi} \arctan(\sqrt{3/2})$, and therefore $\hat{p}_4 = \hat{p}_0$ and $\hat{p}_2 = 2 \arctan(\sqrt{3/2})/\pi - 2\hat{p}_0$. Since our Monte Carlo simulations give

$$(\tilde{p}_0, \tilde{p}_2, \tilde{p}_4) = \left(\frac{2056203}{20000000}, \frac{7169499}{20000000}, \frac{10285619}{100000000} \right) \simeq (0.10281, 0.35847, 0.10286),$$

the above relations show that $(\hat{p}_0, \hat{p}_2, \hat{p}_4) \simeq (0.10282, 0.35846, 0.10282)$.

All our results are collected in Table 7.1.

Order	Observed frequency	Least squares	Relations (Prop. 7.3 and Cor. 7.4))
$n = 1$	$\tilde{p}_0 = 0.49991$ $\tilde{p}_1 = 0.50009$		$p_0 = 0.5$ $p_1 = 0.5$
$n = 2$	$\tilde{p}_0 = 0.30410$ $\tilde{p}_1 = 0.39184$ $\tilde{p}_2 = 0.30406$		$p_0 = \frac{1}{\pi} \arctan(\sqrt{2}) \simeq 0.304087$ $p_1 = \frac{2}{\pi} \arctan\left(\frac{1}{\sqrt{2}}\right) \simeq 0.391826$ $p_2 = \frac{1}{\pi} \arctan(\sqrt{2}) \simeq 0.304087$
$n = 3$	$\tilde{p}_0 = 0.17251$ $\tilde{p}_1 = 0.32752$ $\tilde{p}_2 = 0.32753$ $\tilde{p}_3 = 0.17244$	$\hat{p}_0 = 0.17248$ $\hat{p}_1 = 0.32752$ $\hat{p}_2 = 0.32752$ $\hat{p}_3 = 0.17248$	p_0 $p_1 = \frac{1}{2} - p_0$ $p_2 = \frac{1}{2} - p_0$ $p_3 = p_0$
$n = 4$	$\tilde{p}_0 = 0.10281$ $\tilde{p}_1 = 0.21792$ $\tilde{p}_2 = 0.35847$ $\tilde{p}_3 = 0.21794$ $\tilde{p}_4 = 0.10286$	$\hat{p}_0 = 0.10282$ $\hat{p}_1 = 0.21795$ $\hat{p}_2 = 0.35846$ $\hat{p}_3 = 0.21795$ $\hat{p}_4 = 0.10282$	p_0 $p_1 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{2}{3}}\right) \simeq 0.217953$ $p_2 = \frac{2}{\pi} \arctan\left(\sqrt{\frac{3}{2}}\right) - 2p_0$ $p_3 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{2}{3}}\right) \simeq 0.217953$ $p_4 = p_0$
$n = 5$	$\tilde{p}_0 = 0.05909$ $\tilde{p}_1 = 0.15331$ $\tilde{p}_2 = 0.28760$ $\tilde{p}_3 = 0.28756$ $\tilde{p}_4 = 0.15335$ $\tilde{p}_5 = 0.05908$	$\hat{p}_0 = 0.05909$ $\hat{p}_1 = 0.15333$ $\hat{p}_2 = 0.28758$ $\hat{p}_3 = 0.28758$ $\hat{p}_4 = 0.15333$ $\hat{p}_5 = 0.05909$	p_0 p_1 $p_2 = \frac{1}{2} - p_0 - p_1$ $p_3 = \frac{1}{2} - p_0 - p_1$ $p_4 = p_1$ $p_5 = p_0$
$n = 6$	$\tilde{p}_0 = 0.03501$ $\tilde{p}_1 = 0.09726$ $\tilde{p}_2 = 0.23777$ $\tilde{p}_3 = 0.25985$ $\tilde{p}_4 = 0.23781$ $\tilde{p}_5 = 0.09724$ $\tilde{p}_6 = 0.03505$	$\hat{p}_0 = 0.03502$ $\hat{p}_1 = 0.09726$ $\hat{p}_2 = 0.23779$ $\hat{p}_3 = 0.25986$ $\hat{p}_4 = 0.23779$ $\hat{p}_5 = 0.09726$ $\hat{p}_6 = 0.03502$	p_0 p_1 $p_2 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{4}{3}}\right) - p_0$ $p_3 = \frac{2}{\pi} \arctan\left(\sqrt{\frac{3}{4}}\right) - 2p_1$ $p_4 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{4}{3}}\right) - p_0$ $p_5 = p_1$ $p_6 = p_0$
$n = 7$	$\tilde{p}_0 = 0.02025$ $\tilde{p}_1 = 0.06432$ $\tilde{p}_2 = 0.17174$ $\tilde{p}_3 = 0.24376$ $\tilde{p}_4 = 0.24361$ $\tilde{p}_5 = 0.17177$ $\tilde{p}_6 = 0.06428$ $\tilde{p}_7 = 0.02025$	$\hat{p}_0 = 0.02025$ $\hat{p}_1 = 0.06430$ $\hat{p}_2 = 0.17176$ $\hat{p}_3 = 0.24369$ $\hat{p}_4 = 0.24369$ $\hat{p}_5 = 0.17176$ $\hat{p}_6 = 0.06430$ $\hat{p}_7 = 0.02025$	p_0 p_1 p_2 $p_3 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_4 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_5 = p_2$ $p_6 = p_1$ $p_7 = p_0$
$n = 8$	$\tilde{p}_0 = 0.01194$ $\tilde{p}_1 = 0.03994$ $\tilde{p}_2 = 0.12272$ $\tilde{p}_3 = 0.19230$ $\tilde{p}_4 = 0.25701$ $\tilde{p}_5 = 0.19238$ $\tilde{p}_6 = 0.12724$ $\tilde{p}_7 = 0.03994$ $\tilde{p}_8 = 0.01197$	$\hat{p}_0 = 0.01196$ $\hat{p}_1 = 0.03994$ $\hat{p}_2 = 0.12726$ $\hat{p}_3 = 0.19234$ $\hat{p}_4 = 0.25700$ $\hat{p}_5 = 0.19234$ $\hat{p}_6 = 0.12726$ $\hat{p}_7 = 0.03994$ $\hat{p}_8 = 0.01196$	p_0 p_1 p_2 $p_3 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{4}{5}}\right) - p_1$ $p_4 = \frac{2}{\pi} \arctan\left(\sqrt{\frac{5}{4}}\right) - 2p_0 - 2p_2$ $p_5 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{4}{5}}\right) - p_1$ $p_6 = p_2$ $p_7 = p_1$ $p_8 = p_0$

Order	Observed frequency	Least squares	Relations (Prop. 7.3 and Cor. 7.4)
$n = 9$	$\tilde{p}_0 = 0.00693$ $\tilde{p}_1 = 0.02556$ $\tilde{p}_2 = 0.08711$ $\tilde{p}_3 = 0.15653$ $\tilde{p}_4 = 0.22389$ $\tilde{p}_5 = 0.22382$ $\tilde{p}_6 = 0.15654$ $\tilde{p}_7 = 0.08712$ $\tilde{p}_8 = 0.02557$ $\tilde{p}_9 = 0.00693$	$\hat{p}_0 = 0.00693$ $\hat{p}_1 = 0.02556$ $\hat{p}_2 = 0.08711$ $\hat{p}_3 = 0.15653$ $\hat{p}_4 = 0.22386$ $\hat{p}_5 = 0.22386$ $\hat{p}_6 = 0.15653$ $\hat{p}_7 = 0.08711$ $\hat{p}_8 = 0.02556$ $\hat{p}_9 = 0.00693$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_5 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_6 = p_3$ $p_7 = p_2$ $p_8 = p_1$ $p_9 = p_0$
$n = 10$	$\tilde{p}_0 = 0.00409$ $\tilde{p}_1 = 0.01567$ $\tilde{p}_2 = 0.06089$ $\tilde{p}_3 = 0.11500$ $\tilde{p}_4 = 0.19950$ $\tilde{p}_5 = 0.20978$ $\tilde{p}_6 = 0.19941$ $\tilde{p}_7 = 0.11499$ $\tilde{p}_8 = 0.06088$ $\tilde{p}_9 = 0.01570$ $\tilde{p}_{10} = 0.00408$	$\hat{p}_0 = 0.00411$ $\hat{p}_1 = 0.01566$ $\hat{p}_2 = 0.06091$ $\hat{p}_3 = 0.11497$ $\hat{p}_4 = 0.19947$ $\hat{p}_5 = 0.20976$ $\hat{p}_6 = 0.19947$ $\hat{p}_7 = 0.11497$ $\hat{p}_8 = 0.06091$ $\hat{p}_9 = 0.01566$ $\hat{p}_{10} = 0.00411$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{6}{5}}\right) - p_0 - p_2$ $p_5 = \frac{2}{\pi} \arctan\left(\sqrt{\frac{5}{6}}\right) - 2p_1 - 2p_3$ $p_6 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{6}{5}}\right) - p_0 - p_2$ $p_7 = p_3$ $p_8 = p_2$ $p_9 = p_1$ $p_{10} = p_0$

Table 7.1: Stability indexes for order n linear random homogeneous difference equations.

Acknowledgments

We want to thank Professors Maria Jolis and Joan Torregrosa for their helpful indications. We also acknowledge the anonymous reviewer for the careful reading and for addressing to us very interesting suggestions that have allowed us to improve our work.

The authors are supported by Ministry of Science and Innovation–State Research Agency of the Spanish Government through grants PID2019-104658GB-I00 (MICINN/AEI, first and second authors) and DPI2016-77407-P (MICINN/AEI/FEDER, UE, third author). The first and second authors are also supported by the grant 2017-SGR-1617 from AGAUR, Generalitat de Catalunya. The third author acknowledges the group’s research recognition 2017-SGR-388 from AGAUR, Generalitat de Catalunya.

References

- [1] J. C. ARTÉS, J. LLIBRE, Statistical measure of quadratic vector fields, *Resenhas* 6(2003), 85–97. [MR2056449](#); [Zbl 1156.34314](#)

- [2] S. ASMUSSEN, P. W. GLYNN, *Stochastic simulation: algorithms and analysis*, Springer, New York, 2007. <https://doi.org/10.1007/978-0-387-69033-9>; MR2331321; Zbl 1126.65001
- [3] P. BRATLEY, B. L. FOX, L. E. SCHRAGE, *A guide to simulation*, 2nd ed., Springer, New York, 1987. Zbl 0515.68070
- [4] K. BRIGGS. *Integrals involving erf*, available at <http://keithbriggs.info/documents/erf-integrals.pdf> (accessed February 19, 2021).
- [5] B. M. CHEN-CHARPENTIER, D. STANESCU, Epidemic models with random coefficients, *Math. Comput. Modelling* **52**(2010), 1004–1010. <https://doi.org/10.1016/j.mcm.2010.01.014>; MR2718410; Zbl 1205.60127
- [6] B.M. CHEN-CHARPENTIER, D. STANESCU, Virus propagation with randomness, *Math. Comput. Modelling* **57**(2013), 1816–1821. <https://doi.org/10.1016/j.mcm.2011.11.065>; MR3033413; Zbl 1305.92065
- [7] A. CIMA, A. GASULL, V. MAÑOSA, Phase portraits of random planar homogeneous vector fields. *Qual. Theory Dyn. Syst.* **20**(2021), No. 1, Paper No. 3, 27 pp. <https://doi.org/10.1007/s12346-020-00437-7>; MR4195169; Zbl 07302066
- [8] K. CONRAD, *Probability distributions and maximum entropy*, available at <http://www.math.uconn.edu/~kconrad/blurbs/analysis/entropypost.pdf> (accessed February 19, 2021).
- [9] G. DAHLQUIST, Å. BJÖRCK, *Numerical methods*, Prentice-Hall, Englewood Cliffs NJ, 1974. MR0368379; Zbl 1029.65002
- [10] I. S. GRADSHTEYN, I. M. RYZHIK, *Table of Integrals, Series, and Products*, 6th ed., Academic Press, San Diego CA, 2000. <https://doi.org/10.1016/C2010-0-64839-5>; MR1773820; Zbl 0981.65001
- [11] M. GRIGORIU, T. SOONG, *Random vibration in mechanical and structural systems*, Prentice Hall, New Jersey, 1993.
- [12] F. B. JONES, Connected and disconnected plane sets and the functional equation $f(x) + f(y) = f(x + y)$, *Bull. Amer. Math. Soc.* **48**(1942), 115–120. <https://doi.org/10.1090/S0002-9904-1942-07615-4>; MR0005906; Zbl 0063.03063
- [13] C. LEMIEUX, *Monte Carlo and quasi-Monte Carlo sampling*, Springer, New York, 2009. <https://doi.org/10.1007/978-0-387-78165-5>; MR2723077; Zbl 1269.65001
- [14] H. LOPES, B. PAGNONCELLI, C. PALMEIRA, Coeficientes aleatórios de equações diferenciais ordinárias lineares (in Portuguese), *Matemática Universitária* **14**(2008), 44–50; translated to English at <http://bernardokp.uai.cl/preprint.pdf> with title *Random linear systems and simulation* (accessed February 19, 2021).
- [15] G. MARSAGLIA, Choosing a point from the surface of a sphere, *Ann. Math. Statist.* **43**(1972), 645–646. <https://doi.org/10.1214/aoms/1177692644>; Zbl 0248.65008
- [16] G. MARSAGLIA, W. TSANG, The Ziggurat method for generating random variables, *J. Stat. Softw.* **5**(2000), No. 8, 1–7. <https://doi.org/10.18637/jss.v005.i08>

- [17] M. MATSUMOTO, T. NISHIMURA, Mersenne twister: a 623-dimensionally equidistributed uniform pseudo-random number generator, *ACM Trans. Model. Comput. Simul.* **8**(1998), No. 1, 3–30. <https://doi.org/10.1145/272991.272995>; Zbl 0917.65005
- [18] B.J.T. MORGAN, *Applied stochastic modelling*, Arnold Publishers, London, 2000. MR2468192; Zbl 0985.62003
- [19] M.E. MULLER, *A note on a method for generating points uniformly on N-dimensional spheres*, *Commun. ACM* **2**(1959), 19–20. <https://doi.org/10.1145/377939.377946>; Zbl 0086.11605
- [20] J. R. SCHOTT, *Matrix analysis for statistics*, John Wiley & Sons, New York, 1997. MR1421574; Zbl 0872.15002
- [21] D. STANESCU, B. M. CHEN-CHARPENTIER, Random coefficient differential equation models for bacterial growth, *Math. Comput. Modelling* **50**(2009), 885–895. <https://doi.org/10.1016/j.mcm.2009.05.017>; MR2569250; Zbl 1185.34075
- [22] J. STOER, R. BULIRSCH, *Introduction to numerical analysis*, 3rd ed. Springer, New York, 2002. <https://doi.org/10.1214/aoms/1177692644>; MR1923481; Zbl 1004.65001
- [23] S. H. STROGATZ, *Nonlinear dynamics and chaos*, Westview Press, Cambridge MA, 1994. <https://doi.org/10.1201/9780429492563>; MR3837141; Zbl 1343.37001