

MAXIMAL FLOW THROUGH A NETWORK

L. R. FORD, JR. AND D. R. FULKERSON

Introduction. The problem discussed in this paper was formulated by T. Harris as follows:

“Consider a rail network connecting two cities by way of a number of intermediate cities, where each link of the network has a number assigned to it representing its capacity. Assuming a steady state condition, find a maximal flow from one given city to the other.”

While this can be set up as a linear programming problem with as many equations as there are cities in the network, and hence can be solved by the simplex method (1), it turns out that in the cases of most practical interest, where the network is planar in a certain restricted sense, a much simpler and more efficient hand computing procedure can be described.

In §1 we prove the minimal cut theorem, which establishes that an obvious upper bound for flows over an arbitrary network can always be achieved. The proof is non-constructive. However, by specializing the network (§2), we obtain as a consequence of the minimal cut theorem an effective computational scheme. Finally, we observe in §3 the duality between the capacity problem and that of finding the shortest path, via a network, between two given points.

1. The minimal cut theorem. A *graph* G is a finite, 1-dimensional complex, composed of *vertices* a, b, c, \dots, e , and *arcs* $\alpha(ab), \beta(ac), \dots, \delta(ce)$. An arc $\alpha(ab)$ joins its end vertices a, b ; it passes through no other vertices of G and intersects other arcs only in vertices. A *chain* is a set of distinct arcs of G which can be arranged as $\alpha(ab), \beta(bc), \gamma(cd), \dots, \delta(gh)$, where the vertices a, b, c, \dots, h are distinct, i.e., a chain does not intersect itself; a chain joins its end vertices a and h .

We distinguish two vertices of G : a , the *source*, and b , the *sink*.¹ A *chain flow* from a to b is a couple $(C; k)$ composed of a chain C joining a and b , and a non-negative number k representing the flow along C from source to sink.

Each arc in G has associated with it a positive number called its *capacity*. We call the graph G , together with the capacities of its individual arcs, a *network*. A *flow* in a network is a collection of chain flows which has the property that the sum of the numbers of all chain flows that contain any arc is no greater than the capacity of that arc. If equality holds, we say the arc is *saturated* by the flow. A chain is *saturated* with respect to a flow if it contains

Received September 20, 1955.

¹The case in which there are many sources and sinks with shipment permitted from any source to any sink is obviously reducible to this.

a saturated arc. The value of a flow is the sum of the numbers of all the chain flows which compose it.

It is clear that the above definition of flow is not broad enough to include everything that one intuitively wishes to think of as a flow, for example, sending trains out a dead end and back or around a circuit, but as far as effective transportation is concerned, the definition given suffices.

A *disconnecting set* is a collection of arcs which has the property that every chain joining a and b meets the collection. A disconnecting set, no proper subset of which is disconnecting, is a *cut*. The *value* of a disconnecting set D (written $v(D)$) is the sum of the capacities of its individual members. Thus a disconnecting set of minimal value is automatically a cut.

THEOREM 1. (Minimal cut theorem). *The maximal flow value obtainable in a network N is the minimum of $v(D)$ taken over all disconnecting sets D .*

Proof. There are only finitely many chains joining a and b , say n of them. If we associate with each one a coordinate in n -space, then a flow can be represented by a point whose j th coordinate is the number attached to the chain flow along the j th chain. With this representation, the class of all flows is a closed, convex polytope in n -space, and the value of a flow is a linear functional on this polytope. Hence, there is a maximal flow, and the set of all maximal flows is convex.

Now let S be the class of all arcs which are saturated in every maximal flow.

LEMMA 1. *S is a disconnecting set.*

Suppose not. Then there exists a chain $\alpha_1, \alpha_2, \dots, \alpha_m$ joining a and b with $\alpha_i \notin S$ for each i . Hence, corresponding to each α_i , there is a maximal flow f_i in which α_i is unsaturated. But the average of these flows,

$$f = \frac{1}{m} \sum f_i,$$

is maximal and α_i is unsaturated by f for each i . Thus the value of f may be increased by imposing a larger chain flow on $\alpha_1, \alpha_2, \dots, \alpha_m$, contradicting maximality.

Notice that the orientation assigned to an arc of S by a positive chain flow of a maximal flow is the same for all such chain flows. For suppose first that $(C_1, k_1), (C_2, k_2)$ are two chain flows occurring in a maximal flow $f, k_1 \geq k_2 > 0$, where

$$C_1 = \alpha_1(a a_1), \alpha_2(a_1 a_2), \dots, \alpha_j(a_{j-1}, a_j), \dots, \alpha_r(a_{r-1}, b)$$

$$C_2 = \beta_1(a b_1), \beta_2(b_1 b_2), \dots, \beta_k(b_{k-1}, b_k), \dots, \beta_s(b_{s-1}, b),$$

and $\alpha_j(a_{j-1}, a_j) = \beta_k(b_{k-1}, b_k) \in S, a_{j-1} = b_k, a_j = b_{k-1}$. Then

$$C'_1 = \alpha_1, \alpha_2, \dots, \alpha_{j-1}, \beta_{k+1}, \dots, \beta_s$$

$$C'_2 = \beta_1, \beta_2, \dots, \beta_{k-1}, \alpha_{j+1}, \dots, \alpha_r$$

contain chains C_1'', C_2'' joining a and b , and another maximal flow can be obtained from f as follows. Reduce the C_1 and C_2 components of f each by k_2 , and increase each of the C_1'' and C_2'' components by k_2 . This unsaturates the arc α_j , contradicting its definition as an element of S . On the other hand, if $(C_1, k_1), (C_2, k_2)$ were members of distinct maximal flows f_1, f_2 , consideration of $f = \frac{1}{2}(f_1 + f_2)$ brings us back to the former case. Hence, the arcs of S have a definite orientation assigned to them by maximal flows. We refer to that vertex of an arc $\alpha \in S$ which occurs first in a positive chain flow of a maximal flow as the *left vertex* of α .

Now define a *left arc* of S as follows: an arc α of S is a left arc if and only if there is a maximal flow f and a chain $\alpha_1, \alpha_2, \dots, \alpha_k$ (possibly null) joining a and the left vertex of α with no α_i saturated by f . Let L be the set of left arcs of S .

LEMMA 2. L is a disconnecting set.

Given an arbitrary chain $\alpha_1(a a_1), \alpha_2(a_1 a_2), \dots, \alpha_m(a_{m-1} b)$ joining a and b , it must intersect S by Lemma 1. Let $\alpha_t(a_{t-1}, a_t)$ be the first $\alpha_i \in S$. Then for each $\alpha_i, i < t$, there is a maximal flow f_i in which α_i is unsaturated. The average of these flows provides a maximal flow f in which $\alpha_1, \alpha_2, \dots, \alpha_{t-1}$ are unsaturated. It remains to show that this chain joins a to the left vertex of α_t , i.e., a_{t-1} is the left vertex of α_t . Suppose not. Then the maximal flow f contains a chain flow

$$[\beta_1(ab_1), \beta_2(b_1, b_2), \dots, \beta_r(b_{r-1}, b); k], \quad k > 0, \quad \beta_s = \alpha_t, \quad b_{s-1} = a_t, \quad b_s = a_{t-1}.$$

Let the amount of unsaturation in f of $\alpha_i (i = 1, \dots, t - 1)$, be $k_i > 0$. Now alter f as follows: decrease the flow along the chain $\beta_1, \beta_2, \dots, \beta_r$ by $\min [k, k_i] > 0$ and increase the flow along the chain contained in

$$\alpha_1, \alpha_2, \dots, \alpha_{t-1}, \beta_{s+1}, \dots, \beta_r$$

by this amount. The result is a maximal flow in which α_t is unsaturated, a contradiction. Hence $\alpha_t \in L$.

LEMMA 3. No positive chain flow of a maximal flow can contain more than one arc of L .

Assume the contrary, that is, there is a maximal flow f_1 containing a chain flow

$$[\beta_1(ab_1), \beta_2(b_1 b_2), \dots, \beta_r(b_{r-1}, b); k], \quad k > 0,$$

with arcs $\beta_i, \beta_j \in L, \beta_i$ occurring before β_j , say, in the chain. Let f_2 be that maximal flow for which there is an unsaturated chain

$$\alpha_1(a a_1), \alpha_2(a_1, a_2), \dots, \alpha_s(a_{s-1}, b_{j-1})$$

from a to the left vertex of β_j . Consider $f = \frac{1}{2}(f_1 + f_2)$. This maximal flow contains the chain flow $[\beta_1, \beta_2, \dots, \beta_r; k']$ with $k' \geq \frac{1}{2}k$, and each $\alpha_i (i = 1, \dots, s)$ is unsaturated by $k_i > 0$ in f . Again alter f : decrease the flow along

$\beta_1, \beta_2, \dots, \beta_r$ by $\min [k', k_i] > 0$ and increase the flow along the chain contained in $\alpha_1, \alpha_2, \dots, \alpha_s, \beta_j, \dots, \beta_r$ by the same amount, obtaining a maximal flow in which β_i is unsaturated, a contradiction.

Now to prove the theorem it suffices only to remark that the value of every flow is no greater than $v(D)$ where D is any disconnecting set; and on the other hand we see from Lemma 3 and the definition of S that in adding the capacities of arcs of L we have counted each chain flow of a maximal flow just once. Since by Lemma 2 L is a disconnecting set, we have the reverse inequality. Thus L is a minimal cut and the value of a maximal flow is $v(L)$.

We shall refer to the value of a maximal flow through a network N as the *capacity* of N ($\text{cap}(N)$). Then note the following corollary of the minimal cut theorem.

COROLLARY. *Let A be a collection of arcs of a network N which meets each cut of N in just one arc. If N' is a network obtained from N by adding k to the capacity of each arc of A , then $\text{cap}(N') = \text{cap}(N) + k$.*

It is worth pointing out that the minimal cut theorem is not true for networks with several sources and corresponding sinks, where shipment is restricted to be from a source to its sink. For example, in the network (Fig. 1) with shipment from a_i to b_i and capacities as indicated, the value of a minimal disconnecting set (i.e., a set of arcs meeting all chains joining sources and corresponding sinks) is 4, but the value of a maximal flow is 3.




Fig. 1

2. A computing procedure for source-sink planar networks.² We say that a network N is *planar* with respect to its source and sink, or briefly, N is *ab-planar*, provided the graph G of N , together with arc ab , is a planar

²It was conjectured by G. Dantzig, before a proof of the minimal cut theorem was obtained, that the computing procedure described in this section would lead to a maximal flow for planar networks.

graph (2; 3). (For convenience, we suppose there is no arc in G joining a and b .) The importance of ab -planar networks lies in the following theorem.

THEOREM 2. *If N is ab -planar, there exists a chain joining a and b which meets each cut of N precisely once.*

Proof. We may assume, without loss of generality, that the arc ab is part of the boundary of the outside region, and that G lies in a vertical strip with a located on the left bounding line of the strip, b on the right. Let T be the chain joining a and b which is top-most in N . T has the desired property, as we now show. Suppose not. Then there is a cut D , at least two arcs of which are in T . Let these be α_1 and α_2 , with α_1 occurring before α_2 in following T from a to b . Since D is a cut, there is a chain C_1 joining a and b which meets D in α_1 only. Similarly there is a chain C_2 meeting D in α_2 only. Let C_2' be that part of C_2 joining a to an end point of α_2 . It follows from the definition of T that C_1 and C_2' must intersect. But now, starting at a , follow C_2' to its last intersection with C_1 , then C_1 to b . We thus have a chain from a to b not meeting D , contradicting the fact that D is a cut.

Symmetrically, of course, the bottom-most chain of N has the same property.

Notice that this theorem is not valid for networks which are not ab -planar. A simple example showing this is provided by the "gas, water, electricity" graph (Fig. 2), in which every chain joining a and b meets some cut in three arcs.




Fig. 2

Theorem 2 and the corollary to Theorem 1 provide an easy computational procedure for determining a maximal flow in a network of the kind here considered. Simply locate a chain having the property of Theorem 2; this can be done at a glance by finding the two regions separated by arc ab , and taking the rest of the boundary of either region (throwing out portions of the boundary where it has looped back and intersected itself, so as to get a chain). Impose as large a chain flow ($T; k$) as possible on this chain, thereby saturating one or more of its arcs. By the corollary, subtracting k from each capacity in T reduces the capacity of N by k . Delete the saturated arcs, and proceed as

before. Eventually, the graph disconnects, and a maximal flow has been constructed.

3. A minimal path problem. For source-sink planar networks, there is an interesting duality between the problem of finding a chain of minimal capacity-sum joining source and sink and the network capacity problem, which lies in the fact that chains of N joining source and sink correspond to cuts (relative to two particular vertices) of the dual³ of N and vice versa. More precisely, suppose one has a network N , planar relative to two vertices a and b , and wishes to find a chain joining a and b such that the sum of the numbers assigned to the arcs of the chain is minimal. An easy way to solve this problem is as follows. Add the arc ab , and construct the dual of the resulting graph G . Let a' and b' be the vertices of the dual which lie in the regions of G separated by ab . Assign each number of the original network to the corresponding arc in the dual. Then solve the capacity problem relative to a' and b' for the dual network by the procedure of §2. A minimal cut thus constructed corresponds to a minimal chain in the original network.

³The *dual* of a planar graph G is formed by taking a vertex inside each region of G and connecting vertices which lie in adjacent regions by arcs. See (2; 3).

REFERENCES

1. G. B. Dantzig, *Maximization of a linear function of variables subject to linear inequalities: Activity analysis of production and allocation* (Cowles Commission, 1951).
2. H. Whitney, *Non-separable and planar graphs*, Trans. Amer. Math. Soc., *34* (1932), 339-362.
3. ———, *Planar graphs*, Fundamenta Mathematicae, *21* (1933), 73-84.

Rand Corporation,
Santi Monica, California