



The roots of the independence polynomial of a clawfree graph

Maria Chudnovsky¹, Paul Seymour²

Princeton University, Princeton, NJ 08544, USA

Received 2 February 2005

Available online 5 July 2006

Abstract

The *independence polynomial* of a graph G is the polynomial $\sum_A x^{|A|}$, summed over all independent subsets $A \subseteq V(G)$. We prove that if G is clawfree, then all the roots of its independence polynomial are real. This extends a theorem of Heilmann and Lieb [O.J. Heilmann, E.H. Lieb, Theory of monomer–dimer systems, *Comm. Math. Phys.* 25 (1972) 190–232], answering a question posed by Hamidoune [Y.O. Hamidoune, On the numbers of independent k -sets in a clawfree graph, *J. Combin. Theory Ser. B* 50 (1990) 241–244] and Stanley [R.P. Stanley, Graph colorings and related symmetric functions: Ideas and applications, *Discrete Math.* 193 (1998) 267–286].

© 2006 Elsevier Inc. All rights reserved.

Keywords: Clawfree graphs; Roots; Independence polynomial

1. Introduction

All graphs in this paper are finite and simple. An *independent set* in a graph is a set of pairwise non-adjacent vertices. The *independence polynomial* of a graph G is the polynomial

$$I(G, x) = \sum_A x^{|A|},$$

where the sum is over all independent subsets $A \subseteq V(G)$. (See for instance [2–5,8–10,13] for work on these polynomials.)

E-mail addresses: mchudnov@math.princeton.edu (M. Chudnovsky), pds@math.princeton.edu (P. Seymour).

¹ This research was conducted during the period the author served as a Clay Mathematics Institute Research Fellow.

² Partially supported by ONR Grant N00014-01-1-0608, and NSF Grant DMS-0070912.

Given a graph H , its *line graph* $L(H)$ is the graph whose vertex set is the set of edges of H , and two vertices are adjacent if they share an end in H . In [12] Heilmann and Lieb proved that if G is a line graph, then all the roots of $I(G, x)$ are real. This property does not hold for all graphs, since the independence polynomial of a claw is

$$1 + 4x + 3x^2 + x^3$$

and not all its roots are real (a *claw* is the graph with vertex set $\{v_1, v_2, v_3, v_4\}$ and three edges v_1v_2, v_1v_3, v_1v_4).

A graph G is said to be *clawfree* if no induced subgraph of it is a claw. The main result of this paper is the following, answering a question of Hamidoune [11] that was later posed as a conjecture by Stanley [15].

1.1. *If G is clawfree then all roots of $I(G, x)$ are real.*

Since all line graphs are clawfree, this extends the result of [12].

2. Proof of the main result

Let $f_1(x), \dots, f_k(x)$ be polynomials in one variable with real coefficients. We say they are *compatible* if for all $c_1, \dots, c_k \geq 0$, all the roots of the polynomial $\sum_{i=1}^k c_i f_i(x)$ are real.

For $v \in V(G)$ we denote by $N[v]$ the union of $\{v\}$ and the set of neighbours of v in G , and for an induced subgraph H of G let $N_H[v] = N[v] \cap V(H)$. For $A \subseteq V(G)$ we denote by $G \setminus A$ the subgraph of G induced on $V(G) \setminus A$. Our proof of 1.1 is essentially by induction on the number of vertices, making use of the following fundamental identity, the proof of which is clear:

2.1. *If v is a vertex of a graph G , then $I(G, x) = I(G \setminus \{v\}, x) + xI(G \setminus N[v], x)$.*

When we apply this, there is a problem; while we would know inductively that all roots of $G \setminus \{v\}$ and $G \setminus N[v]$ are real, how can we deduce that all roots of G are real? It is not true that the sum of two polynomials with all roots real again has all roots real. The trick is to prove a stronger statement inductively, that certain pairs of polynomials are compatible. Thus we would know inductively that all pairs of the appropriate polynomials in the smaller graph are compatible; but it seems that we have not gained anything, because when we try to apply 2.1, we find that we need to know that all *quadruples* of these polynomials are compatible. But there is a general lemma that if all pairs of certain polynomials are compatible then so are all larger subsets of them, as we explain next.

If $f_1(x), \dots, f_k(x)$ are as before, we say they are *pairwise compatible* if for all $i, j \in \{1, \dots, k\}$, the polynomials $f_i(x)$ and $f_j(x)$ are compatible. In this section we assume the truth of the following lemma, which we prove in Section 3. (This lemma is new as far as we know, but is closely related to a result of Bartlett, Hollot and Lin [1].)

2.2. *Let $f_1(x), \dots, f_k(x)$ be pairwise compatible polynomials with positive leading coefficients. Then $f_1(x), \dots, f_k(x)$ are compatible.*

We start with some lemmas. A *clique* in G is a set of vertices all pairwise adjacent. The following is a useful extension of 2.1 (again the proof is clear).

2.3. Let K be a clique of a graph G . Then

$$I(G, x) = I(G \setminus K, x) + \sum_{k \in K} xI(G \setminus N[k], x).$$

A clique K is *simplicial* if for every $k \in K$ the set $N[k] \setminus K$ is a clique.

2.4. Let G be a clawfree graph and let K be a simplicial clique in G . Then $N[k] \setminus K$ is a simplicial clique in $G \setminus K$ for all $k \in K$.

Proof. Since K is a simplicial clique, $N[k] \setminus K$ is a clique for all $k \in K$. Suppose for some $k \in K$ and some $n \in N[k] \setminus K$, the set $N[n] \setminus (K \cup N[k])$ is not a clique. Let x, y be two non-adjacent vertices in $N[n] \setminus (K \cup N[k])$. Then $\{n, k, x, y\}$ induces a claw in G , a contradiction. This proves 2.4. \square

Let us say a graph G is *real-rooted* if for every induced subgraph H of G , all roots of $I(H, x)$ are real. We wish to prove that every clawfree graph is real-rooted, and we need some lemmas about real-rooted graphs.

2.5. Let G be a real-rooted clawfree graph. Then:

- (1) for every two simplicial cliques K, L in G , the polynomials $I(G \setminus K, x)$ and $I(G \setminus L, x)$ are compatible,
- (2) for every simplicial clique K , the polynomials $I(G, x)$ and $xI(G \setminus K, x)$ are compatible.

Proof. We prove both statements simultaneously by induction on $|V(G)|$. First we prove 2.5(1). Since G is real-rooted, we may assume that $K \cup L \neq \emptyset$. Write $H = G \setminus (K \cup L)$. By 2.3 applied to $G \setminus L$ and $K \setminus L$, we have

$$I(G \setminus L, x) = I(H, x) + \sum_{v \in K \setminus L} xI(H \setminus N_H[v], x)$$

and similarly

$$I(G \setminus K, x) = I(H, x) + \sum_{v \in L \setminus K} xI(H \setminus N_H[v], x).$$

By 2.2, in order to prove that $I(G \setminus K, x)$ and $I(G \setminus L, x)$ are compatible, it is enough to show that the polynomials

$$I(H, x), \quad xI(H \setminus N_H[v], x), \quad xI(H \setminus N_H[u], x)$$

are pairwise compatible for all $u, v \in K \cup L$, since they all have positive leading coefficients. Since $|V(H)| < |V(G)|$ and by 2.4, it follows inductively from 2.5(2) that $I(H, x)$ is compatible with $xI(H \setminus N_H[v], x)$ for $v \in K \cup L$; and from 2.5(1) that $xI(H \setminus N_H[u], x)$ and $xI(H \setminus N_H[v], x)$ are compatible for all $u, v \in K \cup L$. This proves 2.5(1).

To prove 2.5(2), let K be a simplicial clique in G . Since G is real-rooted, we may assume that $K \neq \emptyset$. By 2.3,

$$I(G, x) = I(G \setminus K, x) + x \sum_{k \in K} I(G \setminus N[k], x).$$

By 2.2, to prove that $I(G, x)$ and $xI(G \setminus K, x)$ are compatible, it is enough to prove that the polynomials

$$xI(G \setminus K, x), \quad I(G \setminus K, x), \quad xI(G \setminus N[k], x), \quad xI(G \setminus N[k'], x)$$

are pairwise compatible for all $k, k' \in K$, since they all have positive leading coefficients.

Since G is real-rooted, it follows that the roots of $I(G \setminus K, x)$ are real, so $xI(G \setminus K, x)$ and $I(G \setminus K, x)$ are compatible. By 2.4 and 2.5(2) (applied inductively to $G \setminus K$), $I(G \setminus K, x)$ and $xI(G \setminus N[k], x)$ are compatible for all $k \in K$, and by 2.5(1) with $L = \emptyset$, $xI(G \setminus K, x)$ and $xI(G \setminus N[k], x)$ are compatible for all $k \in K$. Inductively by 2.5(1), $xI(G \setminus N[k], x)$ and $xI(G \setminus N[k'], x)$ are compatible for all $k, k' \in K$. This proves that

$$xI(G \setminus K, x), \quad I(G \setminus K, x), \quad xI(G \setminus N[k], x), \quad xI(G \setminus N[k'], x)$$

are pairwise compatible for all $k, k' \in K$, and so completes the proof of 2.5(2) and therefore of 2.5. \square

We use 2.5 to prove the following.

2.6. *Let G be a clawfree graph, and let $v \in V(G)$ such that $G \setminus v$ is real-rooted. Then the polynomials $I(G \setminus v, x)$ and $xI(G \setminus N[v], x)$ are compatible.*

Proof. The proof is by induction on $|V(G)|$. If v has no neighbours, then by 2.1 $I(G \setminus N[v], x) = I(G \setminus v, x)$, and since $G \setminus v$ is real-rooted, it follows that $I(G \setminus v, x)$ and $xI(G \setminus N[v], x)$ are compatible. We may therefore assume that there is a vertex u adjacent to v . Let $H = G \setminus (N[u] \cap N[v])$.

(1) $N_H[u]$ and $N_H[v]$ are both simplicial cliques in H .

For $N_H[u]$ is a clique, since $\{u, v, x, y\}$ does not induce a claw for $x, y \in N_H[u]$. Suppose that for some $w \in N_H[u]$ there exist two non-adjacent vertices $s, t \in N_H[w] \setminus N_H[u]$. Then $\{w, u, s, t\}$ induces a claw in G , a contradiction. So $N_H[u]$, and similarly $N_H[v]$, is a simplicial clique in H . This proves (1). By 2.1 applied to $G \setminus \{v\}$,

$$I(G \setminus \{v\}, x) = I(G \setminus \{u, v\}, x) + xI(G \setminus N[u], x),$$

so to complete the proof of the theorem, by 2.2 it is enough to show that the polynomials

$$xI(G \setminus N[v], x), \quad I(G \setminus \{u, v\}, x), \quad xI(G \setminus N[u], x)$$

are pairwise compatible. But the pairs $xI(G \setminus N[v], x)$, $I(G \setminus \{u, v\}, x)$ and $xI(G \setminus N[u], x)$, $I(G \setminus \{u, v\}, x)$ are compatible by the inductive hypothesis applied, respectively, to the graphs $G \setminus \{u\}$ and $G \setminus \{v\}$; and $xI(G \setminus N[u], x)$, $xI(G \setminus N[v], x)$ are compatible by (1) and the first assertion of 2.5, since $I(G \setminus N[u], x) = I(H \setminus N_H[u], x)$ and $I(G \setminus N[v], x) = I(H \setminus N_H[v], x)$. This completes the proof of 2.6. \square

Proof of 1.1. We proceed by induction on $|V(G)|$. We may assume that there exists $v \in V(G)$; and from the inductive hypothesis, $G \setminus \{v\}$ is real-rooted. By 2.6, $I(G \setminus \{v\}, x)$, $xI(G \setminus N[v], x)$ are compatible; and since $I(G, x) = I(G \setminus \{v\}, x) + xI(G \setminus N[v], x)$ by 2.1, it follows that all roots of $I(G, x)$ are real. This proves 1.1. \square

3. Proof of the lemma

It remains, therefore, to prove 2.2, and that is the goal of this section. We start by proving some general facts about compatible pairs of polynomials.

3.1. *If the polynomials f and g are compatible then so are their derivatives.*

Proof. We note that between every two real roots of a polynomial h there is a root of its derivative h' (counting according to multiplicity), and so if all roots of h are real then the same holds for h' . Thus the fact that f and g are compatible implies that their derivatives are compatible. This proves 3.1. \square

If f is a polynomial, we denote its degree by $\deg(f)$.

3.2. *If f, g are compatible polynomials with positive leading coefficients then $|\deg(f) - \deg(g)| \leq 1$.*

Proof. The proof is by induction on $\max\{\deg(f), \deg(g)\}$. Assume first that one of f, g is a constant function, say $f(x) = c$. Since the leading coefficient of f is positive, it follows that $c > 0$. Then adding a large enough multiple of f to g produces a polynomial with at most one real root (since the leading coefficient of g is positive). Since the pair f, g is compatible it follows that $\deg(g) \leq 1$ and the result holds.

So we may assume that both f and g have degree at least 1. By 3.1 the derivatives f' and g' of f and g respectively are compatible. Both f' and g' have positive leading coefficients and it follows inductively that

$$|\deg(f) - \deg(g)| = |\deg(f') - \deg(g')| \leq 1.$$

This proves 3.2. \square

When $f(x)$ is a polynomial, we denote by $n_f(x)$ the number of real roots of $f(x)$ that lie in the interval $[x, \infty)$ (counted with their multiplicities). We say that f and g agree at $a \in \mathfrak{R}$ if $f(a)$ and $g(a)$ are either both positive or both negative (and both non-zero).

3.3. *If f and g are compatible polynomials that agree at a and b for some $a < b \in \mathfrak{R}$, then*

$$n_f(b) - n_f(a) = n_g(b) - n_g(a).$$

Proof. For $0 \leq t \leq 1$ let $p_t(x) = tf(x) + (1-t)g(x)$. Since f and g agree at a and b , for all t the polynomial $p_t(x)$ has no roots on the boundary of the interval $[a, b]$. For $0 \leq t \leq 1$ the roots of p_t move continuously with t in the complex plane, and therefore in the real line, since all the roots of p_t are real because f and g are compatible. In particular, the number of roots of $p_t(x)$ in the open interval (a, b) is independent of t . But the polynomials $p_0(x)$ and $p_1(x)$ have $n_g(b) - n_g(a)$ and $n_f(b) - n_f(a)$ roots respectively in this interval. This proves 3.3. \square

3.4. *Let f, g be compatible polynomials with positive leading coefficients. Then $|n_f(x) - n_g(x)| \leq 1$ for all $x \in \mathfrak{R}$.*

Proof. The proof is by induction on $\max\{\deg(f), \deg(g)\}$. Since the common roots of f and g contribute the same amount to n_f and n_g we may assume that f, g have no common roots by factoring out the greatest common divisor (the compatibility property is preserved). Suppose that $n_f(x_0) - n_g(x_0) \geq 2$ for some $x_0 \in \mathfrak{R}$. We may assume that x_0 is a root of f , and indeed x_0 is the largest root such that $n_f(x_0) - n_g(x_0) \geq 2$. Since x_0 is a root of f , it is not a root of g .

First we claim that $n_f(x_0) - n_g(x_0) = 2$. Assume for a contradiction that $n_f(x_0) - n_g(x_0) \geq 3$. Since between every two real roots of a polynomial there is a real root of its derivative, it follows that $n_{f'}(x_0) = n_f(x_0) - 1$ and $n_{g'}(x_0) \leq n_g(x_0)$. Thus

$$n_{f'}(x_0) - n_{g'}(x_0) \geq n_f(x_0) - n_g(x_0) - 1 \geq 2,$$

contrary to the inductive hypothesis. This proves that $n_f(x_0) - n_g(x_0) = 2$.

Choose y_1 strictly larger than all roots of f and all roots of g . Since f and g both have positive leading coefficients, it follows that f and g agree at y_1 . For the same reason and since $n_f(x_0) - n_g(x_0)$ is even, we can choose $y_2 < x_0$ such that f and g agree at y_2 and have no roots in the interval $[y_2, x_0)$. But now

$$n_f(y_2) - n_f(y_1) \neq n_g(y_2) - n_g(y_1),$$

contrary to 3.3. This proves 3.4. \square

For two monotone non-increasing sequences (a_1, \dots, a_m) and (b_1, \dots, b_n) of real numbers, we say that the first *interleaves* the second if $n \leq m \leq n + 1$ and $(a_1, b_1, a_2, b_2, \dots)$ is another monotone non-increasing sequence. (This does not imply that the second sequence interleaves the first.)

If f is a polynomial of degree d with all roots real, let $r_1 \geq \dots \geq r_d$ be the roots of f . We call the sequence (r_1, \dots, r_d) the *root sequence* of f . Let $f_1(x), \dots, f_k(x)$ be polynomials with positive leading coefficients and all roots real. A *common interleaver* for $f_1(x), \dots, f_k(x)$ is a sequence that interleaves the root sequence of each f_i .

We observe the following (the proof is easy and we leave it to the reader).

3.5. Let $f(x), g(x)$ be polynomials with all roots real. They have a common interleaver if and only if $|n_f(x) - n_g(x)| \leq 1$ for all $x \in \mathfrak{R}$.

We now complete the proof of 2.2, by proving it in the strengthened form below. If $k = 2$ and f_1, f_2 have the same degree, this was essentially proved by Dedieu [6,7] (thanks to Alan Sokal for bringing this reference to our attention).

3.6. Let $f_1(x), \dots, f_k(x)$ be polynomials with positive leading coefficients and all roots real. The following four statements are equivalent:

- (1) f_1, \dots, f_k are pairwise compatible,
- (2) for all s, t such that $1 \leq s < t \leq k$, the polynomials f_s, f_t have a common interleaver,
- (3) f_1, \dots, f_k have a common interleaver,
- (4) f_1, \dots, f_k are compatible.

Proof. For $1 \leq i \leq k$ let $d_i = \deg(f_i)$ and let $d = \max_{1 \leq i \leq k} d_i$. Let $(r_1^i, \dots, r_{d_i}^i)$ be the root sequence of f_i . If $d_i \geq 1$, let $I_1^i, \dots, I_{d_i+1}^i$ be the intervals of \mathfrak{R} defined as follows: $I_1^i = [r_1^i, \infty)$, $I_{d_i+1}^i = (-\infty, r_{d_i}^i]$ and $I_j^i = [r_j^i, r_{j-1}^i]$ for $2 \leq j \leq d_i$. If $d_i = 0$ we set $I_1^i = \mathfrak{R}$. We note

that a sequence (p_1, \dots, p_m) interleaves the root sequence of f_i if and only if $d_i - 1 \leq m \leq d_i$ and $p_j \in I_j^i$ for $i \leq j \leq m$.

3.6(1) implies 3.6(2). For let $1 \leq s < t \leq k$ and let $d^* = \min\{d_s, d_t\}$. By 3.2 $d^* \geq \max\{d_s, d_t\} - 1$. It is enough to prove that for $1 \leq j \leq d^* + 1$ the intersection $I_j^s \cap I_j^t$ is non-empty. Suppose not, and let j be minimum such that $I_j^s \cap I_j^t = \emptyset$. Since the leading coefficients of both f_s and f_t are positive, $j \geq 2$. From the symmetry we may assume that $r_{j-1}^s \leq r_{j-1}^t$, and so r_j^t exists and $r_{j-1}^s < r_j^t$. But then $n_{f_t}(r_j^t) = j$ and $n_{f_s}(r_j^t) \leq j - 2$, contrary to 3.4. This proves that 3.6(1) implies 3.6(2).

3.6(2) implies 3.6(3). For it follows from 3.6(2) that for all $s, t \in \{1, \dots, k\}$ and all $j \in \{1, \dots, d\}$, the intersection $I_j^s \cap I_j^t$ is non-empty. From the Helly property of linear intervals we deduce that $\bigcap_{i=1}^k I_j^i \neq \emptyset$ for all $j \in \{1, \dots, d\}$. For $1 \leq j \leq d$ choose $p_j \in \bigcap_{i=1}^k I_j^i$; then (p_1, \dots, p_d) is a common interleaver for f_1, \dots, f_k . This proves that 3.6(2) implies 3.6(3).

3.6(3) implies 3.6(4). The proof is by induction on d . We may assume that no $x_0 \in \mathfrak{R}$ is a root of all f_1, \dots, f_k , for otherwise the theorem follows from the inductive hypothesis applied to the family $f_i/(x - x_0)$ ($1 \leq i \leq k$), which still has a common interleaver by 3.5. Let c_1, \dots, c_k be non-negative real numbers, and let $f = \sum_{i=1}^k c_i f_i$. We need to prove that all the roots of f are real. We may assume that c_1, \dots, c_k are all positive.

Since f_1, \dots, f_k have a common interleaver, it follows that $d - 1 \leq d_i \leq d$ for all $1 \leq i \leq k$, and so we may assume that there is a common interleaver with d terms, say (p_1, \dots, p_d) .

Let $1 \leq i \leq k$. Since the leading coefficient of f_i is positive and (p_1, \dots, p_d) interleaves the root sequence of f_i , it follows that for all $j \in \{1, \dots, d\}$, $f_i(p_j) \geq 0$ if j is odd and $f_i(p_j) \leq 0$ if j is even. Since no p_j is a common root of f_1, \dots, f_k , it follows that for $1 \leq j \leq d$ we have $f(p_j) > 0$ if j is odd and $f(p_j) < 0$ if j is even.

So for $1 \leq j < d$ there exists r_j with $p_{j+1} < r_j < p_j$ such that $f(r_j) = 0$. Since f is a real polynomial of degree d (and therefore has an even number of non-real roots), it follows that all the roots of f are real. This proves that 3.6(3) implies 3.6(4).

Since 3.6(4) clearly implies 3.6(1), this completes the proof of 3.6. \square

The equivalence of 3.6(1) and 3.6(4) is closely related to a result of Bartlett, Hollot and Lin [1]. They prove the same assertion (indeed, they weaken statement 3.6(1), just requiring the compatibility of all pairs of f_1, \dots, f_k that are adjacent in the 1-skeleton of the boundary of the convex hull of f_1, \dots, f_k); but they require that f_1, \dots, f_k all have the same degree.

If we remove the hypothesis that the leading coefficients of f_1, \dots, f_k are all positive, 2.2 is no longer true. For example, the polynomials $x^2 - 2x, x^2 + 2x, 1 - x^2$ are pairwise compatible, but their sum does not have all roots real.

It is therefore natural to ask for an analogue of 3.6 without the assumption that the leading coefficients of the polynomials are all positive. We have already seen that statements 3.6(1) and 3.6(4) are not equivalent, but it is possible that statements 3.6(3) and 3.6(4) are equivalent under an appropriate modification of the definition of a common interleaver. We have not been able to decide this.

What if we ask for *all* linear combinations of the polynomials to have all roots real, instead of just all non-negative linear combinations? Let us say that a set f_1, \dots, f_k of polynomials is *strongly compatible* if for all $c_1, \dots, c_k \in \mathfrak{R}$, all the roots of the polynomial $\sum_{i=1}^k c_i f_i(x)$ are real. When $k = 2$, there is an analogue of 3.6; a theorem of Obreschkoff [14] when there are no common roots, and of Dedieu [6] in general, asserts that f_1, f_2 are strongly compatible if and only if their degrees differ by at most one, and the root sequence of one of them interleaves that

of the other. But there is no similar analogue for general k , because it is easy to see (thanks to David Moulton for pointing this out to us) that three polynomials are never strongly compatible unless they are linearly dependent.

Proof (Sketch). If two non-zero polynomials are strongly compatible, then their degrees differ by at most one; given three linearly independent polynomials f_1, f_2, f_3 , where $\deg(f_1) \geq \deg(f_2), \deg(f_3)$, there is a non-zero linear combination of them of degree at most $\deg(f_1) - 2$; so this linear combination is not strongly compatible with f_1 . \square

Acknowledgments

We thank Anders Björner, who told us about this conjecture at a graph theory meeting in Oberwolfach in January 2004. We also thank Alan Sokal and the referees, for their helpful comments and a number of useful references.

References

- [1] A.C. Bartlett, C.V. Hollot, Huang Lin, Root locations of an entire polytope of polynomials: It suffices to check the edges, *Math. Control Signals Systems* 1 (1988) 61–71.
- [2] J.I. Brown, R. Nowakowski, Bounding the roots of independence polynomials, *Ars Combin.* 58 (2001) 113–120.
- [3] J.I. Brown, R. Nowakowski, Average independence polynomials, *J. Combin. Theory Ser. B* 93 (2005) 313–318.
- [4] J.I. Brown, K. Dilcher, R.J. Nowakowski, Roots of independence polynomials of well-covered graphs, *J. Algebraic Combin.* 11 (2000) 197–210.
- [5] J.I. Brown, C.A. Hickman, R.J. Nowakowski, On the location of roots of independence polynomials, *J. Algebraic Combin.* 19 (2004) 273–282.
- [6] J.-P. Dedieu, Obreschkoff’s theorem revisited: What convex sets are contained in the set of hyperbolic polynomials?, *J. Pure Appl. Algebra* 81 (1992) 269–278.
- [7] J.-P. Dedieu, R.J. Gregorac, Corrigendum: Obreschkoff’s theorem revisited: What convex sets are contained in the set of hyperbolic polynomials?, *J. Pure Appl. Algebra* 93 (1994) 111–112.
- [8] D.C. Fisher, A.E. Solow, Dependence polynomials, *Discrete Math.* 82 (1990) 251–258.
- [9] I. Gutman, An identity for the independence polynomial of trees, *Publ. Inst. Math. (Belgrade)* 50 (1991) 19–23.
- [10] I. Gutman, Some analytic properties of the independence and matching polynomials, *MATCH Commun. Math. Comput. Chem.* 28 (1992) 139–150.
- [11] Y.O. Hamidoune, On the numbers of independent k -sets in a clawfree graph, *J. Combin. Theory Ser. B* 50 (1990) 241–244.
- [12] O.J. Heilmann, E.H. Lieb, Theory of monomer–dimer systems, *Comm. Math. Phys.* 25 (1972) 190–232.
- [13] C. Hoede, X. Li, Clique polynomials and independent set polynomials of graphs, *Discrete Math.* 25 (1994) 219–228.
- [14] N. Obreschkoff, *Verteilung und Berechnung der Nullstellen reeller Polynome*, VEB Deutscher, Verlag der Wissenschaftler, Berlin, 1963.
- [15] R.P. Stanley, Graph colorings and related symmetric functions: Ideas and applications, *Discrete Math.* 193 (1998) 267–286.