



# Szigetek

K. Horváth Eszter

Szeged, 2010. április 27.



# Definition/1

Grid, neighbourhood relation



## Definition/2

We call a rectangle/triangle an *island*, if for the cell  $t$ , if we denote its height by  $a_t$ , then for each cell  $\hat{t}$  neighbouring with a cell of the rectangle/triangle  $T$ , the inequality  $a_{\hat{t}} < \min\{a_t : t \in T\}$  holds.



## Coding theory

S. Földes and N. M. Singhi: On instantaneous codes, J. of Combinatorics, Information and System Sci., 31 (2006), 317-326.

## Coding theory

S. Földes and N. M. Singhi: On instantaneous codes, J. of Combinatorics, Information and System Sci., 31 (2006), 317-326.

## Rectangular islands

G. Czédli: The number of rectangular islands by means of distributive lattices, European Journal of Combinatorics 30 (2009), 208-215.

The maximum number of rectangular islands in a  $m \times n$  rectangular board on square grid:

$$f(m, n) = \left\lfloor \frac{mn + m + n - 1}{2} \right\rfloor.$$



## Rectangular islands

G. Czédli: The number of rectangular islands by means of distributive lattices, European Journal of Combinatorics 30 (2009), 208-215.

The maximum number of rectangular islands in a  $m \times n$  rectangular board on square grid:

$$f(m, n) = \left\lfloor \frac{mn + m + n - 1}{2} \right\rfloor.$$




## Rectangular islands

G. Czédli: The number of rectangular islands by means of distributive lattices, European Journal of Combinatorics 30 (2009), 208-215.

The maximum number of rectangular islands in a  $m \times n$  rectangular board on square grid:

$$f(m, n) = \left\lfloor \frac{mn + m + n - 1}{2} \right\rfloor.$$

$m$



$n$

## Rectangular islands in higher dimensions

G. Pluhár: The number of brick islands by means of distributive lattices, *Acta Sci. Math.*, to appear.

## Rectangular islands in higher dimensions


G. Pluhár: The number of brick islands by means of distributive lattices, *Acta Sci. Math.*, to appear.

## Triangular islands

E. K. Horváth, Z. Németh and G. Pluhár: The number of triangular islands on a triangular grid, *Periodica Mathematica Hungarica*, 58 (2009), 25–34.

Available at <http://www.math.u-szeged.hu/~horvath>

For the maximum number of triangular islands in an equilateral triangle of side length  $n$ ,  $\frac{n^2+3n}{5} \leq f(n) \leq \frac{3n^2+9n+2}{14}$  holds.




## Triangular islands

E. K. Horváth, Z. Németh and G. Pluhár: The number of triangular islands on a triangular grid, *Periodica Mathematica Hungarica*, 58 (2009), 25–34.

Available at <http://www.math.u-szeged.hu/~horvath>

For the maximum number of triangular islands in an equilateral triangle of side length  $n$ ,  $\frac{n^2+3n}{5} \leq f(n) \leq \frac{3n^2+9n+2}{14}$  holds.




## Triangular islands

E. K. Horváth, Z. Németh and G. Pluhár: The number of triangular islands on a triangular grid, *Periodica Mathematica Hungarica*, 58 (2009), 25–34.

Available at <http://www.math.u-szeged.hu/~horvath>

For the maximum number of triangular islands in an equilateral triangle of side length  $n$ ,  $\frac{n^2+3n}{5} \leq f(n) \leq \frac{3n^2+9n+2}{14}$  holds.




# History/5

## Square islands (also in higher dimensions)

E. K. Horváth, G. Horváth, Z. Németh, Cs. Szabó: The number of square islands on a rectangular sea, Acta Sci. Math., to appear.  
Available at <http://www.math.u-szeged.hu/~horvath>


$$\frac{1}{3}(rs - 2r - 2s) \leq f(r, s) \leq \frac{1}{3}(rs - 1)$$



## Square islands (also in higher dimensions)

E. K. Horváth, G. Horváth, Z. Németh, Cs. Szabó: The number of square islands on a rectangular sea, *Acta Sci. Math.*, to appear.  
Available at <http://www.math.u-szeged.hu/~horvath>


$$\frac{1}{3}(rs - 2r - 2s) \leq f(r, s) \leq \frac{1}{3}(rs - 1)$$



## Square islands (also in higher dimensions)

E. K. Horváth, G. Horváth, Z. Németh, Cs. Szabó: The number of square islands on a rectangular sea, *Acta Sci. Math.*, to appear.  
Available at <http://www.math.u-szeged.hu/~horvath>


$$\frac{1}{3}(rs - 2r - 2s) \leq f(r, s) \leq \frac{1}{3}(rs - 1)$$



## Square islands (also in higher dimensions)

E. K. Horváth, G. Horváth, Z. Németh, Cs. Szabó: The number of square islands on a rectangular sea, *Acta Sci. Math.*, to appear.  
Available at <http://www.math.u-szeged.hu/~horvath>

$$\frac{1}{3}(rs - 2r - 2s) \leq f(r, s) \leq \frac{1}{3}(rs - 1)$$



## LATTICE THEORETICAL METHOD

G. Czédli, A. P. Huhn and E. T. Schmidt: Weakly independent subsets in lattices, *Algebra Universalis* 20 (1985), 194-196.

Any two weak bases of a finite distributive lattice have the same number of elements.

## LATTICE THEORETICAL METHOD

G. Czédli, A. P. Huhn and E. T. Schmidt: Weakly independent subsets in lattices, *Algebra Universalis* 20 (1985), 194-196.


Any two weak bases of a finite distributive lattice have the same number of elements.

## LATTICE THEORETICAL METHOD


G. Czédli, A. P. Huhn and E. T. Schmidt: Weakly independent subsets in lattices, *Algebra Universalis* 20 (1985), 194-196.

Any two weak bases of a finite distributive lattice have the same number of elements.



## TREE-GRAPH METHOD



$T_0(I)$



## TREE-GRAPH METHOD



## TREE-GRAPH METHOD

### Lemma 2 (folklore)

- (i) Let  $T$  be a binary tree with  $\ell$  leaves. Then the number of vertices of  $T$  depends only on  $\ell$ , moreover  $|V| = 2\ell - 1$ .
- (ii) Let  $T$  be a rooted tree such that any non-leaf node has at least 2 sons. Let  $\ell$  be the number of leaves in  $T$ . Then  $|V| \leq 2\ell - 1$ .

We have  $4s + 2d \leq (n+1)(m+1)$ .

The number of leaves of  $T(\mathcal{I})$  is  $\ell = s + d$ . Hence by Lemma 2 the number of islands is

$$|V| - d \leq (2\ell - 1) - d = 2s + d - 1 \leq \frac{1}{2}(n+1)(m+1) - 1.$$

## TREE-GRAPH METHOD

### Lemma 2 (folklore)

- (i) Let  $T$  be a binary tree with  $\ell$  leaves. Then the number of vertices of  $T$  depends only on  $\ell$ , moreover  $|V| = 2\ell - 1$ .**
- (ii) Let  $T$  be a rooted tree such that any non-leaf node has at least 2 sons. Let  $\ell$  be the number of leaves in  $T$ . Then  $|V| \leq 2\ell - 1$ .

We have  $4s + 2d \leq (n + 1)(m + 1)$ .

The number of leaves of  $T(\mathcal{I})$  is  $\ell = s + d$ . Hence by Lemma 2 the number of islands is

$$|V| - d \leq (2\ell - 1) - d = 2s + d - 1 \leq \frac{1}{2}(n + 1)(m + 1) - 1.$$

## TREE-GRAPH METHOD

### Lemma 2 (folklore)

- (i) Let  $T$  be a binary tree with  $\ell$  leaves. Then the number of vertices of  $T$  depends only on  $\ell$ , moreover  $|V| = 2\ell - 1$ .
- (ii) Let  $T$  be a rooted tree such that any non-leaf node has at least 2 sons. Let  $\ell$  be the number of leaves in  $T$ . Then  $|V| \leq 2\ell - 1$ .

We have  $4s + 2d \leq (n + 1)(m + 1)$ .

The number of leaves of  $T(\mathcal{I})$  is  $\ell = s + d$ . Hence by Lemma 2 the number of islands is

$$|V| - d \leq (2\ell - 1) - d = 2s + d - 1 \leq \frac{1}{2}(n + 1)(m + 1) - 1.$$

## TREE-GRAPH METHOD

### Lemma 2 (folklore)

- (i) Let  $T$  be a binary tree with  $\ell$  leaves. Then the number of vertices of  $T$  depends only on  $\ell$ , moreover  $|V| = 2\ell - 1$ .
- (ii) Let  $T$  be a rooted tree such that any non-leaf node has at least 2 sons. Let  $\ell$  be the number of leaves in  $T$ . Then  $|V| \leq 2\ell - 1$ .

We have  $4s + 2d \leq (n + 1)(m + 1)$ .

The number of leaves of  $T(\mathcal{I})$  is  $\ell = s + d$ . Hence by Lemma 2 the number of islands is

$$|V| - d \leq (2\ell - 1) - d = 2s + d - 1 \leq \frac{1}{2}(n + 1)(m + 1) - 1.$$

.

## ELEMENTARY METHOD

We define

$$\mu(R) = \mu(u, v) := (u + 1)(v + 1).$$

Now

$$\begin{aligned} f(m, n) &= 1 + \sum_{R \in \max \mathcal{I}} f(R) = 1 + \sum_{R \in \max \mathcal{I}} \left( \left\lceil \frac{(u+1)(v+1)}{2} \right\rceil - 1 \right) \\ &= 1 + \sum_{R \in \max \mathcal{I}} \left( \left\lceil \frac{\mu(u, v)}{2} \right\rceil - 1 \right) \leq 1 - |\max \mathcal{I}| + \left\lceil \frac{\mu(C)}{2} \right\rceil. \end{aligned}$$

If  $|\max \mathcal{I}| \geq 2$ , then the proof is ready. Case  $|\max \mathcal{I}| = 1$  is an easy exercise.

## ELEMENTARY METHOD

We define

$$\mu(R) = \mu(u, v) := (u + 1)(v + 1).$$

Now

$$\begin{aligned} f(m, n) &= 1 + \sum_{R \in \max \mathcal{I}} f(R) = 1 + \sum_{R \in \max \mathcal{I}} \left( \left\lceil \frac{(u+1)(v+1)}{2} \right\rceil - 1 \right) \\ &= 1 + \sum_{R \in \max \mathcal{I}} \left( \left\lceil \frac{\mu(u, v)}{2} \right\rceil - 1 \right) \leq 1 - |\max \mathcal{I}| + \left\lceil \frac{\mu(C)}{2} \right\rceil. \end{aligned}$$

If  $|\max \mathcal{I}| \geq 2$ , then the proof is ready. Case  $|\max \mathcal{I}| = 1$  is an easy exercise.

## ELEMENTARY METHOD

We define

$$\mu(R) = \mu(u, v) := (u + 1)(v + 1).$$

Now

$$\begin{aligned} f(m, n) &= 1 + \sum_{R \in \max \mathcal{I}} f(R) = 1 + \sum_{R \in \max \mathcal{I}} \left( \left\lceil \frac{(u+1)(v+1)}{2} \right\rceil - 1 \right) \\ &= 1 + \sum_{R \in \max \mathcal{I}} \left( \left\lceil \frac{\mu(u, v)}{2} \right\rceil - 1 \right) \leq 1 - |\max \mathcal{I}| + \left\lceil \frac{\mu(C)}{2} \right\rceil. \end{aligned}$$

If  $|\max \mathcal{I}| \geq 2$ , then the proof is ready. Case  $|\max \mathcal{I}| = 1$  is an easy exercise.

## ELEMENTARY METHOD

We define

$$\mu(R) = \mu(u, v) := (u + 1)(v + 1).$$

Now

$$\begin{aligned} f(m, n) &= 1 + \sum_{R \in \max \mathcal{I}} f(R) = 1 + \sum_{R \in \max \mathcal{I}} \left( \left\lceil \frac{(u+1)(v+1)}{2} \right\rceil - 1 \right) \\ &= 1 + \sum_{R \in \max \mathcal{I}} \left( \left\lceil \frac{\mu(u, v)}{2} \right\rceil - 1 \right) \leq 1 - |\max \mathcal{I}| + \left\lceil \frac{\mu(C)}{2} \right\rceil. \end{aligned}$$

If  $|\max \mathcal{I}| \geq 2$ , then the proof is ready. Case  $|\max \mathcal{I}| = 1$  is an easy exercise.

## Some exact formulas

Peninsulas (semi islands) (J. Barát, P. Hajnal, E.K. Horváth):

$$p(m, n) = f(m, n) = \lfloor (mn + m + n - 1)/2 \rfloor.$$

Cylindric board, rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

If  $n \geq 2$ , then  $h_1(m, n) = \lfloor \frac{(m+1)n}{2} \rfloor$ .

Cylindric board, cylindric and rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

If  $n \geq 2$ , then  $h_2(m, n) = \lfloor \frac{(m+1)n}{2} \rfloor + \lfloor \frac{(m-1)}{2} \rfloor$ .

Torus board, rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

If  $m, n \geq 2$ , then  $t(m, n) = \lfloor \frac{mn}{2} \rfloor$ .

## Some exact formulas

Peninsulas (semi islands) (J. Barát, P. Hajnal, E.K. Horváth):

$$p(m, n) = f(m, n) = \lfloor (mn + m + n - 1)/2 \rfloor.$$

Cylindric board, rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

$$\text{If } n \geq 2, \text{ then } h_1(m, n) = \lfloor \frac{(m+1)n}{2} \rfloor.$$

Cylindric board, cylindric and rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

$$\text{If } n \geq 2, \text{ then } h_2(m, n) = \lfloor \frac{(m+1)n}{2} \rfloor + \lfloor \frac{(m-1)}{2} \rfloor.$$

Torus board, rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

$$\text{If } m, n \geq 2, \text{ then } t(m, n) = \lfloor \frac{mn}{2} \rfloor.$$

## Some exact formulas

Peninsulas (semi islands) (J. Barát, P. Hajnal, E.K. Horváth):

$$p(m, n) = f(m, n) = \lfloor (mn + m + n - 1)/2 \rfloor.$$

Cylindric board, rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

$$\text{If } n \geq 2, \text{ then } h_1(m, n) = \lfloor \frac{(m+1)n}{2} \rfloor.$$

Cylindric board, cylindric and rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

$$\text{If } n \geq 2, \text{ then } h_2(m, n) = \lfloor \frac{(m+1)n}{2} \rfloor + \lfloor \frac{(m-1)}{2} \rfloor.$$

Torus board, rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

$$\text{If } m, n \geq 2, \text{ then } t(m, n) = \lfloor \frac{mn}{2} \rfloor.$$

## Some exact formulas

Peninsulas (semi islands) (J. Barát, P. Hajnal, E.K. Horváth):

$$p(m, n) = f(m, n) = [(mn + m + n - 1)/2].$$

Cylindric board, rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

$$\text{If } n \geq 2, \text{ then } h_1(m, n) = \left\lfloor \frac{(m+1)n}{2} \right\rfloor.$$

Cylindric board, cylindric and rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

$$\text{If } n \geq 2, \text{ then } h_2(m, n) = \left\lfloor \frac{(m+1)n}{2} \right\rfloor + \left\lfloor \frac{(m-1)}{2} \right\rfloor.$$

Torus board, rectangular islands (J. Barát, P. Hajnal, E.K. Horváth):

$$\text{If } m, n \geq 2, \text{ then } t(m, n) = \left\lfloor \frac{mn}{2} \right\rfloor.$$

## Further results on rectangular islands

Zs. Lengvárszky: The minimum cardinality of maximal systems of rectangular islands, *European Journal of Combinatorics*, **30** (2009), 216-219.

## Further results on rectangular islands

Zs. Lengvárszky: The minimum cardinality of maximal systems of rectangular islands, *European Journal of Combinatorics*, **30** (2009), 216-219.

The board consists of all vertices of a hypercube, i.e. the elements of a Boolean algebra  $BA = \{0, 1\}^n$ .

We consider two cells neighbouring if their Hamming distance is 1.

We denote the maximum number of islands in  $BA = \{0, 1\}^n$  by  $b(n)$ .

Island formula for Boolean algebras (P. Hajnal, E.K. Horváth)

$$b(n) = 1 + 2^{n-1}.$$

The board consists of all vertices of a hypercube, i.e. the elements of a Boolean algebra  $BA = \{0, 1\}^n$ .

We consider two cells neighbouring if their Hamming distance is 1.

We denote the maximum number of islands in  $BA = \{0, 1\}^n$  by  $b(n)$ .

Island formula for Boolean algebras (P. Hajnal, E.K. Horváth)

$$b(n) = 1 + 2^{n-1}.$$

The board consists of all vertices of a hypercube, i.e. the elements of a Boolean algebra  $BA = \{0, 1\}^n$ .

We consider two cells neighbouring if their Hamming distance is 1.

We denote the maximum number of islands in  $BA = \{0, 1\}^n$  by  $b(n)$ .

Island formula for Boolean algebras (P. Hajnal, E.K. Horváth)  
 $b(n) = 1 + 2^{n-1}$ .

The board consists of all vertices of a hypercube, i.e. the elements of a Boolean algebra  $BA = \{0, 1\}^n$ .

We consider two cells neighbouring if their Hamming distance is 1.

We denote the maximum number of islands in  $BA = \{0, 1\}^n$  by  $b(n)$ .

Island formula for Boolean algebras (P. Hajnal, E.K. Horváth)  
 $b(n) = 1 + 2^{n-1}$ .

# Rectangular height functions/1

Joint work with Branimir Šešelja and Andreja Tepavčević

A *height function*  $h$  is a mapping from  $\{1, 2, \dots, m\} \times \{1, 2, \dots, n\}$  to  $\mathbb{N}$ ,  $h : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$ .

The co-domain of the height function is the lattice  $(\mathbb{N}, \leq)$ , where  $\mathbb{N}$  is the set of natural numbers under the usual ordering  $\leq$  and suprema and infima are max and min, respectively.

For every  $p \in \mathbb{N}$ , the *cut of the height function*, i.e. the  $p$ -cut of  $h$  is an ordinary relation  $h_p$  on  $\{1, 2, \dots, m\} \times \{1, 2, \dots, n\}$  defined by

$$(x, y) \in h_p \text{ if and only if } h(x, y) \geq p.$$

# Rectangular height functions/1

Joint work with Branimir Šešelja and Andreja Tepavčević

A *height function*  $h$  is a mapping from  $\{1, 2, \dots, m\} \times \{1, 2, \dots, n\}$  to  $\mathbb{N}$ ,  $h : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$ .

The co-domain of the height function is the lattice  $(\mathbb{N}, \leq)$ , where  $\mathbb{N}$  is the set of natural numbers under the usual ordering  $\leq$  and suprema and infima are max and min, respectively.

For every  $p \in \mathbb{N}$ , the *cut of the height function*, i.e. the  $p$ -cut of  $h$  is an ordinary relation  $h_p$  on  $\{1, 2, \dots, m\} \times \{1, 2, \dots, n\}$  defined by

$$(x, y) \in h_p \text{ if and only if } h(x, y) \geq p.$$

# Rectangular height functions/2

We say that two rectangles  $\{\alpha, \dots, \beta\} \times \{\gamma, \dots, \delta\}$  and  $\{\alpha_1, \dots, \beta_1\} \times \{\gamma_1, \dots, \delta_1\}$  are *distant* if they are disjoint and for every two cells, namely  $(a, b)$  from the first rectangle and  $(c, d)$  from the second, we have  $(a - c)^2 + (b - d)^2 \geq 4$ .

The height function  $h$  is called *rectangular* if for every  $p \in \mathbb{N}$ , every nonempty  $p$ -cut of  $h$  is a union of distant rectangles.

# Rectangular height functions/2

We say that two rectangles  $\{\alpha, \dots, \beta\} \times \{\gamma, \dots, \delta\}$  and  $\{\alpha_1, \dots, \beta_1\} \times \{\gamma_1, \dots, \delta_1\}$  are *distant* if they are disjoint and for every two cells, namely  $(a, b)$  from the first rectangle and  $(c, d)$  from the second, we have  $(a - c)^2 + (b - d)^2 \geq 4$ .

The height function  $h$  is called *rectangular* if for every  $p \in \mathbb{N}$ , every nonempty  $p$ -cut of  $h$  is a union of distant rectangles.

# Rectangular height functions/3

|   |   |   |   |   |
|---|---|---|---|---|
| 5 | 5 | 3 | 5 | 5 |
| 4 | 4 | 2 | 4 | 4 |
| 2 | 2 | 1 | 2 | 2 |

$$\Gamma_1 = \{1, 2, 3, 4, 5\} \times \{1, 2, 3\},$$

$$\Gamma_2 = \{1, 2, 3, 4, 5\} \times \{1, 2, 3\} \setminus \{(3, 1)\},$$

$$\Gamma_3 = \{(1, 2), (1, 3), (2, 2), (2, 3), (3, 3), (4, 2), (4, 3), (5, 2), (5, 3)\},$$

$$\Gamma_4 = \{(1, 2), (1, 3), (2, 2), (2, 3), (4, 2), (4, 3), (5, 2), (5, 3)\} \text{ and}$$

$$\Gamma_5 = \{(1, 3), (2, 3), (4, 3), (5, 3)\}$$

# Rectangular height functions/3

|   |   |   |   |   |
|---|---|---|---|---|
| 5 | 5 | 3 | 5 | 5 |
| 4 | 4 | 2 | 4 | 4 |
| 2 | 2 | 1 | 2 | 2 |

$$\Gamma_1 = \{1, 2, 3, 4, 5\} \times \{1, 2, 3\},$$

$$\Gamma_2 = \{1, 2, 3, 4, 5\} \times \{1, 2, 3\} \setminus \{(3, 1)\},$$

$$\Gamma_3 = \{(1, 2), (1, 3), (2, 2), (2, 3), (3, 3), (4, 2), (4, 3), (5, 2), (5, 3)\},$$

$$\Gamma_4 = \{(1, 2), (1, 3), (2, 2), (2, 3), (4, 2), (4, 3), (5, 2), (5, 3)\} \text{ and}$$

$$\Gamma_5 = \{(1, 3), (2, 3), (4, 3), (5, 3)\}$$

# Rectangular height functions/4

## CHARACTERIZATION THEOREM

### Theorem 1

A height function  $h_{\mathbb{N}} : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  is rectangular if and only if for all  $(\alpha, \gamma), (\beta, \delta) \in \{1, 2, \dots, m\} \times \{1, 2, \dots, n\}$  either

- these are not neighboring cells and there is a cell  $(\mu, \nu)$  between  $(\alpha, \gamma)$  and  $(\beta, \delta)$  such that  $h_{\mathbb{N}}(\mu, \nu) < \min\{h_{\mathbb{N}}(\alpha, \gamma), h_{\mathbb{N}}(\beta, \delta)\}$ , or
- for all  $(\mu, \nu) \in [\min\{\alpha, \beta\}, \max\{\alpha, \beta\}] \times [\min\{\gamma, \delta\}, \max\{\gamma, \delta\}]$ ,

$$h_{\mathbb{N}}(\mu, \nu) \geq \min\{h_{\mathbb{N}}(\alpha, \gamma), h_{\mathbb{N}}(\beta, \delta)\}.$$

# Rectangular height functions/4

## CHARACTERIZATION THEOREM

### Theorem 1


A height function  $h_{\mathbb{N}} : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  is rectangular if and only if for all  $(\alpha, \gamma), (\beta, \delta) \in \{1, 2, \dots, m\} \times \{1, 2, \dots, n\}$  either

- these are not neighboring cells and there is a cell  $(\mu, \nu)$  between  $(\alpha, \gamma)$  and  $(\beta, \delta)$  such that  $h_{\mathbb{N}}(\mu, \nu) < \min\{h_{\mathbb{N}}(\alpha, \gamma), h_{\mathbb{N}}(\beta, \delta)\}$ , or
- for all  $(\mu, \nu) \in [\min\{\alpha, \beta\}, \max\{\alpha, \beta\}] \times [\min\{\gamma, \delta\}, \max\{\gamma, \delta\}]$ ,

$$h_{\mathbb{N}}(\mu, \nu) \geq \min\{h_{\mathbb{N}}(\alpha, \gamma), h_{\mathbb{N}}(\beta, \delta)\}.$$

## Theorem 2

For every height function  $h : \{1, 2, \dots, n\} \times \{1, 2, \dots, m\} \rightarrow \mathbb{N}$ , there is a rectangular height function  $h^* : \{1, 2, \dots, n\} \times \{1, 2, \dots, m\} \rightarrow \mathbb{N}$ , such that  $\mathcal{I}_{rect}(h) = \mathcal{I}_{rect}(h^*)$ .



# Rectangular height functions/6

## CONSTRUCTING ALGORITHM

1. FOR  $i = t$  TO 0
2. FOR  $y = 1$  TO  $n$
3. FOR  $x = 1$  TO  $m$
4. IF  $h(x, y) = a_i$  THEN
5.  $j := i$
6. WHILE there is no island of  $h$  which is a subset of  $h_{a_j}$  that contains  $(x, y)$  DO  $j := j - 1$
7. ENDWHILE
8. Let  $h^*(x, y) := a_j$ .
9. ENDIF
10. NEXT  $x$
11. NEXT  $y$
12. NEXT  $i$
13. END.

# Rectangular height functions/7

## LATTICE-VALUED REPRESENTATION

### Theorem 3

Let  $h : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  be a rectangular height function. Then there is a lattice  $L$  and an  $L$ -valued mapping  $\Phi$ , such that the cuts of  $\Phi$  are precisely all islands of  $h$ .

# Rectangular height functions/8

Let  $h : \{1, 2, 3, 4, 5\} \times \{1, 2, 3, 4\} \rightarrow \mathbb{N}$  be a height function.

|   |   |   |   |   |   |
|---|---|---|---|---|---|
| 4 | 9 | 8 | 7 | 1 | 5 |
| 3 | 8 | 8 | 7 | 1 | 4 |
| 2 | 7 | 7 | 7 | 1 | 5 |
| 1 | 2 | 2 | 2 | 1 | 6 |
|   | 1 | 2 | 3 | 4 | 5 |

# Rectangular height functions/9

$h$  is a rectangular height function. Its islands are:

$$I_1 = \{(1, 4)\},$$

$$I_2 = \{(1, 3), (1, 4), (2, 3), (2, 4)\},$$

$$I_3 = \{(1, 2), (1, 3), (1, 4), (2, 2), (2, 3), (2, 4), (3, 2), (3, 3), (3, 4)\},$$

$$I_4 = \{(5, 1)\},$$

$$I_5 = \{(5, 1), (5, 2)\},$$

$$I_6 = \{(5, 4)\},$$

$$I_7 = \{(5, 1), (5, 2), (5, 3), (5, 4)\},$$

$$I_8 = \{(1, 2), (1, 3), (1, 4), (2, 2), (2, 3), (2, 4), (3, 2), (3, 3), (3, 4), (1, 1), (2, 1), (3, 1)\},$$

$$I_9 = \{1, 2, 3, 4, 5\} \times \{1, 2, 3, 4\}.$$

# Rectangular height functions/10

Its cut relations are:

$$h_{10} = \emptyset$$

$$h_9 = I_1 \text{ (one-element island)}$$

$$h_8 = I_2 \text{ (four-element square island)}$$

$$h_7 = I_3 \text{ (nine-element square island)}$$

$$h_6 = I_3 \cup I_4 \text{ (this cut is a disjoint union of two islands)}$$


$$h_5 = I_3 \cup I_5 \cup I_6 \text{ (union of three islands)}$$

$$h_4 = I_3 \cup I_7 \text{ (union of two islands)}$$

$$h_2 = I_7 \cup I_8 \text{ (union of two islands)}$$

$$h_1 = \{1, 2, 3, 4, 5\} \times \{1, 2, 3, 4\} = I_9 \text{ (the whole domain)}$$

# Rectangular height functions/11



$$L = (\mathcal{I}_0(\Gamma), \supseteq)$$

## Theorem 4

For every rectangular height function

$$h^* : \{1, 2, \dots, n\} \times \{1, 2, \dots, m\} \rightarrow \mathbb{N},$$

there is a rectangular height function

$$h^{**} : \{1, 2, \dots, n\} \times \{1, 2, \dots, m\} \rightarrow \mathbb{N},$$

such that  $\mathcal{I}_{rect}(h^*) = \mathcal{I}_{rect}(h^{**})$  and in  $h^{**}$  every island appears exactly in one cut.

If a rectangular height function  $h^{**}$  has the property that each island appears exactly in one cut, then we call it *standard rectangular height function*.

## Theorem 4

For every rectangular height function

$$h^* : \{1, 2, \dots, n\} \times \{1, 2, \dots, m\} \rightarrow \mathbb{N},$$

there is a rectangular height function

$$h^{**} : \{1, 2, \dots, n\} \times \{1, 2, \dots, m\} \rightarrow \mathbb{N},$$

such that  $\mathcal{I}_{rect}(h^*) = \mathcal{I}_{rect}(h^{**})$  and in  $h^{**}$  every island appears exactly in one cut.

If a rectangular height function  $h^{**}$  has the property that each island appears exactly in one cut, then we call it *standard rectangular height function*.


We denote by  $\Lambda_{\max}(m, n)$  the maximum number of different nonempty  $p$ -cuts of a standard rectangular height function on the rectangular table of size  $m \times n$ .

**Theorem 5**  $\Lambda_{\max}(m, n) = m + n - 1$ .

We denote by  $\Lambda_{\max}(m, n)$  the maximum number of different nonempty  $p$ -cuts of a standard rectangular height function on the rectangular table of size  $m \times n$ .

**Theorem 5**  $\Lambda_{\max}(m, n) = m + n - 1$ .

# Rectangular height functions/14



The maximum number of different nonempty  $p$ -cuts of a standard rectangular height function is equal to the minimum cardinality of maximal systems of islands.

## Lemma 1

If  $m \geq 3$  and  $n \geq 3$  and a height function

$h : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  has maximally many islands, then it has exactly two maximal islands.

## Lemma 2

If  $m \geq 3$  or  $n \geq 3$ , then for any odd number  $t = 2k + 1$  with  $1 \leq t \leq \max\{m - 2, n - 2\}$ , there is a standard rectangular height function  $h : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  having the maximum number of islands  $f(m, n)$ , such that one of the side-lengths of one of the maximal islands is equal to  $t$ .

(Remark: The statement is not true for even side-lengths, one can construct counterexample easily!)

## Lemma 1

If  $m \geq 3$  and  $n \geq 3$  and a height function

$h : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  has maximally many islands, then it has exactly two maximal islands.

## Lemma 2

If  $m \geq 3$  or  $n \geq 3$ , then for any odd number  $t = 2k + 1$  with  $1 \leq t \leq \max\{m - 2, n - 2\}$ , there is a standard rectangular height function  $h : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  having the maximum number of islands  $f(m, n)$ , such that one of the side-lengths of one of the maximal islands is equal to  $t$ .

(Remark: The statement is not true for even side-lengths, one can construct counterexample easily!)

We denote by  $\Lambda_h^{cz}(m, n)$  the number of different nonempty cuts of a standard rectangular height function  $h$  in the case  $h$  has maximally many islands, i.e., when the number of islands is

$$f(m, n) = \left\lfloor \frac{mn + m + n - 1}{2} \right\rfloor.$$

## Theorem 6

Let  $h : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  be a standard rectangular height function having maximally many islands  $f(m, n)$ . Then,  
 $\Lambda_h^{cz}(m, n) \geq \lceil \log_2(m+1) \rceil + \lceil \log_2(n+1) \rceil - 1.$

We denote by  $\Lambda_h^{cz}(m, n)$  the number of different nonempty cuts of a standard rectangular height function  $h$  in the case  $h$  has maximally many islands, i.e., when the number of islands is

$$f(m, n) = \left\lfloor \frac{mn + m + n - 1}{2} \right\rfloor.$$

## Theorem 6

Let  $h : \{1, 2, \dots, m\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  be a standard rectangular height function having maximally many islands  $f(m, n)$ . Then,

$$\Lambda_h^{cz}(m, n) \geq \lceil \log_2(m+1) \rceil + \lceil \log_2(n+1) \rceil - 1.$$

# Islands and formal concepts

Neither every island is a formal concept, nor every formal concept is an island.

Let  $(C, D)$  be a formal concept of the context  $\mathbb{K} := (A, B, I)$ . Then, there are linear orderings  $\leq_1$  and  $\leq_2$  on  $A$  and  $B$ , respectively, such that  $C \times D$  is an island of  $I \subseteq (A, \leq_1) \times (B, \leq_2)$  if and only if there is an element  $a \in A$  such that  $a \notin C$  and  $b \in B$  such that  $b \notin D$ , with  $(a, b) \notin I$ ,  $(a, y) \notin I$  for every  $y \in D$  and  $(x, b) \notin I$  for every  $x \in C$ .

If  $A_1 \times B_1$  is an island of a relation  $I \subseteq (A, \leq_1) \times (B, \leq_2)$ , then  $(A_1, B_1)$  is a concept if and only if there is no  $a \in A \setminus A_1$  such that  $(a, b) \in I$  for all  $b \in B_1$  and there is no  $b \in B \setminus B_1$  such that  $(a, b) \in I$  for all  $a \in A_1$ .

Let  $I$  be a relation  $I \subseteq (A, \leq_1) \times (B, \leq_2)$ . Then, every island in a relation  $I$  is a concept if and only if every  $x \in A$  belongs to not more than one island and every  $y \in B$  belongs to not more than one island.

# Islands and formal concepts

Neither every island is a formal concept, nor every formal concept is an island.

Let  $(C, D)$  be a formal concept of the context  $\mathbb{K} := (A, B, I)$ . Then, there are linear orderings  $\leq_1$  and  $\leq_2$  on  $A$  and  $B$ , respectively, such that  $C \times D$  is an island of  $I \subseteq (A, \leq_1) \times (B, \leq_2)$  if and only if there is an element  $a \in A$  such that  $a \notin C$  and  $b \in B$  such that  $b \notin D$ , with  $(a, b) \notin I$ ,  $(a, y) \notin I$  for every  $y \in D$  and  $(x, b) \notin I$  for every  $x \in C$ .

If  $A_1 \times B_1$  is an island of a relation  $I \subseteq (A, \leq_1) \times (B, \leq_2)$ , then  $(A_1, B_1)$  is a concept if and only if there is no  $a \in A \setminus A_1$  such that  $(a, b) \in I$  for all  $b \in B_1$  and there is no  $b \in B \setminus B_1$  such that  $(a, b) \in I$  for all  $a \in A_1$ .

Let  $I$  be a relation  $I \subseteq (A, \leq_1) \times (B, \leq_2)$ . Then, every island in a relation  $I$  is a concept if and only if every  $x \in A$  belongs to not more than one island and every  $y \in B$  belongs to not more than one island.

# Islands and formal concepts

Neither every island is a formal concept, nor every formal concept is an island.

Let  $(C, D)$  be a formal concept of the context  $\mathbb{K} := (A, B, I)$ . Then, there are linear orderings  $\leq_1$  and  $\leq_2$  on  $A$  and  $B$ , respectively, such that  $C \times D$  is an island of  $I \subseteq (A, \leq_1) \times (B, \leq_2)$  if and only if there is an element  $a \in A$  such that  $a \notin C$  and  $b \in B$  such that  $b \notin D$ , with  $(a, b) \notin I$ ,  $(a, y) \notin I$  for every  $y \in D$  and  $(x, b) \notin I$  for every  $x \in C$ .

If  $A_1 \times B_1$  is an island of a relation  $I \subseteq (A, \leq_1) \times (B, \leq_2)$ , then  $(A_1, B_1)$  is a concept if and only if there is no  $a \in A \setminus A_1$  such that  $(a, b) \in I$  for all  $b \in B_1$  and there is no  $b \in B \setminus B_1$  such that  $(a, b) \in I$  for all  $a \in A_1$ .

Let  $I$  be a relation  $I \subseteq (A, \leq_1) \times (B, \leq_2)$ . Then, every island in a relation  $I$  is a concept if and only if every  $x \in A$  belongs to not more than one island and every  $y \in B$  belongs to not more than one island.

# Islands and formal concepts

Neither every island is a formal concept, nor every formal concept is an island.

Let  $(C, D)$  be a formal concept of the context  $\mathbb{K} := (A, B, I)$ . Then, there are linear orderings  $\leq_1$  and  $\leq_2$  on  $A$  and  $B$ , respectively, such that  $C \times D$  is an island of  $I \subseteq (A, \leq_1) \times (B, \leq_2)$  if and only if there is an element  $a \in A$  such that  $a \notin C$  and  $b \in B$  such that  $b \notin D$ , with  $(a, b) \notin I$ ,  $(a, y) \notin I$  for every  $y \in D$  and  $(x, b) \notin I$  for every  $x \in C$ .

If  $A_1 \times B_1$  is an island of a relation  $I \subseteq (A, \leq_1) \times (B, \leq_2)$ , then  $(A_1, B_1)$  is a concept if and only if there is no  $a \in A \setminus A_1$  such that  $(a, b) \in I$  for all  $b \in B_1$  and there is no  $b \in B \setminus B_1$  such that  $(a, b) \in I$  for all  $a \in A_1$ .

Let  $I$  be a relation  $I \subseteq (A, \leq_1) \times (B, \leq_2)$ . Then, every island in a relation  $I$  is a concept if and only if every  $x \in A$  belongs to not more than one island and every  $y \in B$  belongs to not more than one island.

G. Czédli, A. P. Huhn and E. T. Schmidt: Weakly independent subsets in lattices, *Algebra Universalis* 20 (1985), 194-196.

G. Czédli, M. Hartmann and E.T.Schmidt: CD-independent subsets in distributive lattices, *Publicationes Mathematicae Debrecen*, 74/1-2 (2009), 127–134.

G. Czédli and E. T. Schmidt: CDW-independent subsets in distributive lattices, *Acta Sci. Math. (Szeged)*, 75 (2009), 49–53.

G. Czédli, A. P. Huhn and E. T. Schmidt: Weakly independent subsets in lattices, *Algebra Universalis* 20 (1985), 194-196.

G. Czédli, M. Hartmann and E.T.Schmidt: CD-independent subsets in distributive lattices, *Publicationes Mathematicae Debrecen*, 74/1-2 (2009), 127–134.

G. Czédli and E. T. Schmidt: CDW-independent subsets in distributive lattices, *Acta Sci. Math. (Szeged)*, 75 (2009), 49–53.

G. Czédli, A. P. Huhn and E. T. Schmidt: Weakly independent subsets in lattices, *Algebra Universalis* 20 (1985), 194-196.

G. Czédli, M. Hartmann and E.T.Schmidt: CD-independent subsets in distributive lattices, *Publicationes Mathematicae Debrecen*, 74/1-2 (2009), 127–134.

G. Czédli and E. T. Schmidt: CDW-independent subsets in distributive lattices, *Acta Sci. Math. (Szeged)*, 75 (2009), 49–53.

# CD-independence in posets

Join work with Sándor Radeleczki

Let  $\mathbb{P} = (P, \leq)$  be a partially ordered set and  $a, b \in P$ . The elements  $a$  and  $b$  are called *disjoint*, and we write  $a \perp b$ , if

$\inf\{a, b\} = 0$ , whenever  $\mathbb{P}$  has a least element  $0 \in P$ ,  
 $a$  and  $b$  have no common lowerbound, whenever  $\mathbb{P}$  is without  $0$ .

Notice, that  $a \perp b$  implies  $x \perp y$  for all  $x, y \in P$  with  $x \leq a$  and  $y \leq b$ . A nonempty set  $X \subseteq P$  is called *CD-independent*, if for any  $x, y \in X$  either  $x \leq y$  or  $y \leq x$  or  $x \perp y$  holds.

Maximal CD-independent sets (with respect to  $\subseteq$ ) are called *CD-bases* in  $\mathbb{P}$ .

If  $\mathbb{P}$  contains a least element  $0$  (a greatest element  $1$ ) and  $B$  is a CD-base, then obviously,  $0 \in B$  ( $1 \in B$ ).

A nonempty set  $D$  of nonzero elements of  $P$  is called a *disjoint system* in  $\mathbb{P}$ , if  $x \perp y$  holds for all  $x, y \in D$ ,  $x \neq y$ .

If  $\mathbb{P}$  is with 0-element, then  $\{0\}$  is considered to be a disjoint system, too. Clearly, any disjoint system  $D \subseteq P$  and any chain  $C \subseteq P$  is a CD-independent set, and observe also, that  $D$  is a disjoint system, if and only if it is a CD-independent antichain in  $\mathbb{P}$ .

Let  $\mathcal{D}(P)$  stand for the set of all disjoint systems of  $\mathbb{P}$ . Since the disjoint systems of  $\mathbb{P}$  are also antichains, restricting  $\leq$  to  $\mathcal{D}(P)$ , we obtain a poset  $(\mathcal{D}(P), \leq)$ . Clearly, if  $\mathbb{P}$  has a least element 0, then  $\{0\}$  itself is the least element of  $(\mathcal{D}(P), \leq)$ .

**Proposition 1.** *Any poset  $\mathbb{P} = (P, \leq)$  has at least one CD-base, and the set  $P$  is covered by the CD-bases of  $\mathbb{P}$ .*

We recall that any antichain  $A = \{a_i \mid i \in I\}$  of a poset  $\mathbb{P}$  determines a unique order-ideal  $I(A)$  of  $\mathbb{P}$ , namely

$$I(A) = \bigcup_{i \in I} (a_i] = \{x \in P \mid x \leq a_i, \text{ for some } i \in I\},$$

where  $(a]$  stands for the principal ideal of an element  $a \in P$ .

**Proposition 2.** *If  $B$  is a CD-base and  $D \subseteq B$  is a disjoint system in the poset  $(P, \leq)$ , then  $I(D) \cap B$  is also a CD-base in the subposet  $(I(D), \leq)$ .*

**Theorem 7.** *Let  $(P, \leq)$  be a finite poset and  $B$  a CD-base of it. Then the following assertions hold:*

(i) *There exists a maximal chain  $D_1 \succ \dots \succ D_n$  in  $\mathcal{D}(P)$ , such that*

$$B = \bigcup_{i=1}^n D_i \text{ and } n = |B|.$$

(ii) *For any maximal chain  $D_1 \prec \dots \prec D_m$  in  $\mathcal{D}(P)$  the set  $D = \bigcup_{i=1}^m D_i$  is a CD-base in  $(P, \leq)$  with  $|D| = m$ .*

$\mathbb{P}$  is called *graded*, if all its maximal chains have the same cardinality.

**Proposition 3.** *Let  $\mathbb{P} = (P, \leq)$  be a finite poset. Then the CD-bases of  $\mathbb{P}$  have the same number of elements if and only if the poset  $\mathcal{D}(P)$  is graded.*

**Corollary 2.** *Let  $(P, \leq)$  be a finite poset and  $(B, \leq)$  its subposet corresponding to a CD-base  $B \subseteq P$ . Then any maximal chain  $\mathcal{C} : D_1 \prec \dots \prec D_n$  in  $\mathcal{D}(B)$  is also a maximal chain in  $\mathcal{D}(P)$ .*

The complete disjoint systems of  $\mathbb{P}$  form a principal filter  $[A(P))$  in  $\mathcal{D}(P)$ . Their subposet  $([A(P)), \leq)$  will be denoted by  $\mathcal{DC}(P)$ .

**Proposition 4.** *Let  $\mathbb{P} = (P, \leq)$  be a finite poset with 0. Then the following conditions are equivalent.*

- (i) *The CD-bases of  $\mathbb{P}$  have the same number of elements,*
- (ii)  *$\mathcal{D}(P)$  is graded.*
- (iii)  *$\mathcal{DC}(P)$  is graded.*

Let  $\mathbb{P} = (P, \wedge)$  be a semilattice with 0. Now, for any  $a, b \in P$  the relation  $a \perp b$  means that  $a \wedge b = 0$ . Hence, a set  $\{a_i \mid i \in I\}$  of nonzero elements is a disjoint system if and only if  $a_i \wedge a_j = 0$ , for all  $i, j \in I$ ,  $i \neq j$ . A pair of elements  $a, b \in P$  with a least upperbound  $a \vee b$  in  $\mathbb{P}$  is called a *distributive pair*, if  $(c \wedge a) \vee (c \wedge b)$  there exists in  $\mathbb{P}$  for any  $c \in P$ , and  $c \wedge (a \vee b) = (c \wedge a) \vee (c \wedge b)$ .

**Theorem 8.** *If  $\mathbb{P} = (P, \wedge)$  is a semilattice with 0, then  $\mathcal{D}(P)$  is a semilattice with 0; if  $D_1 \cup D_2$  is a CD-independent set for some  $D_1, D_2 \in \mathcal{D}(P)$ , then  $D_1, D_2$  is a distributive pair in  $\mathcal{D}(P)$ . If  $\mathbb{P}$  is a complete lattice, then  $\mathcal{D}(P)$  is a complete lattice, too.*

We say that  $(P, \wedge)$  is *dp-distributive* (distributive with respect to disjoint pairs), if any  $a, b \in P$  with  $a \wedge b = 0$  is a distributive pair.

**Corollary 2.** *If  $\mathbb{P} = (P, \wedge)$  is a semilattice with 0, then  $\mathcal{D}(P)$  is a dp-distributive semilattice.*

Let  $(P, \leq)$  be a poset and  $A \subseteq P$ .  $(A, \leq)$  is called a *sublattice* of  $(P, \leq)$ , if  $(A, \leq)$  is a lattice such that for any  $a, b \in A$  the infimum and the supremum of  $\{a, b\}$  is the same in the subposet  $(A, \leq)$  and in  $(P, \leq)$ . If the relation  $x \prec y$  in  $(A, \leq)$ , for some  $x, y \in A$  implies  $x \prec y$  in the poset  $(P, \leq)$ , then we say that  $(A, \leq)$  is an *cover-preserving subposet* of  $(P, \leq)$ .

**Proposition 5.** *Let  $\mathbb{P} = (P, \leq)$  be a poset with 0 and  $B$  a CD-base of it. Then  $(\mathcal{D}(B), \leq)$  is a distributive cover-preserving sublattice of the poset  $(\mathcal{D}(P), \leq)$ . If  $\mathbb{P}$  is a  $\wedge$ -semilattice, then for any  $D \in \mathcal{D}(P)$  and  $D_1, D_2 \in \mathcal{D}(B)$  we have  $(D_1 \vee D_2) \wedge D = (D_1 \wedge D) \vee (D_2 \wedge D)$ .*

A lattice  $L$  with  $0$  is called *pseudocomplemented* if for each  $x \in L$  there exists an element  $x^* \in L$  such that for any  $y \in L$ ,  $y \wedge x = 0 \Leftrightarrow y \leq x^*$ . It is known that an algebraic lattice  $L$  is pseudocomplemented if and only if it is *0-distributive*, that is, for any  $a, b, x \in L$ ,  $x \wedge a = 0$  and  $x \wedge b = 0$  imply  $x \wedge (a \vee b) = 0$ . We say that  $L$  is *weakly 0-distributive* if this implication holds under the assumption  $a \wedge b = 0$ .

We say that two elements  $a, b \in L$  form a *modular pair* in the lattice  $L$ , and we write  $(a, b)M$ , if for any  $x \in L$ ,  $x \leq b$  implies  $x \vee (a \wedge b) = (x \vee a) \wedge b$ .  $a, b$  is called a *dual-modular pair* if for any  $x \in L$ ,  $x \geq b$  implies  $x \wedge (a \vee b) = (x \wedge a) \vee b$ . This is denoted by  $(a, b)M^*$ . Clearly, if  $a, b$  is a distributive pair, then  $(a, b)M^*$  is satisfied. We say that a lattice  $L$  satisfies condition  $M_0^*$ , if for all  $a, b \in L$  with  $a \wedge b = 0$ ,  $(a, b)M^*$  holds.

**Theorem 9.** *Let  $L$  be a finite weakly 0-distributive lattice that satisfies condition  $M_0^*$ . Then the CD-bases of  $L$  have the same number of elements if and only if  $L$  is graded.*

If  $a \wedge b \neq 0$ , then  $(x \leq a \vee b \text{ and } x \wedge a = 0) \implies x \leq b$ , for any  $a, b, x \in L$ . ( $\mathcal{I}$ )

**Theorem 10.** *Let  $L$  be a finite, weakly modular lattice satisfying condition ( $\mathcal{I}$ ). Then the CD-bases of  $L$  have the same number of elements.*