

COMPLETE POLYNOMIAL VECTOR FIELDS IN BALL

N.M. BEN YOUSIF

ABSTRACT. We describe the complete polynomial vector fields in the unit ball of a Euclidean space.

1. INTRODUCTION

By definition, given any subset K in \mathbb{R}^N the set real n -tuples and a mapping $v: \mathbb{R}^N \rightarrow \mathbb{R}^N$, we say that v is a complete vector field in K if for every point $k_0 \in K$ there exists a curve $x: \mathbb{R} \rightarrow K$ such that $x(0) = k_0$ and $\frac{dx(t)}{dt} = v(x(t))$ for all $t \in \mathbb{R}$. The mapping $v: \mathbb{R}^N \rightarrow \mathbb{R}^N$ is said to be a polynomial vector field if $v(x) = (P_1(x), \dots, P_N(x))$, $x \in \mathbb{R}^N$ for some polynomials $P_1, \dots, P_N: \mathbb{R}^N \rightarrow \mathbb{R}$ of N variables (that is each p_i is a finite linear combination of functions of the form $x_1^{m_1} \dots x_N^{m_N}$ with non-negative integers m_j where $x_j: (\xi_1, \dots, \xi_j) \mapsto \xi_j$ denotes the j -th canonical coordinate function \mathbb{R}^N). By writing $\langle (\xi_1, \dots, \xi_N), (\eta_1, \dots, \eta_N) \rangle = \sum_{i=1}^N \xi_i \eta_i$ for the inner product in \mathbb{R}^N , it is easy to see [1] that a polynomial (or even smooth) vector field is complete in the ball $B := (\langle x, x \rangle < 1)$ iff it is complete in the sphere $S := (\langle x, x \rangle = 1)$. Furthermore, v is complete in S iff it is orthogonal to the radius vector on S , i.e. if $\langle v(x), x \rangle = 0$ for $x \in S$.

In 2001 L.L. Stachó [2] described the complete real polynomial vector fields on the unit disc \mathbb{D} of the space \mathbb{C} of complex numbers. He has shown that a real polynomial vector field $P: \mathbb{C} \rightarrow \mathbb{C}$ is complete in \mathbb{D} iff P is a finite real linear combination of the functions iz , $\gamma \bar{z}^m - \bar{\gamma} z^{m+2}$ ($\gamma \in \mathbb{C}$, $m = 0, 1, \dots$) and $(1 - |z|^2)Q$ where Q is any real polynomial: $\mathbb{C} \rightarrow \mathbb{C}$. In this paper we describe the complete polynomial vector fields of the Euclidean unit ball B (or equivalently the unit Euclidean sphere S) of \mathbb{R}^N . We show that $P: \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a complete polynomial vector field in B if and only if $P(x) = R(x) - \langle R(x), x \rangle x + (1 - \langle x, x \rangle)Q(x)$ for some polynomial vector fields $Q, R: \mathbb{R}^N \rightarrow \mathbb{R}^N$.

Our result not only generalizes the result of [2] on \mathbb{D} , but it even simplifies it by showing that the complete polynomial vector fields on the unit disc of \mathbb{C} have the form $(ip(z)z + q(z)(1 - |z|^2))$ where $p, q: \mathbb{C} \rightarrow \mathbb{R}$ are any real polynomials. In our previous work [4] we represented complete vector fields on a simplex as polynomial combinations of finitely many basis complete vector fields. This idea motivated the formulation of our main result.

2. MAIN RESULT

Lemma 2.1. *Let $f: \mathbb{R}^N \rightarrow \mathbb{R}$ be a polynomial such that $f(x) = 0$ for $x \in S$ where $S = (\langle x, x \rangle = 1)$. Then there exists a polynomial $Q: \mathbb{R}^N \rightarrow \mathbb{R}$, such that $f(x) = (1 - \langle x, x \rangle)Q(x)$.*

2000 *Mathematics Subject Classification.* 46L70. Secondary: 17C65, 46G20.
Key words and phrases. Polynomial vector field, complete vector field.

Proof. Let $g: B \rightarrow \mathbb{R}$ be the function on the unit ball $B := (\langle x, x \rangle < 1)$, defined by $g(x) = \frac{f(x)}{(1-\langle x, x \rangle)}$. The function g is analytic, since it is the quotient of two polynomials. Thus $g(x) = \sum_{k=0}^{\infty} g_k(x)$ where g_k are k -homogeneous polynomials on \mathbb{R}^N . We have $f(\pm e) = 0$ if $\langle e, e \rangle = 1$, where e is the unit vector. So given $e \in \mathbb{R}^N$ with $\langle e, e \rangle = 1$, there exists a polynomial $P_e: \mathbb{R} \rightarrow \mathbb{R}$ of degree $\leq \deg f - 2$ such that $(1-t^2)P_e(t) = f(te)$. It follows that, for every fixed unit vector $e \in \mathbb{R}^N$ $g(te) = \frac{f(te)}{(1-t^2)} = P_e(t) = \sum_{k=0}^{\deg f - 2} \alpha_k(e)t^k$, with suitable constants $\alpha_0(e), \dots, \alpha_{\deg f - 2}(e) \in \mathbb{R}$. Hence we deduce that $g_k(te) = 0$ for $k > \deg f - 2$ and for all $t \in \mathbb{R}$ and unit vectors e . Then $g = \sum_{k=0}^{\deg f - 2} g_k$ is a polynomial. \square

Remark 2.2. In classical algebraic geometry [3] an analogous results is known for irreducible sets in \mathbb{K}^N where \mathbb{K} is an algebraically closed field. However we do not know any reference for the simple case of the Lemma.

Theorem 2.3. *Let $P: \mathbb{R}^N \rightarrow \mathbb{R}^N$ be a polynomial. Then P is a complete polynomial vector field in the sphere $S := (\langle x, x \rangle = 1)$ if and only if*

$$[P(x) = R(x) - \langle R(x), x \rangle x + (1 - \langle x, x \rangle)Q(x)]$$

for some polynomials; $R, Q: \mathbb{R}^N \rightarrow \mathbb{R}^N$.

Proof. Suppose $P(x) = R(x) - \langle R(x), x \rangle x + (1 - \langle x, x \rangle)Q(x)$ where $R, Q: \mathbb{R}^N \rightarrow \mathbb{R}^N$ are polynomials.

Then $\langle R(x) - \langle R(x), x \rangle x, x \rangle = \langle (x), x \rangle - \langle R(x), x \rangle \langle x, x \rangle = 0$ on S . Since $\langle x, x \rangle = 1$ for $x \in S$ and also $(1 - \langle x, x \rangle)Q(x) = 0$ on S . Therefore $\langle P(x), x \rangle = 0$ for $x \in S$, that is P is tangent to S . Conversely, suppose the polynomial vector field $P: \mathbb{R}^N \rightarrow \mathbb{R}^N$ is complete in S . Let $\tilde{P}(x) = P(x) - \langle P(x), x \rangle x$. Since P is tangent to S , we have $P(x) \perp x$ (i.e. $\langle P(x), x \rangle = 0$). This implies that $P(x) = \tilde{P}(x)$ or $P(x) - \tilde{P}(x) = 0$ on S . By the Lemma $P(x) - \tilde{P}(x) = (1 - \langle x, x \rangle)Q(x)$ for some $Q \in Plo.(\mathbb{R}^N, \mathbb{R}^N)$. Therefore $P(x) = \tilde{P}(x) + (1 - \langle x, x \rangle)Q(x) = P(x) - \langle P(x), x \rangle x + (1 - \langle x, x \rangle)Q(x)$. \square

REFERENCES

- [1] L. L. Stachó. On nonlinear projections of vector fields. In *NLA98: Convex analysis and chaos (Sakado, 1998)*, volume 1 of *Josai Math. Monogr.*, pages 47–54. Josai Univ., Sakado, 1999.
- [2] L. L. Stachó. A counterexample concerning contractive projections of real JB^* -triples. *Publ. Math. Debrecen*, 58(1-2):223–230, 2001.
- [3] Robert J. Walker. *Algebraic Curves*. Princeton Mathematical Series, vol. 13. Princeton University Press, Princeton, N. J., 1950.
- [4] N. M. Yousif. Complete polynomial vector fields on simplex set. *Electronic Journal of Qualitative Theory of Differential Equations (Szeged)*. Accepted for publication.

Received February 06, 2004.

BOLYAI INSTITUTE,
UNIVERSITY OF SZEGED,
ARADI VÉRTANÚK TERE 1.
H-6720, SZEGED, HUNGARY