

# MATRICES IN MODULAR LATTICES

BENEDEK SKUBLICS

Let  $(a_1, \dots, a_m, c_{12}, \dots, c_{1m})$  be a spanning von Neumann  $m$ -frame of a modular lattice  $L$ , and let  $(u_1, \dots, u_n, v_{12}, \dots, v_{1n})$  be a spanning von Neumann  $n$ -frame of the interval  $[0, a_1]$ . Assume that either  $m \geq 4$ , or  $L$  is Arguesian and  $m \geq 3$ . Let  $R^*$  denote the coordinate ring of  $(a_1, \dots, a_m, c_{12}, \dots, c_{1m})$ . If  $n \geq 2$ , then there is a ring  $S^*$  such that  $R^*$  is isomorphic to the ring of all  $n \times n$  matrices over  $S^*$ . If  $n \geq 4$  or  $L$  is Arguesian and  $n \geq 3$ , then we can choose  $S^*$  as the coordinate ring of  $(u_1, \dots, u_n, v_{12}, \dots, v_{1n})$ .

The proof uses product frames which were defined by Czedli [1]. The talk is based on [2].

## REFERENCES

- [1] G. Czédli: The product of von Neumann  $n$ -frames, its characteristic, and modular fractal lattices, *Algebra Universalis* 60 (2009), 217-230.
- [2] G. Czédli and B. Skublics: The ring of an outer von Neumann frame in modular lattices, *Algebra Universalis*, to appear.

*URL*: <http://www.math.u-szeged.hu/~bskublics/>

CSM – THE FIRST CONFERENCE OF PHD STUDENTS IN MATHEMATICS, JUNE 29 – JULY 2, 2010