

**The conference is supported by the
National Laboratory for Health Security project
RRF-2.3.1-21-2022-00006**

Optimal subsets in the stability regions of multistep methods

LAJOS LÓCZI

ELTE Faculty of Informatics, Hungary

LLoczi@inf.elte.hu

We study the linear stability regions of linear multistep or multiderivative multistep methods for initial value problems by using techniques that are straightforward to implement in modern computer algebra systems. In many applications, one is interested in (i) checking whether a given subset of the complex plane (e.g., a sector, disk, or parabola) is included in the stability region of the numerical method, (ii) finding the largest subset of a certain shape contained in the stability region of a given method, or (iii) finding the numerical method in a parametric family of multistep methods whose stability region contains the largest subset of a given shape. First, we describe a simple procedure to exactly calculate the stability angle α in the definition of $A(\alpha)$ -stability. As an illustration, we exactly compute the stability angles for the k -step BDF methods and for the k -step second-derivative multistep methods of Enright. Next, we determine the exact value of the stability radius in the BDF family for each k , that is, the radius of the largest disk in the left half of the complex plane, symmetric with respect to the real axis, touching the imaginary axis and lying in the stability region of the corresponding method. Finally, we demonstrate how some Schur–Cohn-type theorems of recursive nature and not relying on the root locus curve method can be used to exactly solve some optimization problems within infinite parametric families of multistep methods. As an example, we identify the unique method in a two-parameter family of implicit-explicit (IMEX) methods having the largest stability angle, then we find the unique method whose stability region contains the largest parabola.