

# Existence of solutions of nonlinear fractional differential equations at resonance \*

Wenjuan Rui †

*Department of Mathematics, China University of Mining and Technology,  
Xuzhou 221008, PR China*

**Abstract.** In this paper we study the existence of solutions of nonlinear fractional differential equations at resonance. By using the coincidence degree theory due to Mawhin, the existence of solutions is obtained.

*Keywords:* Fractional differential equations; Boundary value problems; Resonance; Coincidence degree theory

*MR Subject Classification:* 34A08, 34B15.

## 1 Introduction

Fractional calculus is a generalization of ordinary differentiation and integration on an arbitrary order that can be noninteger. This subject, as old as the problem of ordinary differential calculus, can go back to the times when Leibniz and Newton invented differential calculus. As is known to all, the problem for fractional derivative was originally raised by Leibniz in a letter, dated September 30, 1695.

In recent years, the fractional differential equations have received more and more attention. The fractional derivative has been occurring in many physical applications such as a non-Markovian diffusion process with memory [1], charge transport in amorphous semiconductors [2], propagations of mechanical waves in viscoelastic media [3], etc. Phenomena in electromagnetics, acoustics, viscoelasticity, electrochemistry and material science are also described by differential equations of fractional order (see [4-9]).

---

\*This research was supported by the Fundamental Research Funds for the Central Universities (2010LK SX01) and the Science Foundation of China University of Mining and Technology (2008A037).

†Corresponding author. *E-mail address:* wenjuanrui@126.com (W. Rui).

Recently boundary value problems (BVPs for short) for fractional differential equations at nonresonance have been studied in many papers (see [10-16]). Moreover, Kosmatov studied the BVPs for fractional differential equations at resonance (see [17]). Motivated by the work above, in this paper, we consider the following BVP of fractional equation at resonance

$$\begin{cases} D_{0+}^{\alpha}x(t) = f(t, x(t), x'(t), x''(t)), & t \in [0, 1], \\ x(0) = x'(0) = 0, & x''(0) = x''(1), \end{cases} \quad (1.1)$$

where  $D_{0+}^{\alpha}$  denotes the Caputo fractional differential operator of order  $\alpha$ ,  $2 < \alpha \leq 3$ .  $f : [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}$  is continuous.

The rest of this paper is organized as follows. Section 2 contains some necessary notations, definitions and lemmas. In Section 3, we establish a theorem on existence of solutions for BVP (1.1) under nonlinear growth restriction of  $f$ , basing on the coincidence degree theory due to Mawhin (see [18]). Finally, in Section 4, an example is given to illustrate the main result.

## 2 Preliminaries

In this section, we will introduce notations, definitions and preliminary facts which are used throughout this paper.

Let  $X$  and  $Y$  be real Banach spaces and let  $L : \text{dom}L \subset X \rightarrow Y$  be a Fredholm operator with index zero, and  $P : X \rightarrow X$ ,  $Q : Y \rightarrow Y$  be projectors such that

$$\begin{aligned} \text{Im}P &= \text{Ker}L, & \text{Ker}Q &= \text{Im}L, \\ X &= \text{Ker}L \oplus \text{Ker}P, & Y &= \text{Im}L \oplus \text{Im}Q. \end{aligned}$$

It follows that

$$L|_{\text{dom}L \cap \text{Ker}P} : \text{dom}L \cap \text{Ker}P \rightarrow \text{Im}L$$

is invertible. We denote the inverse by  $K_P$ .

If  $\Omega$  is an open bounded subset of  $X$ , and  $\text{dom}L \cap \overline{\Omega} \neq \emptyset$ , the map  $N : X \rightarrow Y$  will be called  $L$ -compact on  $\overline{\Omega}$  if  $QN(\overline{\Omega})$  is bounded and  $K_P(I - Q)N : \overline{\Omega} \rightarrow X$  is compact, where  $I$  is identity operator.

**Lemma 2.1.** ([18]) If  $\Omega$  is an open bounded set, let  $L : \text{dom}L \subset X \rightarrow Y$  be a Fredholm operator of index zero and  $N : X \rightarrow Y$   $L$ -compact on  $\overline{\Omega}$ . Assume that the following conditions are satisfied

- (1)  $Lx \neq \lambda Nx$  for every  $(x, \lambda) \in [(\text{dom}L \setminus \text{Ker}L)] \cap \partial\Omega \times (0, 1)$ ;
- (2)  $Nx \notin \text{Im}L$  for every  $x \in \text{Ker}L \cap \partial\Omega$ ;

(3)  $\text{deg}(QN|_{\text{Ker}L}, \text{Ker}L \cap \Omega, 0) \neq 0$ , where  $Q : Y \rightarrow Y$  is a projection such that  $\text{Im}L = \text{Ker}Q$ .

Then the equation  $Lx = Nx$  has at least one solution in  $\text{dom}L \cap \overline{\Omega}$ .

**Definition 2.1.** The Riemann-Liouville fractional integral operator of order  $\alpha > 0$  of a function  $x$  is given by

$$I_{0+}^{\alpha}x(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1}x(s)ds,$$

provided that the right side integral is pointwise defined on  $(0, +\infty)$ .

**Definition 2.2.** The Caputo fractional derivative of order  $\alpha > 0$  of a function  $x$  with  $x^{(n-1)}$  absolutely continuous on  $[0, 1]$  is given by

$$D_{0+}^{\alpha}x(t) = I_{0+}^{n-\alpha} \frac{d^n x(t)}{dt^n} = \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{n-\alpha-1}x^{(n)}(s)ds,$$

where  $n = -[-\alpha]$ .

**Lemma 2.2.** ([19]) Let  $\alpha > 0$  and  $n = -[-\alpha]$ . If  $x^{(n-1)} \in AC[0, 1]$ , then

$$I_{0+}^{\alpha}D_{0+}^{\alpha}x(t) = x(t) - \sum_{k=0}^{n-1} \frac{x^{(k)}(0)}{k!}t^k.$$

In this paper, we denote  $X = C^2[0, 1]$  with the norm  $\|x\|_X = \max\{\|x\|_{\infty}, \|x'\|_{\infty}, \|x''\|_{\infty}\}$  and  $Y = C[0, 1]$  with the norm  $\|y\|_Y = \|y\|_{\infty}$ , where  $\|x\|_{\infty} = \max_{t \in [0,1]} |x(t)|$ . Obviously, both  $X$  and  $Y$  are Banach spaces.

Define the operator  $L : \text{dom}L \subset X \rightarrow Y$  by

$$Lx = D_{0+}^{\alpha}x, \tag{2.1}$$

where

$$\text{dom}L = \{x \in X \mid D_{0+}^{\alpha}x(t) \in Y, x(0) = x'(0) = 0, x''(0) = x''(1)\}.$$

Let  $N : X \rightarrow Y$  be the Nemytski operator

$$Nx(t) = f(t, x(t), x'(t), x''(t)), \quad \forall t \in [0, 1].$$

Then BVP (1.1) is equivalent to the operator equation

$$Lx = Nx, \quad x \in \text{dom}L.$$

### 3 Main result

In this section, a theorem on existence of solutions for BVP (1.1) will be given.

**Theorem 3.1.** Let  $f : [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}$  be continuous. Assume that

(H<sub>1</sub>) there exist nonnegative functions  $p, q, r, s \in C[0, 1]$  with  $\Gamma(\alpha - 1) - 2(q_1 + r_1 + s_1) > 0$  such that

$$|f(t, u, v, w)| \leq p(t) + q(t)|u| + r(t)|v| + s(t)|w|, \quad \forall t \in [0, 1], (u, v, w) \in \mathbb{R}^3,$$

where  $p_1 = \|p\|_\infty$ ,  $q_1 = \|q\|_\infty$ ,  $r_1 = \|r\|_\infty$ ,  $s_1 = \|s\|_\infty$ .

(H<sub>2</sub>) there exists a constant  $B > 0$  such that for all  $w \in \mathbb{R}$  with  $|w| > B$  either

$$wf(t, u, v, w) > 0, \quad \forall t \in [0, 1], (u, v) \in \mathbb{R}^2$$

or

$$wf(t, u, v, w) < 0, \quad \forall t \in [0, 1], (u, v) \in \mathbb{R}^2.$$

Then BVP (1.1) has at least one solution in  $X$ .

Now, we begin with some lemmas below.

**Lemma 3.1.** Let  $L$  be defined by (2.1), then

$$\text{Ker}L = \{x \in X | x(t) = \frac{x''(0)}{2}t^2, \forall t \in [0, 1]\}, \quad (3.1)$$

$$\text{Im}L = \{y \in Y | \int_0^1 (1-s)^{\alpha-3}y(s)ds = 0\}. \quad (3.2)$$

**Proof.** By Lemma 2.2,  $D_{0+}^\alpha x(t) = 0$  has solution

$$x(t) = x(0) + x'(0)t + \frac{x''(0)}{2}t^2.$$

Combining with the boundary value condition of BVP (1.1), one has (3.1) hold.

For  $y \in \text{Im}L$ , there exists  $x \in \text{dom}L$  such that  $y = Lx \in Y$ . By Lemma 2.2, we have

$$x(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1}y(s)ds + x(0) + x'(0)t + \frac{x''(0)}{2}t^2.$$

Then, we have

$$x'(t) = \frac{1}{\Gamma(\alpha-1)} \int_0^t (t-s)^{\alpha-2}y(s)ds + x'(0) + x''(0)t$$

and

$$x''(t) = \frac{1}{\Gamma(\alpha - 2)} \int_0^t (t - s)^{\alpha-3} y(s) ds + x''(0).$$

By conditions of BVP (1.1), we can get that  $y$  satisfies

$$\int_0^1 (1 - s)^{\alpha-3} y(s) ds = 0.$$

Thus we get (3.2). On the other hand, suppose  $y \in Y$  and satisfies  $\int_0^1 (1 - s)^{\alpha-3} y(s) ds = 0$ . Let  $x(t) = I_{0+}^\alpha y(t)$ , then  $x \in \text{dom}L$  and  $D_{0+}^\alpha x(t) = y(t)$ . So that,  $y \in \text{Im}L$ . The proof is complete.

**Lemma 3.2.** Let  $L$  be defined by (2.1), then  $L$  is a Fredholm operator of index zero, and the linear continuous projector operators  $P : X \rightarrow X$  and  $Q : Y \rightarrow Y$  can be defined as

$$Px(t) = \frac{x''(0)}{2} t^2, \quad \forall t \in [0, 1],$$

$$Qy(t) = (\alpha - 2) \int_0^1 (1 - s)^{\alpha-3} y(s) ds, \quad \forall t \in [0, 1].$$

Furthermore, the operator  $K_P : \text{Im}L \rightarrow \text{dom}L \cap \text{Ker}P$  can be written by

$$K_P y(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - s)^{\alpha-1} y(s) ds, \quad \forall t \in [0, 1].$$

**Proof.** Obviously,  $\text{Im}P = \text{Ker}L$  and  $P^2x = Px$ . It follows from  $x = (x - Px) + Px$  that  $X = \text{Ker}P + \text{Ker}L$ . By simple calculation, we can get that  $\text{Ker}L \cap \text{Ker}P = \{0\}$ . Then we get

$$X = \text{Ker}L \oplus \text{Ker}P.$$

For  $y \in Y$ , we have

$$Q^2y = Q(Qy) = Qy \cdot (\alpha - 2) \int_0^1 (1 - s)^{\alpha-3} ds = Qy.$$

Let  $y = (y - Qy) + Qy$ , where  $y - Qy \in \text{Ker}Q = \text{Im}L$ ,  $Qy \in \text{Im}Q$ . It follows from  $\text{Ker}Q = \text{Im}L$  and  $Q^2y = Qy$  that  $\text{Im}Q \cap \text{Im}L = \{0\}$ . Then, we have

$$Y = \text{Im}L \oplus \text{Im}Q.$$

Thus

$$\dim \text{Ker}L = \dim \text{Im}Q = \text{codim} \text{Im}L = 1.$$

This means that  $L$  is a Fredholm operator of index zero.

From the definitions of  $P, K_P$ , it is easy to see that the generalized inverse of  $L$  is  $K_P$ . In fact, for  $y \in \text{Im}L$ , we have

$$LK_P y = D_{0+}^\alpha I_{0+}^\alpha y = y. \quad (3.3)$$

Moreover, for  $x \in \text{dom}L \cap \text{Ker}P$ , we get  $x(0) = x'(0) = x''(0) = 0$ . By Lemma 2.2, we obtain that

$$I_{0+}^\alpha Lx(t) = I_{0+}^\alpha D_{0+}^\alpha x(t) = x(t) + x(0) + x'(0)t + \frac{x''(0)}{2}t^2,$$

which together with  $x(0) = x'(0) = x''(0) = 0$  yields that

$$K_P Lx = x. \quad (3.4)$$

Combining (3.3) with (3.4), we know that  $K_P = (L|_{\text{dom}L \cap \text{Ker}P})^{-1}$ . The proof is complete.

**Lemma 3.3.** Assume  $\Omega \subset X$  is an open bounded subset such that  $\text{dom}L \cap \overline{\Omega} \neq \emptyset$ , then  $N$  is  $L$ -compact on  $\overline{\Omega}$ .

**Proof.** By the continuity of  $f$ , we can get that  $QN(\overline{\Omega})$  and  $K_P(I - Q)N(\overline{\Omega})$  are bounded. So, in view of the Arzelà-Ascoli theorem, we need only prove that  $K_P(I - Q)N(\overline{\Omega}) \subset X$  is equicontinuous.

From the continuity of  $f$ , there exists constant  $A > 0$  such that  $|(I - Q)Nx| \leq A$ ,  $\forall x \in \overline{\Omega}, t \in [0, 1]$ . Furthermore, denote  $K_{P,Q} = K_P(I - Q)N$  and for  $0 \leq t_1 < t_2 \leq 1$ ,  $x \in \overline{\Omega}$ , we have

$$\begin{aligned} & |(K_{P,Q}x)(t_2) - (K_{P,Q}x)(t_1)| \\ & \leq \frac{1}{\Gamma(\alpha)} \left| \int_0^{t_2} (t_2 - s)^{\alpha-1} (I - Q)Nx(s) ds - \int_0^{t_1} (t_1 - s)^{\alpha-1} (I - Q)Nx(s) ds \right| \\ & \leq \frac{A}{\Gamma(\alpha)} \left[ \int_0^{t_1} (t_2 - s)^{\alpha-1} - (t_1 - s)^{\alpha-1} ds + \int_{t_1}^{t_2} (t_2 - s)^{\alpha-1} ds \right] \\ & = \frac{A}{\Gamma(\alpha + 1)} (t_2^\alpha - t_1^\alpha), \end{aligned}$$

$$\begin{aligned} & |(K_{P,Q}x)'(t_2) - (K_{P,Q}x)'(t_1)| \\ & = \frac{\alpha - 1}{\Gamma(\alpha)} \left| \int_0^{t_2} (t_2 - s)^{\alpha-2} (I - Q)Nx(s) ds - \int_0^{t_1} (t_1 - s)^{\alpha-2} (I - Q)Nx(s) ds \right| \\ & \leq \frac{A}{\Gamma(\alpha - 1)} \left[ \int_0^{t_1} (t_2 - s)^{\alpha-2} - (t_1 - s)^{\alpha-2} ds + \int_{t_1}^{t_2} (t_2 - s)^{\alpha-2} ds \right] \\ & \leq \frac{A}{\Gamma(\alpha)} (t_2^{\alpha-1} - t_1^{\alpha-1}) \end{aligned}$$

and

$$\begin{aligned}
& |(K_{P,Q}x)''(t_2) - (K_{P,Q}x)''(t_1)| \\
&= \frac{(\alpha - 2)(\alpha - 1)}{\Gamma(\alpha)} \left| \int_0^{t_2} (t_2 - s)^{\alpha-3} (I - Q)Nx(s)ds - \int_0^{t_1} (t_1 - s)^{\alpha-3} (I - Q)Nx(s)ds \right| \\
&\leq \frac{A}{\Gamma(\alpha - 2)} \left[ \int_0^{t_1} (t_1 - s)^{\alpha-3} - (t_2 - s)^{\alpha-3} ds + \int_{t_1}^{t_2} (t_2 - s)^{\alpha-3} ds \right] \\
&\leq \frac{A}{\Gamma(\alpha - 1)} [t_1^{\alpha-2} - t_2^{\alpha-2} + 2(t_2 - t_1)^{\alpha-2}].
\end{aligned}$$

Since  $t^\alpha$ ,  $t^{\alpha-1}$  and  $t^{\alpha-2}$  are uniformly continuous on  $[0, 1]$ , we can get that  $K_{P,Q}(\overline{\Omega}) \subset C[0, 1]$ ,  $(K_{P,Q})'(\overline{\Omega}) \subset C[0, 1]$  and  $(K_{P,Q})''(\overline{\Omega}) \subset C[0, 1]$  are equicontinuous. Thus, we get that  $K_{P,Q} : \overline{\Omega} \rightarrow X$  is compact. The proof is completed.

**Lemma 3.4.** Suppose  $(H_1), (H_2)$  hold, then the set

$$\Omega_1 = \{x \in \text{dom}L \setminus \text{Ker}L \mid Lx = \lambda Nx, \lambda \in (0, 1)\}$$

is bounded.

**Proof.** Take  $x \in \Omega_1$ , then  $Nx \in \text{Im}L$ . By (3.2), we have

$$\int_0^1 (1 - s)^{\alpha-3} f(s, x(s), x'(s), x''(s))ds = 0.$$

Then, by the integral mean value theorem, there exists a constant  $\xi \in (0, 1)$  such that  $f(\xi, x(\xi), x'(\xi), x''(\xi)) = 0$ . Then from  $(H_2)$ , we have  $|x''(\xi)| \leq B$ .

From  $x \in \text{dom}L$ , we get  $x(0) = 0$  and  $x'(0) = 0$ . Therefore

$$|x'(t)| = \left| x'(0) + \int_0^t x''(s)ds \right| \leq \|x''\|_\infty.$$

and

$$|x(t)| = \left| x(0) + \int_0^t x'(s)ds \right| \leq \|x'\|_\infty.$$

That is

$$\|x\|_\infty \leq \|x'\|_\infty \leq \|x''\|_\infty. \tag{3.5}$$

By  $Lx = \lambda Nx$  and  $x \in \text{dom}L$ , we have

$$x(t) = \frac{\lambda}{\Gamma(\alpha)} \int_0^t (t - s)^{\alpha-1} f(s, x(s), x'(s), x''(s))ds + \frac{1}{2}x''(0)t^2.$$

Then we get

$$x'(t) = \frac{\lambda}{\Gamma(\alpha - 1)} \int_0^t (t - s)^{\alpha-2} f(s, x(s), x'(s), x''(s)) ds + x''(0)t$$

and

$$x''(t) = \frac{\lambda}{\Gamma(\alpha - 2)} \int_0^t (t - s)^{\alpha-3} f(s, x(s), x'(s), x''(s)) ds + x''(0).$$

Take  $t = \xi$ , we get

$$x''(\xi) = \frac{\lambda}{\Gamma(\alpha - 2)} \int_0^\xi (\xi - s)^{\alpha-3} f(s, x(s), x'(s), x''(s)) ds + x''(0).$$

Together with  $|x''(\xi)| \leq B$ ,  $(H_1)$  and (3.5), we have

$$\begin{aligned} |x''(0)| &\leq |x''(\xi)| + \frac{\lambda}{\Gamma(\alpha - 2)} \int_0^\xi (\xi - s)^{\alpha-3} |f(s, x(s), x'(s), x''(s))| ds \\ &\leq B + \frac{1}{\Gamma(\alpha - 2)} \int_0^\xi (\xi - s)^{\alpha-3} [p(s) + q(s)|x(s)| + r(s)|x'(s)| + s(s)|x''(s)|] ds \\ &\leq B + \frac{1}{\Gamma(\alpha - 2)} \int_0^\xi (\xi - s)^{\alpha-3} (p_1 + q_1\|x\|_\infty + r_1\|x'\|_\infty + s_1\|x''\|_\infty) ds \\ &\leq B + \frac{1}{\Gamma(\alpha - 2)} \int_0^\xi (\xi - s)^{\alpha-3} [p_1 + (q_1 + r_1 + s_1)\|x''\|_\infty] ds \\ &\leq B + \frac{1}{\Gamma(\alpha - 1)} [p_1 + (q_1 + r_1 + s_1)\|x''\|_\infty]. \end{aligned}$$

Then we have

$$\begin{aligned} \|x''\|_\infty &\leq \frac{1}{\Gamma(\alpha - 2)} \int_0^t (t - s)^{\alpha-3} |f(s, x(s), x'(s), x''(s))| ds + |x''(0)| \\ &\leq \frac{1}{\Gamma(\alpha - 2)} \int_0^t (t - s)^{\alpha-3} [p(s) + q(s)|x(s)| + r(s)|x'(s)| + s(s)|x''(s)|] ds + x''(0) \\ &\leq \frac{1}{\Gamma(\alpha - 2)} \int_0^t (t - s)^{\alpha-3} (p_1 + q_1\|x\|_\infty + r_1\|x'\|_\infty + s_1\|x''\|_\infty) ds + |x''(0)| \\ &\leq \frac{1}{\Gamma(\alpha - 2)} \int_0^t (t - s)^{\alpha-3} [p_1 + (q_1 + r_1 + s_1)\|x''\|_\infty] ds + |x''(0)| \\ &\leq \frac{1}{\Gamma(\alpha - 1)} [p_1 + (q_1 + r_1 + s_1)\|x''\|_\infty] + |x''(0)| \\ &\leq B + \frac{2}{\Gamma(\alpha - 1)} [p_1 + (q_1 + r_1 + s_1)\|x''\|_\infty]. \end{aligned}$$



Thus, from  $\Gamma(\alpha - 1) - 2(q_1 + r_1 + s_1) > 0$ , we obtain that

$$\|x''\|_\infty \leq \frac{2p_1 + \Gamma(\alpha - 1)B}{\Gamma(\alpha - 1) - 2(q_1 + r_1 + s_1)} := M_1.$$

Thus, together with (3.5), we get

$$\|x\|_\infty \leq \|x'\|_\infty \leq \|x''\|_\infty \leq M_1.$$

Therefore,

$$\|x\|_X \leq M_1.$$

So  $\Omega_1$  is bounded. The proof is complete.

**Lemma 3.5.** Suppose  $(H_2)$  holds, then the set

$$\Omega_2 = \{x | x \in \text{Ker}L, Nx \in \text{Im}L\}$$

is bounded.

**Proof.** For  $x \in \Omega_2$ , we have  $x(t) = \frac{x''(0)}{2}t^2$  and  $Nx \in \text{Im}L$ . Then we get

$$\int_0^1 (1-s)^{\alpha-3} f(s, \frac{x''(0)}{2}s^2, x''(0)s, x''(0)) ds = 0,$$

which together with  $(H_2)$  implies  $|x''(0)| \leq B$ . Thus, we have

$$\|x\|_X \leq B.$$

Hence,  $\Omega_2$  is bounded. The proof is complete.

**Lemma 3.6.** Suppose the first part of  $(H_2)$  holds, then the set

$$\Omega_3 = \{x | x \in \text{Ker}L, \lambda x + (1-\lambda)QNx = 0, \lambda \in [0, 1]\}$$

is bounded.

**Proof.** For  $x \in \Omega_3$ , we have  $x(t) = \frac{x''(0)}{2}t^2$  and

$$\lambda \frac{x''(0)}{2}t^2 + (1-\lambda)(\alpha-2) \int_0^1 (1-s)^{\alpha-3} f(s, \frac{x''(0)}{2}s^2, x''(0)s, x''(0)) ds = 0. \quad (3.6)$$

If  $\lambda = 0$ , then  $|x''(0)| \leq B$  because of the first part of  $(H_2)$ . If  $\lambda \in (0, 1]$ , we can also obtain  $|x''(0)| \leq B$ . Otherwise, if  $|x''(0)| > B$ , in view of the first part of  $(H_2)$ , one has

$$\lambda [x''(0)]^2 t^2 + (1-\lambda)\alpha \int_0^1 (1-s)^{\alpha-1} x''(0) f(s, \frac{x''(0)}{2}s^2, x''(0)s, x''(0)) ds > 0,$$

which contradicts to (3.6).

Therefore,  $\Omega_3$  is bounded. The proof is complete.

**Remark 3.1.** Suppose the second part of  $(H_2)$  hold, then the set

$$\Omega'_3 = \{x|x \in \text{Ker}L, -\lambda x + (1 - \lambda)QNx = 0, \lambda \in [0, 1]\}$$

is bounded.

**The proof of Theorem 3.1.** Set  $\Omega = \{x \in X |||x||_X < \max\{M_1, B\} + 1\}$ . It follows from Lemma 3.2 and 3.3 that  $L$  is a Fredholm operator of index zero and  $N$  is  $L$ -compact on  $\overline{\Omega}$ . By Lemma 3.4 and 3.5, we get that the following two conditions are satisfied

- (1)  $Lx \neq \lambda Nx$  for every  $(x, \lambda) \in [(\text{dom}L \setminus \text{Ker}L) \cap \partial\Omega] \times (0, 1)$ ;
- (2)  $Nx \notin \text{Im}L$  for every  $x \in \text{Ker}L \cap \partial\Omega$ .

Take

$$H(x, \lambda) = \pm\lambda x + (1 - \lambda)QNx.$$

According to Lemma 3.6 (or Remark 3.1), we know that  $H(x, \lambda) \neq 0$  for  $x \in \text{Ker}L \cap \partial\Omega$ . Therefore

$$\begin{aligned} \deg(QN|_{\text{Ker}L}, \Omega \cap \text{Ker}L, 0) &= \deg(H(\cdot, 0), \Omega \cap \text{Ker}L, 0) \\ &= \deg(H(\cdot, 1), \Omega \cap \text{Ker}L, 0) \\ &= \deg(\pm I, \Omega \cap \text{Ker}L, 0) \neq 0. \end{aligned}$$

So that, the condition (3) of Lemma 2.1 is satisfied. By Lemma 2.1, we can get that  $Lx = Nx$  has at least one solution in  $\text{dom}L \cap \overline{\Omega}$ . Therefore, BVP (1.1) has at least one solution. The proof is complete.

## 4 An example

**Example 4.1.** Consider the following BVP

$$\begin{cases} D_{0+}^{\frac{5}{2}}x(t) = \frac{1}{16}(x'' - 10) + \frac{t^2}{16}e^{-|x'|} + \frac{t^3}{16}\sin(x^2), & t \in [0, 1] \\ x(0) = x'(0) = 0, \quad x''(0) = x''(1). \end{cases} \quad (4.1)$$

Where

$$f(t, u, v, w) = \frac{1}{16}(w - 10) + \frac{t^2}{16}e^{-|v|} + \frac{t^3}{16}\sin(u^2).$$

Choose  $p(t) = \frac{10+2}{16}$ ,  $q(t) = 0$ ,  $r(t) = 0$ ,  $s(t) = \frac{1}{16}$ ,  $B = 10$ . We can get that  $q_1 = 0$ ,  $r_1 = 0$ ,  $s_1 = \frac{1}{16}$  and

$$\Gamma\left(\frac{5}{2} - 1\right) - 2(q_1 + r_1 + s_1) > 0.$$

Then, all conditions of Theorem 3.1 hold, so BVP (4.1) has at least one solution.

## References

- [1] R. Metzler, J. Klafter, Boundary value problems for fractional diffusion equations, *Phys. A* 278, 107-125 (2000).
- [2] H. Scher, E. Montroll, Anomalous transit-time dispersion in amorphous solids, *Phys. Rev. B* 12, 2455-2477 (1975).
- [3] F. Mainardi, Fractional diffusive waves in viscoelastic solids, in: J.L. Wegner, F.R. Norwood (Eds.), *Nonlinear Waves in Solids*, Fairfield, 93-97 (1995).
- [4] K. Diethelm, A.D. Freed, On the solution of nonlinear fractional order differential equations used in the modeling of viscoplasticity, in: F. Keil, W. Mackens, H. Voss, J. Werther (Eds.), *Scientific Computing in Chemical Engineering II-Computational Fluid Dynamics, Reaction Engineering and Molecular Properties*, pp. 217-224. Springer-Verlag, Heidelberg (1999).
- [5] L. Gaul, P. Klein, S. Kempfle, Damping description involving fractional operators, *Mech. Syst. Signal Process.* 5, 81-88 (1991).
- [6] W.G. Glockle, T.F. Nonnenmacher, A fractional calculus approach of self-similar protein dynamics, *Biophys. J.* 68, 46-53 (1995).
- [7] F. Mainardi, Fractional calculus: Some basic problems in continuum and statistical mechanics, in: A. Carpinteri, F. Mainardi (Eds.), *Fractals and Fractional Calculus in Continuum Mechanics*, pp. 291-348. Springer-Verlag, Wien (1997).
- [8] F. Metzler, W. Schick, H.G. Kilian, T.F. Nonnenmacher, Relaxation in filled polymers: A fractional calculus approach, *J. Chem. Phys.* 103, 7180-7186 (1995).
- [9] K.B. Oldham, J. Spanier, *The Fractional Calculus*, Academic Press, New York, London (1974).
- [10] R.P. Agarwal, D. O'Regan, S. Stanek, Positive solutions for Dirichlet problems of singular nonlinear fractional differential equations, *J. Math. Anal. Appl.* 371, 57-68 (2010).
- [11] Z. Bai, H. Lü, Positive solutions for boundary value problem of nonlinear fractional differential equation, *J. Math. Anal. Appl.* 311, 495-505 (2005).
- [12] E.R. Kaufmann, E. Mboumi, Positive solutions of a boundary value problem for a nonlinear fractional differential equation, *Electron. J. Qual. Theory Differ. Equ.* 3, 1-11 (2008).
- [13] H. Jafari, V.D. Gejji, Positive solutions of nonlinear fractional boundary value problems using Adomian decomposition method, *Appl. Math. Comput.* 180, 700-706 (2006).

- [14] M. Benchohra, S. Hamani, S.K. Ntouyas, Boundary value problems for differential equations with fractional order and nonlocal conditions, *Nonlinear Anal.* 71, 2391-2396 (2009).
- [15] S. Liang, J. Zhang, Positive solutions for boundary value problems of nonlinear fractional differential equation, *Nonlinear Anal.* 71, 5545-5550 (2009).
- [16] S. Zhang, Positive solutions for boundary-value problems of nonlinear fractional differential equations, *Electron. J. Differ. Equ.* 36, 1-12 (2006).
- [17] N. Kosmatov, A boundary value problem of fractional order at resonance, *Electron. J. Differ. Equ.* 135, 1-10 (2010).
- [18] J. Mawhin, Topological degree and boundary value problems for nonlinear differential equations in topological methods for ordinary differential equations, *Lecture Notes in Math.* 1537, 74-142 (1993) .
- [19] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, vol. 204 of North-Holland Mathematics Studies, Elsevier Science B.V., Amsterdam, The Netherlands (2006).

(Received July 24, 2011)