

# Eighty-three sublattices and planarity

**Gábor Czédli** (University of Szeged)

<http://www.math.u-szeged.hu/~czedli/>

—

Talk at the 57th SSAOS, Karolinka, September 1–7, 2019

August 28, 2019

Paper: dedicated to George Grätzer on his 83<sup>rd</sup> birthday.

Theorem (G. Czédli, 2019; arXiv:1901.00572, 100 pages, sharp! )  
<http://www.math.u-szeged.hu/~czedli/>

*Let  $L$  be a finite lattice, and let  $n := |L|$  denote the number of its elements. If  $L$  has at least  $83 \cdot 2^{n-8}$  sublattices, then it is planar.*

Planar lattice  $\iff \exists$  Hasse diagram without crossing edges.

Motivating earlier results (Note: 83 is "many")

- Claudia Mureşan (2017) many ideals (more than filters), <https://arxiv.org/abs/1710.10183>
- Czédli, Horváth, Ahmed: the five largest  $|\text{Sub}(L)|$  for  $|L| = n$ , arXiv:1812.11512 , Discuss. Math. Gen. Alg. Appl., to appear
- Czédli (AU 80:16, 2019):  $|\text{Con}(L)| > 2^{n-5} \Rightarrow L$  is planar.

Paper: dedicated to George Grätzer on his 83<sup>rd</sup> birthday.

Theorem (G. Czédli, 2019; arXiv:1901.00572, 100 pages, sharp! )  
<http://www.math.u-szeged.hu/~czedli/>

*Let  $L$  be a finite lattice, and let  $n := |L|$  denote the number of its elements. If  $L$  has at least  $83 \cdot 2^{n-8}$  sublattices, then it is planar.*

Planar lattice  $\iff \exists$  Hasse diagram without crossing edges.

Motivating earlier results (Note: 83 is "many")

- Claudia Mureşan (2017) many ideals (more than filters), <https://arxiv.org/abs/1710.10183>
- Czédli, Horváth, Ahmed: the five largest  $|\text{Sub}(L)|$  for  $|L| = n$ , arXiv:1812.11512, Discuss. Math. Gen. Alg. Appl., to appear
- Czédli (AU 80:16, 2019):  $|\text{Con}(L)| > 2^{n-5} \Rightarrow L$  is planar.

Paper: dedicated to George Grätzer on his 83<sup>rd</sup> birthday.

Theorem (G. Czédli, 2019; arXiv:1901.00572, 100 pages, sharp! )  
<http://www.math.u-szeged.hu/~czedli/>

*Let  $L$  be a finite lattice, and let  $n := |L|$  denote the number of its elements. If  $L$  has at least  $83 \cdot 2^{n-8}$  sublattices, then it is planar.*

Planar lattice  $\iff \exists$  Hasse diagram without crossing edges.

Motivating earlier results (Note: 83 is "many")

- Claudia Mureşan (2017) many ideals (more than filters), <https://arxiv.org/abs/1710.10183>
- Czédli, Horváth, Ahmed: the five largest  $|\text{Sub}(L)|$  for  $|L| = n$ , arXiv:1812.11512 , Discuss. Math. Gen. Alg. Appl., to appear
- Czédli (AU 80:16, 2019):  $|\text{Con}(L)| > 2^{n-5} \Rightarrow L$  is planar.

Paper: dedicated to George Grätzer on his 83<sup>rd</sup> birthday.

Theorem (G. Czédli, 2019; arXiv:1901.00572, 100 pages, sharp! )  
<http://www.math.u-szeged.hu/~czedli/>

*Let  $L$  be a finite lattice, and let  $n := |L|$  denote the number of its elements. If  $L$  has at least  $83 \cdot 2^{n-8}$  sublattices, then it is planar.*

Planar lattice  $\iff \exists$  Hasse diagram without crossing edges.

### Motivating earlier results (Note: 83 is "many")

- Claudia Mureşan (2017) many ideals (more than filters), <https://arxiv.org/abs/1710.10183>
- Czédli, Horváth, Ahmed: the five largest  $|\text{Sub}(L)|$  for  $|L| = n$ , arXiv:1812.11512, Discuss. Math. Gen. Alg. Appl., to appear
- Czédli (AU 80:16, 2019):  $|\text{Con}(L)| > 2^{n-5} \Rightarrow L$  is planar.

$$\sigma(\mathcal{A}) := |\text{Sub}(\mathcal{A})| \cdot 2^{8-n}, \quad \text{for a partial algebra.}$$

### Theorem (The main theorem reformulated)

*If  $L$  is a finite lattice such that  $\sigma(L) > 83$ , then  $L$  is planar.  
Furthermore, for every natural number  $n \geq 9$ , there exists an  $n$ -element lattice  $L$  such that  $\sigma(L) = 83$  and  $L$  is not planar.*

“ $\sigma$ -few” means “ $\sigma$  is at most 83”.

“ $\sigma$ -many” means “ $\sigma$  is strictly more than 83”.

### Theorem (Yet another formulation)

*Finite lattices with  $\sigma$ -many sublattices are planar.*

$$\sigma(\mathcal{A}) := |\text{Sub}(\mathcal{A})| \cdot 2^{8-n}, \quad \text{for a partial algebra.}$$

### Theorem (The main theorem reformulated)

*If  $L$  is a finite lattice such that  $\sigma(L) > 83$ , then  $L$  is planar. Furthermore, for every natural number  $n \geq 9$ , there exists an  $n$ -element lattice  $L$  such that  $\sigma(L) = 83$  and  $L$  is not planar.*

“ **$\sigma$ -few**” means “ $\sigma$  is at most 83”.

“ **$\sigma$ -many**” means “ $\sigma$  is strictly more than 83”.

### Theorem (Yet another formulation)

*Finite lattices with  $\sigma$ -many sublattices are planar.*

$$\sigma(\mathcal{A}) := |\text{Sub}(\mathcal{A})| \cdot 2^{8-n}, \quad \text{for a partial algebra.}$$

### Theorem (The main theorem reformulated)

*If  $L$  is a finite lattice such that  $\sigma(L) > 83$ , then  $L$  is planar. Furthermore, for every natural number  $n \geq 9$ , there exists an  $n$ -element lattice  $L$  such that  $\sigma(L) = 83$  and  $L$  is not planar.*

“ **$\sigma$ -few**” means “ $\sigma$  is at most 83”.

“ **$\sigma$ -many**” means “ $\sigma$  is strictly more than 83”.

### Theorem (Yet another formulation)

*Finite lattices with  $\sigma$ -many sublattices are planar.*

## Lemma (1st lemma)

If  $\mathcal{B} = (B, F_B)$  is a (weak partial) subalgebra of a finite (partial) algebra  $\mathcal{A} = (A, F_A)$ , then  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .

## Proof.

Let  $n = |B|$ ,  $k = |A| - |B|$ . If  $X \in \text{Sub}(\mathcal{A})$ , then  $X' := X \cap B \in \text{Sub}(\mathcal{B})$ .  $\#X' \leq |\text{Sub}(\mathcal{B})|$ . Each  $X'$  extends to a subset of  $A$  in  $2^k$  ways  $\Rightarrow |\text{Sub}(\mathcal{A})| \leq 2^k \cdot \#X' \leq 2^k \cdot |\text{Sub}(\mathcal{B})|$ . Dividing this by  $2^{k+n}$ , we get  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .  $\square$

If  $\mathcal{B} = (B, F_B)$  is a weak partial subalgebra of  $\mathcal{A} = (A, F_A)$  if  $B \subseteq A$ ,  $\text{Dom}(f_B) \subseteq B^2 \cap \text{Dom}(f_A)$  for every  $f \in F$  and  $f_B(x, y) = f_A(x, y)$  for all  $(x, y) \in \text{Dom}(f_B)$ .

## Lemma (1st lemma)

If  $\mathcal{B} = (B, F_B)$  is a (weak partial) subalgebra of a finite (partial) algebra  $\mathcal{A} = (A, F_A)$ , then  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .

## Proof.

Let  $n = |B|$ ,  $k = |A| - |B|$ . If  $X \in \text{Sub}(\mathcal{A})$ , then  $X' := X \cap B \in \text{Sub}(\mathcal{B})$ .  $\#X' \leq |\text{Sub}(\mathcal{B})|$ . Each  $X'$  extends to a subset of  $A$  in  $2^k$  ways  $\Rightarrow |\text{Sub}(\mathcal{A})| \leq 2^k \cdot \#X' \leq 2^k \cdot |\text{Sub}(\mathcal{B})|$ . Dividing this by  $2^{k+n}$ , we get  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .  $\square$

If  $\mathcal{B} = (B, F_B)$  is a weak partial subalgebra of  $\mathcal{A} = (A, F_A)$  if  $B \subseteq A$ ,  $\text{Dom}(f_B) \subseteq B^2 \cap \text{Dom}(f_A)$  for every  $f \in F$  and  $f_B(x, y) = f_A(x, y)$  for all  $(x, y) \in \text{Dom}(f_B)$ .

## Lemma (1st lemma)

If  $\mathcal{B} = (B, F_B)$  is a (weak partial) subalgebra of a finite (partial) algebra  $\mathcal{A} = (A, F_A)$ , then  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .

## Proof.

Let  $n = |B|$ ,  $k = |A| - |B|$ . If  $X \in \text{Sub}(\mathcal{A})$ , then  $X' := X \cap B \in \text{Sub}(\mathcal{B})$ .  $\#X' \leq |\text{Sub}(\mathcal{B})|$ . Each  $X'$  extends to a subset of  $A$  in  $2^k$  ways  $\Rightarrow |\text{Sub}(\mathcal{A})| \leq 2^k \cdot \#X' \leq 2^k \cdot |\text{Sub}(\mathcal{B})|$ . Dividing this by  $2^{k+n}$ , we get  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .  $\square$

If  $\mathcal{B} = (B, F_B)$  is a weak partial subalgebra of  $\mathcal{A} = (A, F_A)$  if  $B \subseteq A$ ,  $\text{Dom}(f_B) \subseteq B^2 \cap \text{Dom}(f_A)$  for every  $f \in F$  and  $f_B(x, y) = f_A(x, y)$  for all  $(x, y) \in \text{Dom}(f_B)$ .

## Lemma (1st lemma)

If  $\mathcal{B} = (B, F_B)$  is a (weak partial) subalgebra of a finite (partial) algebra  $\mathcal{A} = (A, F_A)$ , then  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .

## Proof.

Let  $n = |B|$ ,  $k = |A| - |B|$ . If  $X \in \text{Sub}(\mathcal{A})$ , then  $X' := X \cap B \in \text{Sub}(\mathcal{B})$ .  $\#X' \leq |\text{Sub}(\mathcal{B})|$ . Each  $X'$  extends to a subset of  $A$  in  $2^k$  ways  $\Rightarrow |\text{Sub}(\mathcal{A})| \leq 2^k \cdot \#X' \leq 2^k \cdot |\text{Sub}(\mathcal{B})|$ .  
Dividing this by  $2^{k+n}$ , we get  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ . □

If  $\mathcal{B} = (B, F_B)$  is a weak partial subalgebra of  $\mathcal{A} = (A, F_A)$  if  $B \subseteq A$ ,  $\text{Dom}(f_B) \subseteq B^2 \cap \text{Dom}(f_A)$  for every  $f \in F$  and  $f_B(x, y) = f_A(x, y)$  for all  $(x, y) \in \text{Dom}(f_B)$ .

## Lemma (1st lemma)

If  $\mathcal{B} = (B, F_B)$  is a (weak partial) subalgebra of a finite (partial) algebra  $\mathcal{A} = (A, F_A)$ , then  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .

## Proof.

Let  $n = |B|$ ,  $k = |A| - |B|$ . If  $X \in \text{Sub}(\mathcal{A})$ , then  $X' := X \cap B \in \text{Sub}(\mathcal{B})$ .  $\#X' \leq |\text{Sub}(\mathcal{B})|$ . Each  $X'$  extends to a subset of  $A$  in  $2^k$  ways  $\Rightarrow |\text{Sub}(\mathcal{A})| \leq 2^k \cdot \#X' \leq 2^k \cdot |\text{Sub}(\mathcal{B})|$ . Dividing this by  $2^{k+n}$ , we get  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .  $\square$

If  $\mathcal{B} = (B, F_B)$  is a weak partial subalgebra of  $\mathcal{A} = (A, F_A)$  if  $B \subseteq A$ ,  $\text{Dom}(f_B) \subseteq B^2 \cap \text{Dom}(f_A)$  for every  $f \in F$  and  $f_B(x, y) = f_A(x, y)$  for all  $(x, y) \in \text{Dom}(f_B)$ .

## Lemma (1st lemma)

If  $\mathcal{B} = (B, F_B)$  is a (weak partial) subalgebra of a finite (partial) algebra  $\mathcal{A} = (A, F_A)$ , then  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .

## Proof.

Let  $n = |B|$ ,  $k = |A| - |B|$ . If  $X \in \text{Sub}(\mathcal{A})$ , then  $X' := X \cap B \in \text{Sub}(\mathcal{B})$ .  $\#X' \leq |\text{Sub}(\mathcal{B})|$ . Each  $X'$  extends to a subset of  $A$  in  $2^k$  ways  $\Rightarrow |\text{Sub}(\mathcal{A})| \leq 2^k \cdot \#X' \leq 2^k \cdot |\text{Sub}(\mathcal{B})|$ . Dividing this by  $2^{k+n}$ , we get  $\sigma(\mathcal{A}) \leq \sigma(\mathcal{B})$ .  $\square$

If  $\mathcal{B} = (B, F_B)$  is a weak partial subalgebra of  $\mathcal{A} = (A, F_A)$  if  $B \subseteq A$ ,  $\text{Dom}(f_B) \subseteq B^2 \cap \text{Dom}(f_A)$  for every  $f \in F$  and  $f_B(x, y) = f_A(x, y)$  for all  $(x, y) \in \text{Dom}(f_B)$ .

By D. Kelly and I. Rival (1974), a finite lattice  $L$  is NON-planar iff one of the following lattices

$$\mathcal{L}_{KR} := \{A_n, E_n, F_n, G_n, H_n : n \geq 0\} \cup \{B, C, D\}$$

is sitting in  $L$  or in the dual of  $L$ .

We need not care with

“dual” since  $\text{Sub}(L) = \text{Sub}(\text{dual of } L)$ .

The plan is clear:

- First, compute  $\sigma(K)$  for each  $K \in \mathcal{L}_{KR}$ ,
- Second, apply the Kelly–Rival Theorem.

By D. Kelly and I. Rival (1974), a finite lattice  $L$  is NON-planar iff one of the following lattices

$$\mathcal{L}_{\text{KR}} := \{A_n, E_n, F_n, G_n, H_n : n \geq 0\} \cup \{B, C, D\}$$

is sitting in  $L$  or in the dual of  $L$ .  
 “dual” since  $\text{Sub}(L) = \text{Sub}(\text{dual of } L)$ .

We need not care with

The plan is clear:

- First, compute  $\sigma(K)$  for each  $K \in \mathcal{L}_{\text{KR}}$ ,
- Second, apply the Kelly–Rival Theorem.

By D. Kelly and I. Rival (1974), a finite lattice  $L$  is NON-planar iff one of the following lattices


$$\mathcal{L}_{\text{KR}} := \{A_n, E_n, F_n, G_n, H_n : n \geq 0\} \cup \{B, C, D\}$$


is sitting in  $L$  or in the dual of  $L$ .  
“dual” since  $\text{Sub}(L) = \text{Sub}(\text{dual of } L)$ .

We need not care with

The plan is clear:

- First, compute  $\sigma(K)$  for each  $K \in \mathcal{L}_{\text{KR}}$ ,
- Second, apply the Kelly–Rival Theorem.





## Lemma (2nd Lemma; on small Kelly–Rival lattices)

$$\begin{array}{llll} \sigma(A_0) = 74, & \sigma(B) = 54, & \sigma(C) = 68.5, & \sigma(D) = 76, \\ \sigma(E_0) = 60.5, & \sigma(F_0) = \underline{83}, & \sigma(G_0) = 54.25, & \\ \sigma(H_0) = 49.75, & \sigma(E_1) = 31.125, & \text{and } \sigma(F_1) = 41.125. & \end{array}$$

Using fences and  $K_5$ , there are arguments for the large Kelly–Rival lattices.

If  $L$  is nonplanar, then some  $K \in \mathcal{L}_{KR}$  is sitting in  $L$ , whereby our lemmas yield  $83 \geq \sigma(K) \geq \sigma(L)$ , i.e.,  $L$  has  $\sigma$ -few sublattices. Consequently, if  $L$  has  $\sigma$ -many sublattices, then it is planar. ???

**WRONG!** What does “sitting” mean? In the Kelly–Rival Theorem, “sitting” means “subposet”. But the first lemma is about subalgebras; this lemma is not applicable! Very tedious work is needed for each  $K \in \mathcal{L}_{KR}$ . We only outline it for  $F_0$ .

If  $L$  is nonplanar, then some  $K \in \mathcal{L}_{KR}$  is sitting in  $L$ , whereby our lemmas yield  $83 \geq \sigma(K) \geq \sigma(L)$ , i.e.,  $L$  has  $\sigma$ -few sublattices. Consequently, if  $L$  has  $\sigma$ -many sublattices, then it is planar. ???

**WRONG!** What does “sitting” mean? In the Kelly–Rival Theorem, “sitting” means “subposet”. But the first lemma is about subalgebras; this lemma is not applicable! Very tedious work is needed for each  $K \in \mathcal{L}_{KR}$ . We only outline it for  $F_0$ .


If  $L$  is nonplanar, then some  $K \in \mathcal{L}_{KR}$  is sitting in  $L$ , whereby our lemmas yield  $83 \geq \sigma(K) \geq \sigma(L)$ , i.e.,  $L$  has  $\sigma$ -few sublattices. Consequently, if  $L$  has  $\sigma$ -many sublattices, then it is planar. ???

**WRONG!** What does “sitting” mean? In the Kelly–Rival Theorem, “sitting” means “subposet”. But the first lemma is about subalgebras; this lemma is not applicable! Very tedious work is needed for each  $K \in \mathcal{L}_{KR}$ . We only outline it for  $F_0$ .


If  $L$  is nonplanar, then some  $K \in \mathcal{L}_{KR}$  is sitting in  $L$ , whereby our lemmas yield  $83 \geq \sigma(K) \geq \sigma(L)$ , i.e.,  $L$  has  $\sigma$ -few sublattices. Consequently, if  $L$  has  $\sigma$ -many sublattices, then it is planar. ???

**WRONG!** What does “sitting” mean? In the Kelly–Rival Theorem, “sitting” means “subposet”. But the first lemma is about subalgebras; this lemma is not applicable! Very tedious work is needed for each  $K \in \mathcal{L}_{KR}$ . We only outline it for  $F_0$ .

Assuming  $F_0$  is a subposet of  $L$ ; we want to show  $\sigma(L) \leq 83$ .



$$b \wedge c = e, \quad e \vee g = c, \quad c \vee d = a, \quad d \wedge e = f, \quad a \vee b = i, \quad f \wedge g = o. \quad (1)$$



(C1):  $b \vee c = i$ ; then  $b \vee g = b \vee e \vee g = b \vee c = i$  also holds.

(C1a):  $e \wedge g = o$ ; then  $b \wedge g = b \wedge c \wedge g = e \wedge g = o$ .

(C1a.1):  $d \vee e = a$ , then  $b \vee d = b \vee e \vee d = b \vee a = i$ .


(C1a.1a):  $c \wedge d = f$ ; then  $d \wedge g = d \wedge c \wedge g = f \wedge g = o$ .

(C1a.1a.1):  $a \wedge b = e$ ; then  $b \wedge d = b \wedge a \wedge d = e \wedge d = f$ .

(C1a.1a.1b):  $f \vee g =: x \neq c$ . Then  $x < c$  is new and  $e \vee x = c$

since  $c = e \vee g \leq e \vee x \leq c$ .  $\sigma(\text{weak part. subalgebra}) = 74.25$ .

Now,  $\sigma(\text{each of the 13 leaves}) \leq 83$ ; Lemma 1  $\Rightarrow \sigma(L) \leq 83\sqrt{}$ .



(C1):  $b \vee c = i$ ; then  $b \vee g = b \vee e \vee g = b \vee c = i$  also holds.

(C1a):  $e \wedge g = o$ ; then  $b \wedge g = b \wedge c \wedge g = e \wedge g = o$ .

(C1a.1):  $d \vee e = a$ , then  $b \vee d = b \vee e \vee d = b \vee a = i$ .


(C1a.1a):  $c \wedge d = f$ ; then  $d \wedge g = d \wedge c \wedge g = f \wedge g = o$ .

(C1a.1a.1):  $a \wedge b = e$ ; then  $b \wedge d = b \wedge a \wedge d = e \wedge d = f$ .

(C1a.1a.1b):  $f \vee g =: x \neq c$ . Then  $x < c$  is new and  $e \vee x = c$

since  $c = e \vee g \leq e \vee x \leq c$ .  $\sigma(\text{weak part. subalgebra}) = 74.25$ .

Now,  $\sigma(\text{each of the 13 leaves}) \leq 83$ ; Lemma 1  $\Rightarrow \sigma(L) \leq 83\sqrt{}$ .



(C1):  $b \vee c = i$ ; then  $b \vee g = b \vee e \vee g = b \vee c = i$  also holds.

(C1a):  $e \wedge g = o$ ; then  $b \wedge g = b \wedge c \wedge g = e \wedge g = o$ .

(C1a.1):  $d \vee e = a$ , then  $b \vee d = b \vee e \vee d = b \vee a = i$ .


(C1a.1a):  $c \wedge d = f$ ; then  $d \wedge g = d \wedge c \wedge g = f \wedge g = o$ .

(C1a.1a.1):  $a \wedge b = e$ ; then  $b \wedge d = b \wedge a \wedge d = e \wedge d = f$ .

(C1a.1a.1b):  $f \vee g =: x \neq c$ . Then  $x < c$  is new and  $e \vee x = c$

since  $c = e \vee g \leq e \vee x \leq c$ .  $\sigma(\text{weak part. subalgebra}) = 74.25$ .

Now,  $\sigma(\text{each of the 13 leaves}) \leq 83$ ; Lemma 1  $\Rightarrow \sigma(L) \leq 83\sqrt{}$ .



(C1):  $b \vee c = i$ ; then  $b \vee g = b \vee e \vee g = b \vee c = i$  also holds.

(C1a):  $e \wedge g = o$ ; then  $b \wedge g = b \wedge c \wedge g = e \wedge g = o$ .

(C1a.1):  $d \vee e = a$ , then  $b \vee d = b \vee e \vee d = b \vee a = i$ .

(C1a.1a):  $c \wedge d = f$ ; then  $d \wedge g = d \wedge c \wedge g = f \wedge g = o$ .

(C1a.1a.1):  $a \wedge b = e$ ; then  $b \wedge d = b \wedge a \wedge d = e \wedge d = f$ .

(C1a.1a.1b):  $f \vee g =: x \neq c$ . Then  $x < c$  is new and  $e \vee x = c$

since  $c = e \vee g \leq e \vee x \leq c$ .  $\sigma(\text{weak part. subalgebra}) = 74.25$ .

Now,  $\sigma(\text{each of the 13 leaves}) \leq 83$ ; Lemma 1  $\Rightarrow \sigma(L) \leq 83\sqrt{}$ .

$K \in \mathcal{L}_{KR}$	$B$	$C$	$D$	$E_0$	$E_1$	$F_0$	$F_0$	$G_0$	$H_0$
$ \{\text{leaves}\} $	11	12	5	37	5	13	19	24	67

Lattice theoretical considerations and some additional (small) parsing trees are necessary to exclude the “large” members of the Kelly–Rival List.

These slides, the paper, the computer program, and the input and output files are all available from

<http://www.math.u-szeged.hu/~czedli/>

THANK YOU FOR YOUR ATTENTION!

$K \in \mathcal{L}_{KR}$	$B$	$C$	$D$	$E_0$	$E_1$	$F_0$	$F_0$	$G_0$	$H_0$
$ \{\text{leaves}\} $	11	12	5	37	5	13	19	24	67

Lattice theoretical considerations and some additional (small) parsing trees are necessary to exclude the “large” members of the Kelly–Rival List.

These slides, the paper, the computer program, and the input and output files are all available from

<http://www.math.u-szeged.hu/~czedli/>

THANK YOU FOR YOUR ATTENTION!