

SUMS AND TOLERANCES OF LATTICES

GÁBOR CZÉDLI AND GEORGE GRÄTZER

Sums and *tolerances* are important concepts in lattice theory, and perhaps, with some exceptions, only in lattice theory.

1. AN EASY WAY TO SUMS

Given a congruence Θ of a lattice L , we can form the quotient lattice $K = L/\Theta$, and the Θ -blocks, which are lattices that form a K -indexed system $\{A \mid A \in K\}$. If we want to recreate L from the lattice K and the system $\{A \mid A \in K\}$, then we form the *sum* of a K -indexed system of lattices. Sum is also called *Płonka sum* or *Graczyńska sum*, and it is one of the main tools for studying products of lattice varieties; see, for instance, [1], [7], [8], [9], [12], [14]. For simplicity, here we consider complete lattices only.

For a K -indexed system of lattices, one has to describe how the summands are related. This is usually done by a pair of functors, one being the right adjoint of the other, see [4] for historical details. Now we offer a single functor, which is an easier to visualize. Let $L_1 = (L_1, \leq_1)$ and $L_2 = (L_2, \leq_2)$ be lattices. Roughly speaking, a relation $\rho \subseteq L_1 \times L_2$ will be called an *atop relation*, if taking disjoint copies of L_1 and L_2 and putting L_2 atop L_1 modulo ρ (that is, adding ρ to the union of \leq_1 and \leq_2), we obtain a complete lattice.

Theorem 1 ([4]). *The class \mathcal{C} of complete lattices, as objects, together with atop relations, as morphisms, the lattice orderings, as identities, and the usual relational product, as operation, constitute a category.*

As described in [4], summable systems of lattices can be defined as a functor from K to \mathcal{C} , and sums can be treated accordingly.

2. TOLERANCES AS HOMOMORPHIC IMAGES OF CONGRUENCES

By a *tolerance* we mean a reflexive, symmetric, compatible relation; see [2] for a survey. While tolerances are congruences in congruence permutable varieties, we have four arguments for their importance in Lattice Theory. Firstly, term functions of a finite lattice L are exactly those isotone functions that preserves all tolerances, see [13]. Secondly,

tolerances are, explicitly or implicitly present in several gluings of lattices, see [6], [11], and [12], for example.

By a *block* of a tolerance $\rho \subseteq L^2$, we mean a maximal subset X of L such that $X^2 \subseteq \rho$. Blocks are convex sublattices. Note that $(x, y) \in \rho$ iff there is a block X of ρ such that $x, y \in X$. If ρ is a tolerance of an algebra A , f is an n -ary operation of A , and X_1, \dots, X_n are blocks of ρ , then Zorn's lemma yields a block Y of ρ such that $f(x_1, \dots, x_n) \in Y$, for all $x_i \in X_i$. While this Y is not unique in general, the following theorem, which is the third argument, holds.

Theorem 2 ([3]). *Let ρ be a tolerance of a lattice L , and let L/ρ be the set of all blocks of ρ . Then Y above is unique; this makes L/ρ an algebra $(L/\rho, \vee, \wedge)$ in the obvious way. This $(L/\rho, \vee, \wedge)$ is a lattice.*

For an alternative approach to L/ρ see [10]. If Θ is a congruence of an algebra A and $\varphi: A \rightarrow B$ is a surjective homomorphism, then $\varphi(\Theta) = \{(\varphi(x), \varphi(y)) : (x, y) \in \Theta\}$ is clearly a tolerance of B . Our fourth argument is

Theorem 3 ([5]). *Let ρ be a tolerance of a lattice L . Then there exist a lattice M , a congruence Θ on M , and a surjective homomorphism $\varphi: M \rightarrow L$ such that $\rho = \varphi(\Theta)$.*

3. A LINK BETWEEN SUMS AND TOLERANCES

The first proof of Theorem 3 (for the finite case only) ran as follows: we took a sum of the blocks of ρ , which are lattices. However, in the talk we give a simpler proof.

REFERENCES

- [1] Bandelt, H.-J.; *Plonka sums of complete lattices*, Simon Stevin 55 (1981), no. 3, 169-171.
- [2] Chajda, I.: *Algebraic Theory of Tolerances*, Olomouc, 1991.
- [3] Czédli, G.: *Factor lattices by tolerances*, Acta Sci. Math. (Szeged), 44 (1982), 35-42.
- [4] Czédli, G.: *Sums of lattices and a relational category*, Order 26 (2009), 309-318.
- [5] Czédli, G.; Grätzer, G.: *Lattice tolerances and congruences*, Algebra Universalis, submitted.
- [6] Day, A.; Herrmann, C.: *Gluings of modular Lattices*, Order 5 (1988), 85-101.
- [7] Graczyńska, G.: *On the sum of double systems of lattices*, Proc. Colloq. Szeged (1975), North Holland 1977.
- [8] Graczyńska, G.; Grätzer, G.: *On double systems of lattices*, Demonstratio Math. 13 (1980), 743-747.
- [9] Grätzer, G.; Kelly, D.: *Products of lattice varieties*, Algebra Universalis 21 (1985), 33-45.

- [10] Grätzer, G; Wenzel, G.: *Notes on tolerance relations of lattices*, Acta Sci. Math. 54 (1990), 229-240.
- [11] Grygiel, J.: *The concept of gluing for lattices*, Czestochowa 2004, 116 pages.
- [12] Herrmann, C.: *S-verklebte Summen von Verbänden*, Math. Z. 130 (1973), 255-274.
- [13] Kindermann, M.: *Über die Äquivalenz von Ordnungspolynomvollständigkeit und Toleranzeinfachheit endlicher Verbände*, Proc. Conf. Klagenfurt 1978.
- [14] Romanowska, A.: *Building bisemilattices from lattices and semilattices*, Contributions to General Algebra, 2 (Klagenfurt, 1982), 343-358.

E-mail address: `czedli@math.u-szeged.hu`

UNIVERSITY OF SZEGED, BOLYAI INSTITUTE, SZEGED, HUNGARY-6720

E-mail address: `gratzer@me.com`

UNIVERSITY OF MANITOBA, WINNIPEG, MB R3T 2N2, CANADA