Coordinatization of finite join-distributive lattices

Gábor Czédli

ABSTRACT. Join-distributive lattices are finite, meet-semidistributive, and semimodular lattices. They are the same as lattices with unique irreducible decompositions introduced by R. P. Dilworth in 1940, and many alternative definitions and equivalent concepts have been discovered or rediscovered since then. Let L be a join-distributive lattice of length n, and let k denote the width of the set of join-irreducible elements of L. We prove that there exist k - 1 permutations acting on $\{1, \ldots, n\}$ such that the elements of L are coordinatized by k-tuples over $\{0, \ldots, n\}$, and the permutations determine which k-tuples are allowed. Since the concept of join-distributive lattices is equivalent to that of antimatroids and convex geometries, our result offers a coordinatization for these combinatorial structures.

1. Introduction

In 1940, R. P. Dilworth [12] introduced an important class of finite lattices. Recently, these lattices are called join-distributive. The concept of antimatroids, which are particular greedoids of B. Korte and L. Lovász [22] and [23], and that of convex geometries were introduced only much later by P. H. Edelman and R. E. Jamison [20], [13], and [15]. Join-distributive lattices, antimatroids, and convex geometries are equivalent concepts in a natural way, see Section 7.

Hence, though the majority of the paper belongs to lattice theory, the result we prove can also be interesting in combinatorics. Note that there were a lot of discoveries and rediscoveries of join-distributive lattices and the corresponding combinatorial structures; see B. Monjardet [24] and M. Stern [26] for surveys.

Target. Let L be a join-distributive lattice of length n. The poset (= partially ordered set) of join-irreducible elements of L is denoted by Ji L. Let C_1, \ldots, C_k be maximal chains of L such that Ji $L \subseteq C_1 \cup \cdots \cup C_k$. The collection of isomorphism classes of systems $\langle L; C_1, \ldots, C_k \rangle$ is denoted by JD(n, k). The symmetric group of degree n, which consists of all $\{1, \ldots, n\} \to \{1, \ldots, n\}$ permutations, is denoted by S_n . Our goal is to establish a bijection between JD(n, k) and S_n^{k-1} . This bijection describes L as subset of $\{0, 1, \ldots, n\}^k$ of k-tuples satisfying a system of equations determined by some member of S_n^{k-1} . This explains why we term our bijection as a "coordinatization". If k is small

²⁰¹⁰ Mathematics Subject Classification: Primary 06C10; secondary 05E99 and 52C99. Key words and phrases: Semimodular lattice, diamond-free lattice, planar lattice, Jordan-Hölder permutation, antimatroid, convex geometry, anti-exchange closure.

This research was supported by the NFSR of Hungary (OTKA), grant numbers K77432 and K83219, and by TÁMOP-4.2.1/B-09/1/KONV-2010-0005.

compared to n, then this coordinatization gives a very economic way to describe $\langle L; C_1, \ldots, C_k \rangle$ and, consequently, L with few data. Our coordinatization can easily be translated to the language of convex geometries and antimatroids. While the case k = 2 was settled, partly rediscovered, in G. Czédli and E. T. Schmidt [11]; the case k > 2 requires a more complex approach.

Outline. Section 2 contains the lattice theoretic prerequisites, and recalls the known characterizations of join-distributive lattices. Trajectories, which represent the main tool used in the paper, were introduced for the planar case in G. Czédli and E. T. Schmidt [10]. Section 3 is devoted to trajectories in arbitrary join-distributive lattices. With the help of trajectories, we develop a new approach to Jordan-Hölder permutations in Section 4. Our main result, the coordinatization theorem for join-distributive lattices, is formulated in Section 5. This theorem is proved in Section 6. Section 7 surveys antimatroids and convex geometries briefly. It also translates our coordinatization theorem to the language of combinatorics.

After an earlier version of this paper, http://arxiv.org/abs/1208.3517, Kira Adaricheva pointed out that the main result here is closely related to P.H. Edelman and R.E. Jamison [15, Theorem 5.2], which is formulated for convex geometries. This connection is analyzed in K. Adaricheva and G. Czédli [2], in which it is shown that our coordinatization result and the Edelman-Jamison description can be derived from each other in less than a page. However, not only the methods and the motivations of [15] and the present paper are entirely different, the results are not exactly the same; see [2] for comparison. In particular, our coordinatization represents a joindistributive lattice L as a meet-subsemilattice of the direct power of a chain, while [15] represents L as a join-subsemilattice of the powerset lattice of the same chain; see [15] for details. While [15] belongs to combinatorics, the coordinatization result is a logical "step" in a chain of purely lattice theoretical papers, starting from G. Grätzer and E. Knapp [17] and G. Grätzer and J.B. Nation [18], and including, among others, G. Czédli, L. Ozsvárt, and B. Udvari [8], G. Czédli and E. T. Schmidt [9], [11], and also the paper G. Czédli and E. T. Schmidt [10], which gives an application of lattice theory. Note that our method motivates a new characterization of join-distributive lattices, see [2], and implies some known characterizations, see Remark 2.2.

2. Preliminaries

The objective of this section is to give various descriptions for the lattices the present paper deals with. The *length* of an (n+1)-element chain is n, while the length of a lattice L, denoted by length L, is the supremum of {length C : Cis a chain of L}. A lattice is *trivial* if it consists of a single element. Let us agree that all lattices in this paper are either finite, or they are explicitly assumed to be of finite length. As usual, \prec stands for the covering relation:

 $x \prec y$ means that the interval [x, y] is 2-element. If $0 \prec a$, then a is an atom. A lattice L is semimodular if $x \prec y$ implies $x \lor z \prec y \lor z$, for all $x, y, z \in L$. An element is *meet-irreducible* if it has exactly one cover. The poset of these elements of L is denoted by Mi L. Note that $1 \notin \text{Mi } L$ and $0 \notin \text{Ji } L$. Since L is of finite length, each element $x \in L$ is of the form $x = \bigwedge Y$ for some $Y \subseteq \operatorname{Mi} L$. Note that $Y = \emptyset$ iff x = 1. The equation $x = \bigwedge Y$ is an *irredundant meet*decomposition of x if $Y \subseteq \operatorname{Mi} L$ and $x \neq \bigwedge Y'$ for every proper subset Y' of Y. If each $x \in L$ has only one irredundant meet-decomposition, then we say that L is a lattice with unique meet-irreducible decompositions. A diamond of L is a 5-element modular but not distributive sublattice M_3 of L. A diamond consists of its top, its bottom, and the rest of its elements form an antichain. If no such sublattice exists, then L is *diamond-free*. If S is a sublattice of Lsuch that, for all $x, y \in S$, $x \prec_S y$ implies $x \prec_L y$, then S is a cover-preserving sublattice of L. If S is a nonempty subset of L such that $x \lor y \in S$ for all $x, y \in S$, then S is a *join-subsemilattice* of L. For $x \in L$, the join of all covers of x is denoted by x^* . If, for all $x, y, z \in L$, $x \wedge y = x \wedge z$ implies $x \wedge y = x \wedge (y \vee z)$, then L is meet-semidistributive. If Ji L is the union of two chains, then L is *slim*. The next statement is known and gives a good understanding of join-distributive lattices within lattice theory. For further characterizations, see Section 7 here, see S.P. Avann [5], which is recalled in P.H. Edelman [14, Theorem 1.1], and see also M. Stern [26, Theorem 7.2.27].

Proposition 2.1. For a finite lattice L, the following properties are equivalent.

- (i) L is join-distributive, that is, semimodular and meet-semidistributive.
- (ii) L has unique meet-irreducible decompositions.
- (iii) For each $x \in L$, the interval $[x, x^*]$ is distributive.
- (iv) For each $x \in L$, the interval $[x, x^*]$ is boolean.
- (v) The length of each maximal chain of L equals |MiL|.
- (vi) L is semimodular and diamond-free.
- (vii) L is semimodular and has no cover-preserving diamond sublattice.
- (viii) L is a cover-preserving join-subsemilattice of a finite distributive lattice.

Now, we explain how Proposition 2.1 can be extracted from the literature. The equivalence of (ii) and (iii) above was proved by R. P. Dilworth [12]. D. Armstrong [4, Theorem 2.7] states (i) \Leftrightarrow (ii) \Leftrightarrow (iii) by extracting it from K. Adaricheva, V. A. Gorbunov, and V. I. Tumanov [3, Theorems 1.7 and 1.9], where the dual statement is given. We know (iii) \Leftrightarrow (v) \Leftrightarrow (vii) from M. Stern [26, Theorem 7.2.27], who attributes it to S. P. Avann [5] and [6]. The implication (vi) \Rightarrow (vii) is trivial. H. Abels [1, Theorem 3.9] contains (vii) \Leftrightarrow (viii). Next, as the fourth sentence in P. H. Edelman [13, Section 3] points out, (iii) \Leftrightarrow (iv) is practically trivial; the argument runs as follows. Assume that $[x, x^*]$ is distributive. Let a_1, \ldots, a_t be the covers of x. They are independent in $[x, x^*]$ by distributivity and G. Grätzer [16, Theorem 360]. Hence they generate a boolean sublattice B of length t and size 2^t , and $[x, x^*]$ is also of length t. Since $|\operatorname{Ji}([x, x^*])| = \operatorname{length}([x, x^*]) = t$ by [16, Corollary

112], we obtain from [16, Theorem 107] that $|[x, x^*]| \leq 2^t$. Thus $[x, x^*] = B$ is boolean, proving (iii) \Leftrightarrow (iv). Finally, suppose for a contradiction that (ii) \Rightarrow (vi) fails for some finite lattice L. Let $\{u < a_1 \parallel a_2 \parallel a_3 < v\}$ be a diamond in L, and let $R_i \subseteq \text{Mi } L$ such that $a_i = \bigwedge R_i$ for $i \in \{1, 2, 3\}$. For $i \in \{2, 3\}, R_1 \cup (R_i \setminus \uparrow v)$ has a subset T_i such that $\bigwedge T_i$ is an irredundant meet-decomposition of $u = a_1 \land a_i$. Clearly, $T_i \not\subseteq R_1$. If we had $T_2 = T_3$, then $T_2 \setminus R_1 = T_3 \setminus R_1$ would contain an element x, and $x \geq a_2 \lor a_3 = v$ would contradict $x \in R_2 \setminus \uparrow v$. This shows that (ii) fails in L, a contradiction.

Remark 2.2. The proof of our coordinatization result offers an alternative way to the implication (vii) \Rightarrow (i), even for lattices of finite length.

Remark 2.3. If L is a lattice of finite length, then each of conditions (i), (iii), ..., (viii) implies that L is finite. (Condition (ii) has not been investigated from this aspect.) This follows either from Propositions 2.1 and 6.1, or from Proposition 2.1 and H. Abels [1, Theorem 3.9]; see also Corollary 4.4.

3. Trajectories

The general assumption in Section 3 is that L is a semimodular lattice of finite length and without cover-preserving diamonds. A prime interval is a 2-element interval. A covering square of L is a cover-preserving 4-element boolean sublattice $S = \{a \land b, a, b, a \lor b\}$. The prime intervals $[a \land b, a]$ and $[b, a \lor b]$ are opposite sides of S, and so are the prime intervals $[a \land b, b]$ and $[a, a \lor b]$. The set of prime intervals of L is denoted by PrInt(L). If two prime intervals are opposite sides of the same covering square, then they are consecutive. As in G. Czédli and E. T. Schmidt [10], the transitive reflexive closure of this consecutiveness relation on PrInt(L) is an equivalence relation, and the blocks of this equivalence relation are the trajectories of L. The collection of all trajectories of L is denoted by Traj(L).

Lemma 3.1. Let L be a semimodular lattice of finite length, having no coverpreserving diamond, and let S be a cover-preserving join-subsemilattice of L. If S has a smallest element, then the following two statements hold.

(A) For each $R \in \operatorname{Traj}(S)$, there is a unique $T \in \operatorname{Traj}(L)$ such that $R \subseteq T$. (B) Let $\kappa \colon \operatorname{Traj}(L) \cup \{\varnothing\} \to \operatorname{Traj}(S) \cup \{\varnothing\}$, defined by $\kappa(T) = T \cap \operatorname{PrInt}(S)$. Then κ is a surjective map. If $\{0_L, 1_L\} \subseteq S$, then κ is a bijection.

Note that S in Lemma 3.1 is a lattice but not a sublattice of L in general.

Proof of Lemma 3.1(A). Denoting the meet in S by \wedge_S , let $\{a \wedge_S b, a, b, a \lor b\}$ be a covering square of S. Then a and b are incomparable, and both cover $a \wedge_S b$ in L since S is a cover-preserving subset of L. This yields $a \wedge_S b = a \wedge_L b$, and we conclude that the covering squares of S are also covering squares of L. This implies part (A) of the lemma.

If L_1 and L_2 are lattices, $\varphi: L_1 \to L_2$ is a join-homomorphism, and $\varphi(x) \preceq \varphi(y)$ holds for all $x, y \in L_1$ with $x \preceq y$, then φ is a cover-preserving join-homomorphism. The kernel of a cover-preserving join-homomorphism is a cover-preserving join-congruence. For a join-congruence $\Theta \subseteq L^2$ and a covering square $S = \{a \land b, a, b, a \lor b\}$, S is a Θ -forbidden covering square if the Θ -blocks $a/\Theta, b/\Theta$, and $(a \land b)/\Theta$ are pairwise distinct but $(a \lor b)/\Theta$ equals a/Θ or b/Θ . The following easy lemma was proved in G. Czédli and E. T. Schmidt [10] and [9, Lemma 6].

Lemma 3.2. Let Θ be a join-congruence of a semimodular lattice of finite length. Then Θ is cover-preserving iff L has no Θ -forbidden covering square.

The initial idea of the present paper is formulated in the next lemma.

Lemma 3.3. Let L be a semimodular lattice of finite length such that L contains no cover-preserving diamond. Then, for each maximal chain C of L and for each trajectory T of L, T contains exactly one prime interval of C.

Proof. Take a prime interval $\mathfrak{p} \in T$, and pick a maximal chain D of L such that $\mathfrak{p} \in \operatorname{PrInt}(D)$. Let $S = [C \cup D]_{\vee}$, the join-subsemilattice generated by $C \cup D$. We know from G. Czédli and E. T. Schmidt [10, Lemma 2.4] that S is a cover-preserving $\{0, 1\}$ -join-subsemilattice of L, and it is a slim semimodular lattice. (But S is not a sublattice of L in general.) It follows from [10, Lemmas 2.4 and 2.8] that there are a unique $\mathfrak{q} \in \operatorname{PrInt}(C)$ and a unique trajectory R of S such that \mathfrak{p} and \mathfrak{q} belong to R. By Lemma 3.1(A), L has a trajectory R' such that $R \subseteq R'$. Since $p \in R' \cap T$, we obtain that T = R' contains \mathfrak{q} . This proves the existence part.

Note that, instead of [10], one could use H. Abels [1, Corollary 3.3] to prove the existence part. Actually, some ideas of [1] were rediscovered in [10]. However, since the concept of trajectories comes from [10], it is more convenient to reference [10].

We prove the uniqueness by contradiction. Suppose that $a \prec b \leq c \prec d$ such that [a, b] and [c, d] belong to the same trajectory T of L. Then there exists a sequence $[a, b] = [x_0, y_0], [x_1, y_1], \ldots, [x_t, y_t] = [c, d]$ of prime intervals such that $H_u = \{x_{u-1}, y_{u-1}, x_u, y_u\}$ is a covering square for $u \in \{1, \ldots, t\}$. Pick a maximal chain $0 = b_0 \prec b_1 \prec \cdots \prec b_s = b$ in the interval [0, b], and consider the join-homomorphisms $\psi_m \colon L \to L$, defined by $\psi_m(z) = b_m \lor z$, for $m \in \{0, \ldots, s\}$. Let Θ_m stand for Ker ψ_m , for $m \in \{0, \ldots, s\}$. By semimodularity, the ψ_m are cover-preserving join-homomorphisms. Since $\psi_s(x_0) = b = \psi_s(y_0)$ but $\psi_s(x_t) = c \neq d = \psi_s(y_t)$, there is a smallest $i \in \{1, \ldots, t\}$ such that $\psi_s(x_i) \neq \psi_s(y_i)$. That is, $\langle x_{i-1}, y_{i-1} \rangle \in \Theta_s$ but $\langle x_i, y_i \rangle \notin \Theta_s$. Clearly, x_i is the bottom of the covering square H_i , while y_{i-1} is its top.

The restriction of a relation ρ to a subset X will be denoted by $\rho|_X$. The equality relation on X is denoted by ω_X or ω . Since $\Theta_0|_{H_i} = \omega_{H_i}$ but $\langle x_{i-1}, y_{i-1} \rangle \in \Theta_s|_{H_i}$, there is a smallest j such that $\langle x_{i-1}, y_{i-1} \rangle \in \Theta_j|_{H_i}$. However, $\langle x_i, y_i \rangle \notin \Theta_j|_{H_i}$ since otherwise $b_j \vee x_i = \psi_j(x_i) = \psi_j(y_i) = b_j \vee y_i$ together with $b_j \leq b_s$ would imply $\psi_s(x_i) = \psi_s(y_i)$, which would contradict $\langle x_i, y_i \rangle \notin \Theta_s$.

Next, to simplify our notation, let

$$\alpha = \psi_{j-1}(x_i), \qquad \beta = \psi_{j-1}(y_i), \qquad \gamma = \psi_{j-1}(x_{i-1}), \qquad \delta = \psi_{j-1}(y_{i-1}), \\ \alpha' = \psi_j(x_i), \qquad \beta' = \psi_j(y_i), \qquad \gamma' = \psi_j(x_{i-1}), \qquad \delta' = \psi_j(y_{i-1}).$$

By the minimality of j, $|\{\alpha, \beta, \gamma, \delta\}| = 4$. Hence, $\{\alpha, \beta, \gamma, \delta\} = \psi_{j-1}(H_i)$ is a covering square with bottom α and top δ in the filter $\uparrow b_{j-1} = [b_{j-1}, 1]$ since ψ_{j-1} is a cover-preserving join-homomorphism. Consider the cover-preserving join-homomorphism $\varphi \colon \uparrow b_{j-1} \to \uparrow b_{j-1}$, defined by $\varphi(z) = b_j \lor z$. Denote the height function on $\uparrow b_{j-1}$ by h. Since b_j is an atom of the filter $\uparrow b_{j-1}$, semimodularity implies that $h(z) \leq h(\varphi(z)) \leq h(z) + 1$ holds for all $z \in \uparrow b_{j-1}$. By definitions, $\varphi(\alpha) = \alpha'$, $\varphi(\beta) = \beta'$, $\varphi(\gamma) = \gamma'$, and $\varphi(\delta) = \delta'$, and we also have $\gamma' = \delta'$ and $\alpha' \neq \beta'$. Let $\Phi = \operatorname{Ker} \varphi$. It is a cover-preserving join-congruence, and Lemma 3.2 yields that $\{\beta, \gamma\} \subseteq \delta/\Phi$ but $\alpha \notin \delta/\Phi$. Thus $\alpha' \neq \beta' = \gamma' = \delta'$. Actually, $\alpha' \prec \beta' = \gamma' = \delta'$ since φ is cover-preserving.

Since $\beta \neq \gamma$, we have $\langle \beta', \gamma' \rangle \neq \langle \beta, \gamma \rangle$. Hence we can assume $\beta' \neq \beta$, and we obtain $\beta \prec \beta'$. Using $h(\gamma) + 1 = h(\beta) + 1 = h(\beta') = h(\gamma')$, we obtain $\gamma \prec \gamma'$. Now $h(\delta') = h(\beta') = h(\beta) + 1 = h(\delta)$, together with $\delta \leq \delta'$, yields $\delta' = \delta$. We have $\alpha' \preceq \beta' = \delta' = \delta$ since φ is cover-preserving. Hence $\alpha' \neq \alpha$, and we obtain $\alpha \prec \alpha'$.

The previous relations imply $b_j \leq \delta$, $b_j \not\leq \alpha$, $b_j \not\leq \beta$, and $b_j \not\leq \gamma$. Since b_j is an atom in the filter $\uparrow b_{j-1}$, we have $\alpha \prec \alpha \lor b_j = \varphi(\alpha) = \alpha'$. We obtain $\alpha' \not\leq \beta$ from $b_j \not\leq \beta$, and we obtain $\alpha' \not\leq \gamma$ similarly. Hence β , α' , and γ are three different covers of α in the interval $[\alpha, \delta]$ of length 2. Thus $\{\alpha, \beta, \alpha', \gamma, \delta\}$ is a cover-preserving diamond of L, which is a contradiction.

Now we are in the position to complete the proof of Lemma 3.1.

Proof of Lemma 3.1(B). Let C be a map from PrInt(S) to the set of maximal chains of S such that, for every $\mathfrak{p} \in PrInt(S), \mathfrak{p} \in PrInt(C(\mathfrak{p}))$.

To show that κ is a map from $\operatorname{Traj}(L)$ to $\operatorname{Traj}(S)$, let $T \in \operatorname{Traj}(L)$ such that $T \cap \operatorname{PrInt}(S) \neq \emptyset$. Pick a prime interval $\mathfrak{p} \in T \cap \operatorname{PrInt}(S)$, and let $R \in \operatorname{Traj}(S)$ be the unique trajectory containing \mathfrak{p} . At present, we know that

$$T \in \operatorname{Traj}(L), R \in \operatorname{Traj}(S), \text{ and } \mathfrak{p} \in T \cap R \cap \operatorname{PrInt}(S),$$
 (3.1)

and this will be the only assumption on T and R we use in the rest of the present paragraph. By Lemma 3.1(A), there is an $R' \in \operatorname{Traj}(L)$ such that $R \subseteq R'$. Since $\mathfrak{p} \in T \cap R'$, we have $R \subseteq R' = T$, which implies $R \subseteq \kappa(T)$. To show the converse inclusion, let $\mathfrak{q} \in \kappa(T) = T \cap \operatorname{PrInt}(S)$. Applying the existence part of Lemma 3.3 to S, we obtain an $\mathfrak{r} \in R \cap \operatorname{PrInt}(C(\mathfrak{q}))$. Since $R \subseteq T$, both \mathfrak{r} and \mathfrak{q} belong to T. Thus the uniqueness part of Lemma 3.3 gives $\mathfrak{q} = \mathfrak{r} \in R$. Hence, $R = \kappa(T)$, and κ is a map from $\operatorname{Traj}(L)$ to $\operatorname{Traj}(S)$.

Since (3.1) implies $R = \kappa(T)$, κ is surjective.

Finally, let $\{0_L, 1_L\} \subseteq S$. Then S contains a maximal chain X of L. Assume that $T_1, T_2 \in \text{Traj}(L)$ such that $\kappa(T_1) = \kappa(T_2)$. By Lemma 3.3, there is a unique $\mathfrak{q} \in T_1 \cap \text{PrInt}(X)$. We have $\mathfrak{q} \in T_1 \cap \text{PrInt}(S) = \kappa(T_1) = \kappa(T_2) \subseteq T_2$. Hence $T_1 \cap T_2 \neq \emptyset$, and we conclude $T_1 = T_2$. Consequently, κ is injective, and we obtain that it is bijective.

4. Jordan-Hölder permutations

Any two maximal chains of a semimodular lattice of length n determine a so-called Jordan-Hölder permutation on the set $\{1, \ldots, n\}$. This was first stated by R. P. Stanley [25]; see also H. Abels [1] for further developments. Independently, the same permutations emerged in G. Grätzer and J. B. Nation [18]. The Jordan-Hölder permutations were rediscovered in G. Czédli and E. T. Schmidt [10], and were successfully applied to add a uniqueness part to the classical Jordan-Hölder Theorem for groups. As an excuse for this rediscovery, note that some results we need here were proved in [10] and the subsequent G. Czédli and E. T. Schmidt [11]. In [11], there are three equivalent definitions for Jordan-Hölder permutations. Here we combine the treatment given in [10] and [11] with Lemma 3.3. As opposed to H. Abels [1], we always assume that L has no cover-preserving diamond.

Definition 4.1. Let *L* be a semimodular lattice of length *n*, and assume that *L* has no cover-preserving diamond. Let $C = \{0 = c_0 \prec c_1 \prec \cdots \prec c_n = 1\}$ and $D = \{0 = d_0 \prec d_1 \prec \cdots \prec d_n = 1\}$ be maximal chains of *L*. Then the Jordan-Hölder permutation $\pi_{CD}: \{1, \ldots, n\} \rightarrow \{1, \ldots, n\}$ is defined by $\pi_{CD}(i) = j$ iff $[c_{i-1}, c_i]$ and $[d_{j-1}, d_j]$ belong to the same trajectory of *L*.

It is clear from Lemma 3.3 that π_{CD} above is a permutation. The next lemma shows that our definition of π_{CD} is the same as that of H. Abels [1, (3.1)], provided L has no cover-preserving diamond.

Lemma 4.2. Let L, C, and D be as in Definition 4.1. Then, for $i \in \{1, \ldots, n\}$,

$$\pi_{CD}(i) = \min\{j : c_{i-1} \lor d_j = c_i \lor d_j\}.$$

The permutation π_{CD} equals the identity permutation id iff C = D. We have $c_{i-1} \vee d_j = c_i \vee d_j$ iff $\pi_{CD}(i) \leq j$.

Proof. It follows from Lemma 3.1 that $\pi_{CD}(i)$ can be computed in $[C \cup D]_{\vee}$. Hence the first part of the statement can be extracted from G. Czédli and E. T. Schmidt [11, Definition 2.5] or, more easily, from G. Czédli, L. Ozsvárt, and B. Udvari [8, Definition 2.4]. Now that we know that our π_{CD} is the same as defined in H. Abels [1], the middle part follows from [1, 3.5.(a)]. It also follows from [11, Theorem 3.3] or [8, Lemma 7.2]. The last part is a trivial consequence of the first part.

While the following lemma needs a proof in H. Abels [1, Theorem 3.9(f)], Definition 4.1 in our setting makes it obvious. We compose permutations and maps from right to left, that is, $(f \circ g)(x) = f(g(x))$.

Lemma 4.3. Let L be a semimodular lattice of finite length and without coverpreserving diamonds, and let C, D, and E be maximal chains of L. Then the following hold.

- (i) $\pi_{CC} = \text{id}$, the identity map. (ii) $\pi_{CD} = \pi_{DC}^{-1}$
- (iii) $\pi_{DE} \circ \pi_{CD} = \pi_{CE}$.

Equivalently, the lemma above asserts that the maximal chains of L with singleton hom-sets hom $(C, D) = \{\pi_{CD}\}$ form a category, namely, a groupoid. We do not recall further details since although this category (equipped with the weak Bruhat order) determines L by D.S. Herscovici [19], it is rather large and complicated for our purposes. The best way for coordinatization is offered by lattice theory.

To give a short illustration of the strength of Lemma 4.3, we prove the following corollary even if it is known; see Remark 2.3.

Corollary 4.4. If L is a semimodular lattice of finite length and L has no cover-preserving diamonds, then L is finite.

Proof. Pick a maximal chain C in L. By Lemma 4.3, if $\pi_{CD} = \pi_{CE}$, then id $= \pi_{CD} \circ \pi_{CD}^{-1} = \pi_{CE} \circ \pi_{DC} = \pi_{DE}$, and Lemma 4.2 implies D = E. Hence, L has only finitely many maximal chains. Thus L is finite.

5. The main result

We always assume that n belongs to $\mathbb{N} = \{1, 2, 3, ...\}$ and, unless otherwise specified, $k \in \{2, 3, ...\}$. For a structure \mathfrak{A} , the class of structures isomorphic to \mathfrak{A} is denoted by I \mathfrak{A} . As usual, S_n stands for the group of permutations of the set $\{1, ..., n\}$. Consider the class

 $JD(n,k) = \{ \mathbf{I} \langle L; C_1, \dots, C_k \rangle : L \text{ is a join-distributive lattice of length } n, \text{ and} \\ C_1, \dots, C_k \text{ are maximal chains of } L \text{ such that } Ji L \subseteq C_1 \cup \dots C_k \}.$

We define a map

$$\xi \colon \mathrm{JD}(n,k) \to S_n^{k-1} \text{ by } \mathbf{I}\langle L; C_1, \dots, C_k \rangle \mapsto \langle \pi_{C_1 C_2}, \pi_{C_1 C_3}, \dots, \pi_{C_1 C_k} \rangle.$$
(5.1)

If $\vec{\pi} = \langle \pi_{12}, \ldots, \pi_{1k} \rangle \in S_n^{k-1}$, then by the corresponding *extended vector* we mean the k^2 -tuple $\langle \pi_{ij} : i, j \in \{1, \ldots, k\} \rangle$, where π_{11} is the identity permutation and $\pi_{ij} = \pi_{1j} \circ \pi_{1i}^{-1}$. In general,

 π_{ij} is always understood as $\pi_{1j} \circ \pi_{1i}^{-1}$, (5.2)

even if this is not emphasized all the time.

Definition 5.1. By an *eligible* $\vec{\pi}$ -tuple we mean a k-tuple $\vec{x} = \langle x_1, \ldots, x_k \rangle \in \{0, 1, \ldots, n\}^k$ such that $\pi_{ij}(x_i + 1) \ge x_j + 1$ holds for all $i, j \in \{1, \ldots, k\}$ such that $x_i < n$. The set of eligible $\vec{\pi}$ -tuples is denoted by $L(\vec{\pi})$. It is a poset with respect to the componentwise order: $\vec{x} \le \vec{y}$ means that, for all $i \in \{1, \ldots, k\}$, $x_i \le y_i$.

For $i \in \{1, \ldots, k\}$, an eligible $\vec{\pi}$ -tuple \vec{x} is *initial in its i-th component* if for all $\vec{y} \in L(\vec{\pi}), x_i = y_i$ implies $\vec{x} \leq \vec{y}$. Let $C_i(\vec{\pi})$ be the set of all eligible $\vec{\pi}$ -tuples that are initial in their *i*-th component. Now we are in the position to define a map

$$\eta: S_n^{k-1} \to \mathrm{JD}(n,k) \text{ by } \vec{\pi} \mapsto \mathbf{I}\langle L(\vec{\pi}); C_1(\vec{\pi}), \dots, C_k(\vec{\pi}) \rangle.$$
 (5.3)

It is not obvious that $\eta(\vec{\pi}) \in JD(n,k)$, but we will prove it soon. Now, we can formulate our main result as follows.

Theorem 5.2. The maps ξ and η are reciprocal bijections between JD(n,k) and S_n^{k-1} .

This theorem gives the desired coordinatization since each $\mathbf{I}\langle L; C_1, \ldots, C_k \rangle$ from JD(n, k) is described by its ξ -image, that is, by k - 1 permutations. The elements of $\langle L; C_1, \ldots, C_k \rangle$ correspond to k-tuples over $\{0, \ldots, n\}$, and the k - 1 permutations specify which k-tuples occur.

Problem 5.3. It would be desirable to characterize those pairs $\langle \vec{\pi}, \vec{\sigma} \rangle$ of (k - 1)-tuples of permutations of S_n for which the lattice part of $\eta(\vec{\pi})$ (without the k chains) coincides with that of $\eta(\vec{\sigma})$. However, in spite of the theory developed in D. S. Herscovici [19], we do not expect an elegant characterization. While the requested characterization for k = 2 is known from G. Czédli and E. T. Schmidt [11] and it was used in G. Czédli, L. Ozsvárt, and B. Udvari [8], even the case k = 3 seems to be quite complicated; this is witnessed by the following example.

Example 5.4. Let $L = \{0, a, b, c, a \lor b, a \lor c, b \lor c, 1\}$ be the 8-element boolean lattice. Consider the maximal chains $C_1 = \{0, a, a \lor b, 1\}, C_2 = \{0, b, a \lor b, 1\}, C_3 = \{0, c, b \lor c, 1\}$, and also the maximal chains $C'_1 = \{0, a, a \lor b, 1\}, C'_2 = \{0, b, b \lor c, 1\}$, and $C'_3 = \{0, c, a \lor c, 1\}$. Then

$$\pi_{C_1C_2} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}, \qquad \pi_{C_1C_3} = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix},$$
$$\pi_{C'_1C'_2} = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}, \qquad \pi_{C'_1C'_3} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}.$$

Let $\vec{\pi} = \langle \pi_{C_1C_2}, \pi_{C_1C_3} \rangle$ and $\vec{\sigma} = \langle \pi_{C'_1C'_2}, \pi_{C'_1C'_3} \rangle$. These two vectors look very different since $\vec{\pi}$ consists of transpositions while $\vec{\sigma}$ does not; furthermore, the second component of $\vec{\sigma}$ is the inverse of the first component, while this is not so for $\vec{\pi}$. However, by construction and Theorem 5.2, $\vec{\pi}$ and $\vec{\sigma}$ determine the same lattice L.

6. Proving the main result

Some of the auxiliary statements we will prove are valid under a seemingly weaker assumption than requiring $\mathbf{IL} \in \mathrm{JD}(n,k)$. Therefore, in accordance with Remark 2.3, we prove a (seemingly) stronger statement with little extra effort. To do so, consider the class

$$CDF(n,k) = \{ \mathbf{I} \langle L; C_1, \dots, C_k \rangle : L \text{ is semimodular of length } n,$$

 $L \text{ contains no cover-preserving diamond, } C_1, \dots, C_k$
are maximal chains of L , and $\text{Ji } L \subseteq C_1 \cup \dots \cup C_k. \}$

This notation comes from "Cover-preserving Diamond-Free". Although we know from Proposition 2.1 and Remark 2.3 that CDF(n, k) equals JD(n, k), we will give a new proof for this equality. We will use only the obvious $\text{JD}(n, k) \subseteq \text{CDF}(n, k)$. Keeping the notation of the original map, we extend its range as follows:

 $\xi \colon \mathrm{CDF}(n,k) \to S_n^{k-1}$, where $\mathbf{I}\langle L; C_1, \ldots, C_k \rangle \mapsto \langle \pi_{C_1C_2}, \pi_{C_1C_3}, \ldots, \pi_{C_1C_k} \rangle$.

Our aim in this section is to prove the following statement, which implies Theorem 5.2 and harmonizes with Remark 2.3.

Proposition 6.1.

(A) CDF(n,k) = JD(n,k).

(B) The maps ξ and η are reciprocal bijections.

The proof of Proposition 6.1 will need the following lemma.

Lemma 6.2. If $\vec{\pi} \in S_n^{k-1}$, then $\eta(\vec{\pi}) \in JD(n,k)$.

Proof. Convention (5.2) should be kept in mind. Clearly, $\langle n, \ldots, n \rangle \in L(\vec{\pi})$. Hence $L(\vec{\pi})$ has a top element. Let $\vec{x} = \langle x_1, \ldots, x_k \rangle$ and $\vec{y} = \langle y_1, \ldots, y_k \rangle$ belong to $L(\vec{\pi})$, and let $z_i = x_i \wedge y_i = \min\{x_i, y_i\}$, for $i \in \{1, \ldots, k\}$. Assume that $j \in \{1, \ldots, k\}$ such that $z_j < n$. Since $\vec{x}, \vec{y} \in L(\vec{\pi})$, we have $\pi_{jt}(x_j+1) \ge x_t+1$ and $\pi_{jt}(y_j+1) \ge y_t+1$, for every $t \in \{1, \ldots, k\}$. Let, say, $z_j = x_j$. We obtain $\pi_{jt}(z_j+1) = \pi_{jt}(x_j+1) \ge x_t+1 \ge z_t+1$. Hence $\vec{z} = \langle z_1, \ldots, z_k \rangle \in L(\vec{\pi})$, and we conclude that $L(\vec{\pi})$ is a lattice.

If we had $\vec{x} = \langle n, x_2, \ldots, x_k \rangle \in L(\vec{\pi})$ and $x_i \neq n$ for some $i \in \{2, \ldots, k\}$, then $\pi_{i1}(x_i + 1) \leq n < x_1 + 1$ would contradict $\vec{x} \in L(\vec{\pi})$. Hence $\langle n, \ldots, n \rangle$ is the only vector in $L(\vec{\pi})$ with first component n, and we conclude that this vector belongs to $C_1(\vec{\pi})$. Furthermore, since the first component of an eligible $\vec{\pi}$ -tuple plays no special role,

$$L(\vec{\pi}) \setminus \{\langle n, \dots, n \rangle\} \subseteq \{0, \dots, n-1\}^k.$$

$$(6.1)$$

Next, assume that $\vec{y} = \langle y_1, \ldots, y_k \rangle$ and $\vec{z} = \langle z_1, \ldots, z_k \rangle$ belong to $C_1(\vec{\pi})$ such that $y_1 \leq z_1$. Since $L(\vec{\pi})$ is meet-closed, $\vec{y} \wedge \vec{z} = \langle y_1, y_2 \wedge z_2, \ldots, y_k \wedge z_k \rangle \in L(\vec{\pi})$. Since \vec{y} is initial in its first component, $\vec{y} \leq \vec{y} \wedge \vec{z}$, which gives $\vec{y} \leq \vec{z}$. Thus $C_1(\vec{\pi})$ is a chain.

A vector \vec{x} in $\{1, \ldots, n\}^k$ is a $\vec{\pi}$ -orbit, if $\pi_{ij}(x_i) = x_j$ for all $i, j \in \{1, \ldots, k\}$. Equivalently, if it is of the form

$$\vec{x} = \langle \pi_{i1}(b), \dots, \pi_{ik}(b) \rangle$$
(6.2)

for some $i \in \{1, \ldots, k\}$ and $b \in \{1, \ldots, n\}$. A $\vec{\pi}$ -orbit need not belong to $L(\vec{\pi})$. A vector $\vec{y} = \langle y_1, \ldots, y_k \rangle$ in $\{0, \ldots, n-1\}^k$ is suborbital with respect to $\vec{\pi}$ if

 $\langle y_1 + 1, \dots, y_k + 1 \rangle$ is a $\vec{\pi}$ -orbit. (6.3)

Clearly, suborbital vectors belong to $L(\vec{\pi})$. For each $b \in \{0, \ldots, n-1\}$, (6.2), applied for i = 1, shows that there exists a vector $\vec{x} \in L(\vec{\pi})$ whose first component is b. Let \vec{y} be the meet of all these vectors. Clearly, $y_1 = b$ and $\vec{y} \in C_1(\vec{\pi})$. This shows that $C_1(\vec{\pi})$ is a chain of length n, and so are the $C_i(\vec{\pi})$ for $i \in \{1, \ldots, k\}$.

Next, let $B(\vec{\pi})$ denote the set of suborbital vectors with respect to $\vec{\pi}$. We have $B(\vec{\pi}) \subseteq L(\vec{\pi})$. By (6.2), each $b \in \{1, \ldots, n\}$ is the *i*-th component of exactly one $\vec{\pi}$ -orbit. Therefore,

$$|B(\vec{\pi})| = n. \tag{6.4}$$

We assert that

for each $\vec{x} \in L(\vec{\pi})$, there is a unique $U \subseteq B(\vec{\pi})$ such

that $\vec{x} = \bigwedge U$ is an irredundant meet decomposition. (6.5)

Let $\vec{x} = \langle x_1, \ldots, x_k \rangle \in L(\vec{\pi}) \setminus \{ \langle n, \ldots, n \rangle \}$, and keep (6.1) in mind. For $i \in \{1, \ldots, k\}$, let

$$\vec{y}^{(i)} = \langle \pi_{i1}(x_i+1) - 1, \dots, \pi_{ik}(x_i+1) - 1 \rangle.$$

It belongs to $B(\vec{\pi})$, whence $\vec{y}^{(i)} \in L(\vec{\pi})$. Since \vec{x} is also in $L(\vec{\pi})$, for $j \in \{1, \ldots, k\}$ we have $x_j = x_j + 1 - 1 \le \pi_{ij}(x_i + 1) - 1 = y_j^{(i)}$. Hence $\vec{x} \le \vec{y}^{(i)}$, and we conclude $\vec{x} \le \bigwedge \{\vec{y}^{(i)} : i \in \{1, \ldots, k\}\}$. The converse inequality also holds since $y_i^{(i)} = \pi_{ii}(x_i + 1) - 1 = x_i$ for $i \in \{1, \ldots, k\}$. That is, $\vec{x} = \bigwedge \{\vec{y}^{(i)} : i \in \{1, \ldots, k\}\}$. Since the meetands here are not necessarily distinct, we let

$$J = \{ j \in \{1, \dots, k\} : \vec{y}^{(i)} \neq \vec{y}^{(j)} \text{ for all } i < j \}.$$

Clearly, $\vec{x} = \bigwedge \{\vec{y}^{(j)} : j \in J\}$. We assert that if $V \subseteq B(\vec{\pi})$ and $\vec{x} = \bigwedge V$, then $\{\vec{y}^{(j)} : j \in J\} \subseteq V$. To show this, let $j \in J$. Since the *j*-th components of the vectors of V form a chain and the meet of these components equals x_j , there exists a $\vec{u} \in V$ such that $u_j = x_j$. The *j*-th component of $\vec{y}^{(j)}$ is also x_j by definition. Since any two orbital vectors with a common *j*-th component are equal, we obtain $\vec{y}^{(j)} = \vec{u} \in V$. Thus we obtain $\{\vec{y}^{(j)} : j \in J\} \subseteq V$. Since the $\vec{y}^{(j)}$, for $j \in J$, are pairwise distinct, we can take $U = \{\vec{y}^{(j)} : j \in J\}$, and we conclude (6.5).

Now, let $\vec{a} \in \operatorname{Mi}(L(\vec{\pi}))$. Since $\vec{a} = \vec{b}^{(1)} \wedge \cdots \wedge \vec{b}^{(s)}$ for appropriate $\vec{b}^{(1)}, \ldots, \vec{b}^{(s)}$ in $B(\vec{\pi})$ by (6.5), we obtain s = 1 and $\vec{a} = \vec{b}^{(1)} \in B(\vec{\pi})$. That is, $\operatorname{Mi}(L(\vec{\pi})) \subseteq B(\vec{\pi})$. For the sake of contradiction, suppose $B(\vec{\pi}) \setminus \operatorname{Mi}(L(\vec{\pi})) \neq \emptyset$, and let $\vec{b} \in B(\vec{\pi}) \setminus \operatorname{Mi}(L(\vec{\pi}))$. Observe that $U' = \{\vec{b}\}$ gives an irredundant meet-decomposition according to (6.5). There are a minimal $t \in \mathbb{N}$ and $\vec{w}^{(1)}, \ldots, \vec{w}^{(t)} \in \operatorname{Mi}(L(\vec{\pi}))$ such that $\vec{b} = \vec{w}^{(1)} \wedge \cdots \wedge \vec{w}^{(t)}$. This meet is irredundant by the minimality of t, and $t \geq 2$ since $\vec{b} \neq \langle n, \ldots, n \rangle$ and $\vec{b} \notin \operatorname{Mi}(L(\vec{\pi}))$. However, $\vec{w}^{(1)}, \ldots, \vec{w}^{(t)} \in B(\vec{\pi})$ since $\operatorname{Mi}(L(\vec{\pi})) \subseteq B(\vec{\pi})$. Thus $U'' = \{\vec{w}^{(1)}, \ldots, \vec{w}^{(t)}\}$ also gives an irredundant meet-decomposition according to (6.5). This is a contradiction since $U' \neq U''$. Hence,

$$\operatorname{Mi}\left(L(\vec{\pi})\right) = B(\vec{\pi}),\tag{6.6}$$

and $L(\vec{\pi})$ has unique meet-irreducible decompositions. Thus we conclude from Proposition 2.1 that $L(\vec{\pi})$ is join-distributive.

Next, we have to show that $C_i(\vec{\pi})$ is a maximal chain for $i \in \{1, \ldots, k\}$. Since $|C_i(\vec{\pi})| = n + 1$, it suffices to show that $L(\vec{\pi})$ is of length at most n. To prove this by contradiction, suppose $H = \{h_0 \prec h_1 \prec \cdots \prec h_{n+1}\}$ is a chain of $L(\vec{\pi})$. Let $W_i = \operatorname{Mi}(L(\vec{\pi})) \cap \uparrow h_i$. Clearly, $W_0 \supseteq W_1 \supseteq \cdots \supseteq W_{n+1}$, which contradicts the fact that $|\operatorname{Mi} L(\vec{\pi})| = |B(\vec{\pi})| = n$ by (6.4) and (6.6).

Finally, let $\vec{x} = \langle x_1, \ldots, x_k \rangle \in L(\vec{\pi})$. For $i \in \{1, \ldots, k\}$, let $\vec{y}^{(i)} \in L(\vec{\pi})$ be the smallest vector whose *i*-th component is x_i . Clearly, $\vec{y}^{(i)} \in C_i(\vec{\pi}), \vec{y}^{(i)} \leq \vec{x}$, and $\vec{y}^{(1)} \vee \cdots \vee \vec{y}^{(k)} = \vec{x}$. Hence Ji $(L(\vec{\pi})) \subseteq C_1(\vec{\pi}) \cup \cdots \cup C_k(\vec{\pi})$.

Lemma 6.3 (Roof Lemma). If $k \in \mathbb{N}$, $\mathbf{IL} = \mathbf{I}\langle L; C_1, \ldots, C_k \rangle$ belongs to $\mathrm{CDF}(n,k), x_1, \ldots, x_k \in L, x_1 \vee \cdots \vee x_k = 1$, and $\mathfrak{p}_i \in \mathrm{PrInt}(\uparrow x_i)$ for $i \in \{1, \ldots, k\}$, then $\mathfrak{p}_1, \ldots, \mathfrak{p}_k$ cannot belong to the same trajectory of L.

Proof. We can assume that $1 \notin \{x_1, \ldots, x_k\}$ since otherwise $PrInt(\uparrow x_i) = \emptyset$ for some $i \in \{1, \ldots, k\}$, and the statement trivially holds. Therefore, we can also assume that k > 1.

First, we deal with k = 2. For the sake of contradiction, suppose $\mathfrak{p}_1 = [u_1, v_1]$ and $\mathfrak{p}_2 = [u_2, v_2]$ belong to the same trajectory T. For $i \in \{1, 2\}$, pick a maximal chain X_i containing x_i such that $\mathfrak{p}_i \in \operatorname{PrInt}(X_i)$. Let $M = [X_1 \cup X_2]_{\vee}$, the join-subsemilattice generated by $X_1 \cup X_2$. We know from G. Czédli and E. T. Schmidt [10, Lemma 2.4] that M is a cover-preserving $\{0, 1\}$ -join-subsemilattice of L, and it is a slim semimodular lattice. Its length, n, is the same as that of L. We obtain from Lemma 3.1 that $T \cap \operatorname{PrInt}(M)$ is a trajectory of M. Thus \mathfrak{p}_1 and \mathfrak{p}_2 belong to the same trajectory of M. By [10, Lemma 2.2], M is a planar lattice, and it has a planar diagram whose left boundary chain is X_1 . The trajectories of planar semimodular lattices join-generated by two chains are well-understood. By [10, Lemma 2.9], there is an interval $\mathfrak{p}_3 = [u_3, v_3]$ such that both \mathfrak{p}_1 and \mathfrak{p}_2 are up-perspective to \mathfrak{p}_3 . This means that $u_i = v_i \wedge u_3$ and $v_3 = v_i \vee u_3$, for $i \in \{1, 2\}$. Hence $1 = x_1 \vee x_2 \le u_1 \vee u_2 \le u_3 < v_3 \le 1$, which is a contradiction that proves the statement for k = 2.

Next, to proceed by induction, assume that k > 2 and the lemma holds for smaller values. To obtain a contradiction, suppose that $\mathfrak{p}_1, \ldots, \mathfrak{p}_k$ belong to the same trajectory T of L. Let $y = x_2 \lor \cdots \lor x_k$. We can assume $y \neq 1$ since otherwise x_1 can be omitted and the induction hypothesis applies. Pick a maximal chain U of L such that $y \in U$. By Lemma 3.3, there is a unique $\mathfrak{q} \in T \cap \operatorname{PrInt}(U)$. We assert that $\mathfrak{q} \in \operatorname{PrInt}(\uparrow y)$. Suppose not, and let $R = T \cap \operatorname{PrInt}(\downarrow y)$. It is nonempty by Lemma 3.3. Hence, it is a trajectory of $\downarrow y$ by Lemma 3.1. Thus the induction hypothesis, together with Lemma 3.3, yields an $i \in \{2, \ldots, k\}$ and a prime interval \mathfrak{r}_i such that $\mathfrak{r}_i \in \operatorname{PrInt}(\downarrow x_i) \cap R \subseteq \operatorname{PrInt}(\downarrow x_i) \cap T$. Since we can clearly take a maximal chain V of L through x_i such that $\mathfrak{p}_i, \mathfrak{r}_i \in \operatorname{PrInt}(V)$, we obtain a contradiction by Lemma 3.3. Thus we conclude $\mathfrak{q} \in \operatorname{PrInt}(\uparrow y)$. Since $\mathfrak{p}_1 \in T \cap \operatorname{PrInt}(\uparrow x_1)$ and $\mathfrak{q} \in T \cap \operatorname{PrInt}(\uparrow y)$, the equality $x_1 \lor y = 1$ contradicts the case k = 2.

Let $\mathfrak{L} = \langle L; C_1, \ldots, C_k \rangle$ such that $I\mathfrak{L} \in CDF(n, k)$. We denote the elements of C_i as follows:

$$C_i = \{0 = c_0^{(i)} \prec c_1^{(i)} \prec \dots \prec c_n^{(i)} = 1\}.$$

A vector $\vec{x} \in \{0, \dots, n\}^k$ is called \mathcal{L} -maximal if either $\vec{x} = \langle n, \dots, n \rangle$, or

$$c_{x_1}^{(1)} \lor \dots \lor c_{x_k}^{(k)} < c_{x_1}^{(1)} \lor \dots \lor c_{x_{i-1}}^{(i-1)} \lor c_{x_i+1}^{(i)} \lor c_{x_{i+1}}^{(i+1)} \lor \dots \lor c_{x_k}^{(k)}$$

holds for all $i \in \{1, \ldots, k\}$.

Lemma 6.4. Let $\mathbf{I}\mathfrak{L} = \mathbf{I}\langle L; C_1, \ldots, C_k \rangle \in \text{CDF}(n, k)$. Let $\vec{\pi}$ denote its ξ -image, and let $\vec{x} \in \{0, \ldots, n\}^k$. Then \vec{x} is \mathfrak{L} -maximal iff it is an eligible $\vec{\pi}$ -tuple.

Proof. By (6.1), we can assume that $\vec{x} \in \{0, \ldots, n-1\}^k$. To prove the "only if" part, let $\vec{x} = \langle x_1, \ldots, x_k \rangle$ be an \mathfrak{L} -maximal vector, and let $i \neq j \in \{1, \ldots, k\}$. By \mathfrak{L} -maximality,

$$\begin{split} c_{x_i}^{(i)} &\lor c_{x_j}^{(j)} \lor \bigvee \big\{ c_{x_t}^{(t)} : t \in \{1, \dots, k\} \setminus \{i, j\} \big\} \\ &< c_{x_i+1}^{(i)} \lor c_{x_j}^{(j)} \lor \bigvee \big\{ c_{x_t}^{(t)} : t \in \{1, \dots, k\} \setminus \{i, j\} \big\}. \end{split}$$

Hence $c_{x_i}^{(i)} \vee c_{x_j}^{(j)} < c_{x_i+1}^{(i)} \vee c_{x_j}^{(j)}$. Thus, by Lemma 4.2, $\pi_{C_i C_j}(x_i + 1) > x_j$. Therefore, \vec{x} is an eligible $\vec{\pi}$ -tuple.

To prove the converse implication by contradiction, suppose that $\vec{x} = \langle x_1, \ldots, x_k \rangle$ is an eligible $\vec{\pi}$ -tuple but it is not \mathfrak{L} -maximal. Let, say, the k-th component of \vec{x} violate \mathfrak{L} -maximality, and let

$$u = c_{x_1}^{(1)} \vee \dots \vee c_{x_{k-1}}^{(k-1)}.$$
(6.7)

We have $u \vee c_{x_k}^{(k)} = u \vee c_{x_k+1}^{(k)}$. For $i \in \{1, \ldots, k-1\}$, extend $(C_i \cap \downarrow c_{x_i}^{(i)}) \cup \{u\}$ to a maximal chain U_i of L. We denote the trajectory of L that contains $[c_{x_k}^{(k)}, c_{x_k+1}^{(k)}]$ by T. Since $u \vee c_{x_k}^{(k)} = u \vee c_{x_k+1}^{(k)}$, Lemma 4.2 yields $\pi_{C_k U_i}(x_k+1) \leq h(u)$, where h is the height function. Hence, by Definition 4.1, there is a $\mathfrak{p}_i \in \operatorname{PrInt}(U_i) \cap T$ such that \mathfrak{p}_i is below u, that is,

$$\mathfrak{p}_i \in \operatorname{PrInt}(U_i \cap \downarrow u) \cap T. \tag{6.8}$$

On the other hand, the eligibility of \vec{x} gives $\pi_{C_kC_i}(x_k+1) \ge x_i+1$. Hence, again by Definition 4.1, C_i contains a prime interval of T above $c_{x_i}^{(i)}$. Therefore,

by Lemma 3.3, C_i does not contain any prime interval of T below $c_{x_i}^{(i)}$. But $C_i \cap \downarrow c_{x_i}^{(i)} = U_i \cap \downarrow c_{x_i}^{(i)}$, and we conclude that $\operatorname{PrInt}(U_i \cap \downarrow c_{x_i}^{(i)}) \cap T = \varnothing$. Combining this with (6.8), we obtain that $T \cap \operatorname{PrInt}(\downarrow u)$, which is a trajectory of $\downarrow u$ by Lemma 3.1, contains a prime interval in $\uparrow c_{x_i}^{(i)}$, for $i \in \{1, \ldots, k-1\}$. This, together with (6.7), contradicts Lemma 6.3.

The following lemma generalizes G. Czédli and E. T. Schmidt [11, Lemma 2.3]. We use the notation preceding Lemma 6.4. The set of suborbital vectors is still denoted by $B(\vec{\pi})$, as above (6.4). For $u \in L$, the *foot* of u is the following vector in $\{0, \ldots, n\}^k$:

$$\vec{f}(u) = \langle f_1(u), \dots, f_k(u) \rangle = \left\langle \max\{j : c_j^{(1)} \le u\}, \dots, \max\{j : c_j^{(k)} \le u\} \right\rangle.$$

Lemma 6.5. If $\mathbf{I}\mathfrak{L} = \mathbf{I}\langle L; C_1, \ldots, C_k \rangle \in \mathrm{CDF}(n, k)$, then $\{\vec{f}(u) : u \in \mathrm{Mi} L\} \subseteq B(\vec{\pi})$, where $\vec{\pi} = \xi(\mathbf{I}\mathfrak{L})$. If $\mathbf{I}\mathfrak{L} \in \mathrm{JD}(n, k)$, then even $\{\vec{f}(u) : u \in \mathrm{Mi} L\} = B(\vec{\pi})$ holds.

Proof. For k = 2, the lemma is only a reformulation of [11, Lemma 2.3]. Namely, in this case, one takes the trajectory through $[u, u^*]$; it contains a unique prime interval $[a_0, a_1]$ on the left boundary chain, and $f_1(u) = h(a_0)$; and analogously for $f_2(u)$.

Hence, we assume k > 2. Let $u \in \operatorname{Mi} L$. We are going to prove

$$\pi_{12}(f_1(u) + 1) = f_2(u) + 1. \tag{6.9}$$

Let $K = [C_1 \cup C_2]_{\vee}$; it is a slim, semimodular, cover-preserving lattice and also a $\{0, 1\}$ -join-subsemilattice of L by G. Czédli and E. T. Schmidt [10, Lemma 2.4]. Let u_0 be the largest element of $K \cap \downarrow u$.

First, if $u_0 \in Mi K$, in particular if $u = u_0$, then (6.9) follows from [11, Lemma 2.3], Lemma 3.1, and the argument detailed in the first paragraph of the present proof.

Second, for the sake of contradiction, suppose that

 u_0 is distinct from u and u_0 is meet-reducible in K.

Pick two distinct covers, a_0 and b_0 of u_0 in K. Let $u_0 \prec u_1 \prec \cdots \prec u_t = u$ be a maximal chain in the interval $[u_0, u]$. The unique cover of u in L is denoted by u^* . By semimodularity, $u \lor a_0$ and $u \lor b_0$ cover u. Hence, $u \lor a_0 = u \lor b_0 = u^*$, which implies $\{a_0, b_0\} \subseteq \downarrow u^*$. Since u_1 is not in K, the elements a_0, b_0, u_1 are three distinct atoms in the filter $\uparrow u_0$. If they are not independent, then we can select two of them that are independent in the sense of G. Grätzer [16, Theorem 380], and we easily obtain that a_0, b_0, u_1 generate a cover-preserving diamond, which is a contradiction. Hence they are independent, and they generate a cover-preserving boolean sublattice, briefly, a *cube*. Define $a_1 = a_0 \lor u_1$ and $b_1 = b_0 \lor u_1$; they belong to $\downarrow u^*$ since so are a_0, b_0, u_1 . Then the cube we have just obtained is $\{u_0, a_0, b_0, a_0 \lor b_0, u_1, a_1, b_1, a_1 \lor b_1\}$, and it is in $\downarrow u^*$. Since $u_1 \prec a_1$ and $u_1 \prec b_1, u_1 \neq u \in \operatorname{Mi} L$.

Now, we repeat the procedure within $[u_2, u^*]$ instead of $[u_1, u^*]$. If we had, say, $u_2 = a_1$, then $a_0 < a_1 = u_2 \le u$ and $a_0 \in K$ would contradict the definition of u_0 . Hence a_1, b_1, u_2 are distinct covers of u_1 . As before, they generate a cube, which is $\{u_1, a_1, b_1, u_2, a_1 \lor b_1, a_2 = a_1 \lor u_2, b_2 = b_1 \lor u_2, a_2 \lor b_2\}$. Since $u_2 \prec a_2$ and $u_2 \prec b_2, u_2 \ne u \in \operatorname{Mi} L$.

And so on. After t steps, we obtain $u = u_t \notin \text{Mi } L$, a contradiction. This proves (6.9). We obtain $\pi_{ij}(f_i(u) + 1) = f_j(u) + 1$ similarly, and we conclude $\vec{f}(u) \in B(\vec{\pi})$. This proves the first part of the lemma.

For $u \in \operatorname{Mi} L$, we have $u = c_{f_1(u)}^{(1)} \vee \cdots \vee c_{f_k(u)}^{(k)}$ since $\operatorname{Ji} L \subseteq C_1 \cup \cdots \cup C_k$. Hence, $\vec{f}(u)$ determines u, the map $\operatorname{Mi} L \to {\{\vec{f}(u) : u \in \operatorname{Mi} L\}}$, defined by $u \mapsto \vec{f}(u)$, is a bijection, and $|\{\vec{f}(u) : u \in \operatorname{Mi} L\}| = |\operatorname{Mi} L|$. Thus the second part of the lemma follows from the first part, Proposition 2.1(v), and (6.4). \Box

Proof of Proposition 6.1. Clearly, $JD(n,k) \subseteq CDF(n,k)$. Hence we obtain from Lemma 6.2 that η is a map from S_n^{k-1} to CDF(n,k).

Let $\mathbf{I}\mathfrak{L} = \mathbf{I}\langle L; C_1, \ldots, C_k \rangle \in \mathrm{CDF}(n, k)$. Denote $\xi(\mathbf{I}\mathfrak{L})$ by $\vec{\pi}$. This makes sense since, clearly, every \mathfrak{L}' isomorphic to \mathfrak{L} gives the same $\vec{\pi}$. It is obvious by Section 4 that $\vec{\pi} \in S_n^{k-1}$. That is, ξ is a map from $\mathrm{CDF}(n, k)$ to S_n^{k-1} .

Now, for $\mathbf{I}\mathfrak{L} \in \mathrm{CDF}(n,k)$ and $\pi = \xi(\mathbf{I}\mathfrak{L})$ above, we use the notation introduced before Lemma 6.4. Since $L(\vec{\pi})$ coincides with the set of \mathfrak{L} -maximal vectors by Lemma 6.4, the map $\mu: L \to L(\vec{\pi})$, defined by $u \mapsto \vec{f}(u)$, is an order-isomorphism. Thus μ is a lattice isomorphism. To deal with $\mu(C_i)$, let $x \in C_i$. For any $y \in L$, if $f_i(y) = h(x) = f_i(x)$, then $x \leq y$ and, hence, $\mu(x) = \vec{f}(x) \leq \vec{f}(y) = \mu(y)$. This shows that $\mu(x)$ is initial in its *i*-th component. Thus $\mu(C_i) \subseteq C_i(\vec{\pi})$, for $i \in \{1, \ldots, k\}$. This implies $\mu(C_i) = C_i(\vec{\pi})$ since $|C_i(\vec{\pi})| = \mathrm{length}(L(\vec{\pi})) = \mathrm{length}\,C_i$. Hence, $\eta(\vec{\pi}) = \mathrm{I}\mathfrak{L}$. This shows that

$$\eta \circ \xi$$
 is the identity map on $\text{CDF}(n, k)$. (6.10)

Now, we are in the position to prove CDF(n, k) = JD(n, k). The inclusion $\text{JD}(n, k) \subseteq \text{CDF}(n, k)$ is trivial. Conversely, let $\mathbf{I}\mathfrak{L}$ belong to CDF(n, k), and denote $\xi(\mathbf{I}\mathfrak{L})$ by $\vec{\pi}$. By (6.10), $\mathbf{I}\mathfrak{L} = \eta(\vec{\pi})$. Hence Lemma 6.2 yields $\mathbf{I}\mathfrak{L} \in \text{JD}(n, k)$. This proves CDF(n, k) = JD(n, k).

Next, let $\vec{\pi} \in S_n^{k-1}$, and consider $\eta(\vec{\pi}) = \mathbf{I}\mathfrak{L}(\vec{\pi}) = \mathbf{I}\langle L(\vec{\pi}); C_1(\vec{\pi}), \ldots, C_k(\vec{\pi}) \rangle$. Let $\vec{x} \in L(\vec{\pi})$. Since $\eta(\vec{\pi}) \in \mathrm{JD}(n,k)$ by Lemma 6.2, the length of $C_1(\vec{\pi})$ is n. This, together with the fact that no two distinct vectors in $C_1(\vec{\pi})$ have the same first component, implies that each $t \in \{0, \ldots, n\}$ is the first component of exactly one vector in $C_1(\vec{\pi})$. Thus there is a unique $\vec{y} \in C_1(\vec{\pi})$ such that $y_1 = x_1$. Since \vec{y} is initial in its first component, it is the largest vector in $C_1(\vec{\pi}) \cap \downarrow \vec{x}$. Clearly, the height of \vec{y} is $y_1 = x_1$. Hence, using the notation given before Lemma 6.5, $f_1(\vec{x}) = x_1$, and similarly for other indices. Therefore, $\vec{f}(\vec{x}) = \vec{x}$ holds for all $\vec{x} \in L(\vec{\pi})$. Applying this observation to $B(\vec{\pi}) = \{$ suborbital vectors with respect to $\vec{\pi} \}$, we conclude $\{\vec{f}(u) : u \in B(\vec{\pi})\} = B(\vec{\pi})$. Hence, by (6.6),

$$\{f(u) : u \in \operatorname{Mi}(L(\vec{\pi}))\} = B(\vec{\pi}).$$
 (6.11)

Algebra univers.

On the other hand, Lemma 6.5, applied to $\mathbf{L}\mathfrak{L}(\vec{\pi}) = \eta(\vec{\pi})$, yields the equality $\{\vec{f}(u) : u \in \operatorname{Mi}(L(\vec{\pi}))\} = B(\xi(\eta(\vec{\pi})))$. Combining this equality with (6.11), we obtain $B(\xi(\eta(\vec{\pi}))) = B(\vec{\pi})$. This means that $\vec{\pi}$ and $\xi(\eta(\vec{\pi}))$ have exactly the same suborbital vectors. Hence, they have the same orbits. Since they are determined by their orbits, we conclude $\xi(\eta(\vec{\pi})) = \vec{\pi}$. Thus $\xi \circ \eta$ is the identity map on S_n^{k-1} .

7. Coordinatizing antimatroids and convex geometries

The concept of antimatroids is due to R. E. Jamison-Waldner [20], who was the first to use the term "antimatroid". At the same time, an equivalent complementary concept was introduced by P. H. Edelman [13] under the name "anti-exchange closures". There are several ways to define antimatroids, see D. Armstrong [4, Lemma 2.1]; here we accept the following one. The set of all subsets of a set E is denoted by P(E).

Definition 7.1. A pair $\langle E, \mathfrak{F} \rangle$ is an *antimatroid* if it satisfies the following properties:

- (i) E is a finite set, and $\emptyset \neq \mathfrak{F} \subseteq P(E)$;
- (ii) \mathfrak{F} is a *feasible set*, that is, for each nonempty $A \in \mathfrak{F}$, there exists an $x \in A$ such that $A \setminus \{x\} \in \mathfrak{F}$;
- (iii) \mathfrak{F} is closed under taking unions;
- (iv) $E = \bigcup \{A : A \in \mathfrak{F} \}.$

Many authors do not stipulate (iv), that is, they allow "dummy points". For technical reasons (see Proposition 7.3), we always require (iv).

Antimatroids were generalized to more general systems called greedoid by B. Korte and L. Lovász [22] and [23]. A closure operator on a set E is an extensive, monotone, and idempotent map $\Phi: P(E) \to P(E)$. That is, $X \subseteq \Phi(X) = \Phi(\Phi(X)) \subseteq \Phi(Y)$, for all $X \subseteq Y \in P(E)$. A closure system on E is a nonempty subset of P(E) that is closed under taking arbitrary intersections. In particular, a closure system on E always contains the empty intersection, E. There is a well-known bijective correspondence between closure operators and closure systems; see S. Burris and H. P. Sankappanavar [7, I.§5]. The closure system corresponding to a closure operator Φ consists of the closed sets, that is, of the sets $X \in P(E)$ satisfying $\Phi(X) = X$. The closure operator corresponding to a closure system \mathfrak{C} is the map $P(E) \to P(E)$, defined by $X \to \bigcap \{Y \in \mathfrak{C} : X \subseteq Y\}$. Now, we define a concept closely related to antimatroids, see P.H. Edelman [13] and K. Adaricheva, V. A. Gorbunov, and V. I. Tumanov [3].

Definition 7.2. A pair $\langle E, \Phi \rangle$ is a *convex geometry*, also called *anti-exchange* system, if it satisfies the following properties:

(i) E is a finite set, and $\Phi: P(E) \to P(E)$ is a closure operator.

16

(ii) If Φ(A) = A ∈ P(E), x, y ∈ E, x ∉ A, y ∉ A, x ≠ y, and x ∈ Φ(A ∪ {y}), then y ∉ Φ(A ∪ {x}). (This is the so-called *anti-exchange property*.)
(iii) Φ(Ø) = Ø.

For a closures system \mathfrak{G} on E with corresponding closure operator Φ , $\langle E, \mathfrak{G} \rangle$ is a *convex geometry* if so is $\langle E, \Phi \rangle$ in the above sense. In what follows, the notations $\langle E, \Phi \rangle$ and $\langle E, \mathfrak{G} \rangle$ can be used interchangeably for the same mathematical object. The members of \mathfrak{G} are called *closed sets*.

It follows easily from (ii) that if $\langle E, \mathfrak{G} \rangle$ is a convex geometry, then

for each $B \in \mathfrak{G} \setminus \{E\}$, there is an $x \in E \setminus B$ such that $B \cup \{x\} \in \mathfrak{G}$. (7.1)

The following statement is taken from the book B. Korte, L. Lovász, and L. Schrader [21, Theorem III.1.3], see also (7.1) together with D. Armstrong [4, Lemma 2.5].

Proposition 7.3. Let E be a finite set, and let $\emptyset \neq \mathfrak{F} \subseteq P(E)$. Then $\mathfrak{A} = \langle E, \mathfrak{F} \rangle$ is an antimatroid iff $\mathfrak{A}^{\delta} = \langle E, \{E \setminus X : X \in \mathfrak{F}\} \rangle$ is a convex geometry.

Part of the following proposition was proved by P. H. Edelman [13, Theorem 3.3], see also D. Armstrong [4, Theorem 2.8]. The rest can be extracted from K. Adaricheva, V. A. Gorbunov, and V. I. Tumanov [3, proof of Theorem 1.9]. Since this extraction is not so obvious, we will give some details for the reader's convenience. Lattices whose duals are join-distributive were called "join-semidistributive and lower semimodular" in [3]; here we return to the original terminology of P. H. Edelman [13], and call them *meet-distributive*.

Lemma 7.4. If L is a meet-distributive lattice and $\mathfrak{M} = \langle E, \mathfrak{G} \rangle$ is a convex geometry, then the following three statements hold.

- (i) $\langle \mathfrak{G}, \subseteq \rangle$ is a meet-distributive lattice; it is denoted by LMd(\mathfrak{M}).
- (ii) $\langle \text{Ji } L, \{ \text{Ji } L \cap \downarrow x : x \in L \} \rangle$ is a convex geometry; it is denoted by Geom(L).
- (iii) $\operatorname{LMd}(\operatorname{Geom}(L)) \cong L$ and $\operatorname{Geom}(\operatorname{LMd}(\mathfrak{M})) \cong \mathfrak{M}$.

Proof. (i) was proved by P. H. Edelman [13, Theorem 3.3], see the first paragraph in his proof. The proof of K. Adaricheva, V. A. Gorbunov, and V. I. Tumanov [3, Theorem 1.9] contains the statement that Geom(L) is a convex geometry. The sixteenth line in the proof of [3, Theorem 1.9] explicitly says $\text{LMd}(\text{Geom}(L)) \cong L$.

Now, we are left with the proof of $\operatorname{Geom}(\operatorname{LMd}(\mathfrak{M})) \cong \mathfrak{M}$. So let \mathfrak{M} be a convex geometry. Its closed sets, the members of \mathfrak{G} , are exactly the elements of $\operatorname{LMd}(\mathfrak{M})$. We assert that

for every
$$A \in \mathfrak{G}$$
, A is 1-generated iff $A \in \text{Ji}(\text{LMd}(\mathfrak{M}))$. (7.2)

To show this, let $A = \{a_1, \ldots, a_t\} \in \mathfrak{G}$. Assume that $A \in \text{Ji}(\text{LMd}(\mathfrak{M}))$. The closure operator corresponding to \mathfrak{G} is denoted by Φ . Since $A = \Phi(\{a_1\}) \lor \cdots \lor \Phi(\{a_t\})$ holds in LMd(\mathfrak{M}), the join-irreducibility of A yields $A = \Phi(\{a_i\})$ for some $i \in \{1, \ldots, t\}$. This means that A is 1-generated. We prove the converse implication by way of contradiction. Suppose that A is 1-generated

but $A \notin \text{Ji}(\text{LMd}(\mathfrak{M}))$. We have $A = \Phi(X) \lor \Phi(Y)$ for some $X, Y \subset A$ such that $A \neq \Phi(X)$ and $A \neq \Phi(Y)$. Clearly, we can pick elements $b_1, \ldots, b_s \in X \cup Y$ such that $A = \Phi(\{b_1\}) \lor \cdots \lor \Phi(\{b_s\})$, this join is irredundant (that is, no joinand can be omitted), and $s \ge 2$. Since A is 1-generated, we can also pick a $c \in A$ such that $A = \Phi(\{c\})$. Since the join we consider is irredundant, $c \notin \Phi(\{b_1, \ldots, b_{s-1}\}), b_s \notin \Phi(\{b_1, \ldots, b_{s-1}\})$, and $c \neq b_s$. So we have

 $c \in \Phi(\Phi(\{b_1, \dots, b_{s-1}\} \cup \{b_s\}), \quad \{c, b_s\} \cap \Phi(\{b_1, \dots, b_{s-1}\}) = \emptyset, \quad \text{and} \ c \neq b_s.$ Thus the anti-exchange property yields that $b_s \notin \Phi(\Phi(\{b_1, \dots, b_{s-1}\} \cup \{c\}))$.

This contradicts $b_s \in A = \Phi(\{c\}) \subseteq \Phi(\Phi(\{b_1, \dots, b_{s-1}\} \cup \{c\}))$, proving (7.2). Next, if we had $x, y \in E$ such that $\Phi(x) = \Phi(y)$ but $x \neq y$, then $y \in C$

 $\Phi(\{y\}) = \Phi(\{x\}) = \Phi(\emptyset \cup \{x\})$ together with the analogous $x \in \Phi(\emptyset \cup \{y\})$ would contradict the anti-exchange property by Definition 7.2(iii). Thus $\Phi(x) = \Phi(y)$ implies x = y. Hence, by (7.2), for each $A \in \text{Ji}(\text{LMd}(\mathfrak{M}))$, there is a unique $e_A \in E$ such that $A = \Phi(\{e_A\})$. Since $\Phi(\{e\}) \in \text{Ji}(\text{LMd}(\mathfrak{M}))$ also holds for all $e \in E$ by (7.2), we have a bijection ψ : Ji (LMd $(\mathfrak{M})) \to E$, defined by $A \mapsto e_A$. Its inverse is denoted by η ; it is defined by $\eta(e) = \Phi(\{e\})$. We assert that ψ is an isomorphism from Geom(LMd (\mathfrak{M})) to \mathfrak{M} . To prove this, let X be a closed set of Geom(LMd (\mathfrak{M})). This means that X is of the form $X = \{A \in \text{Ji}(\text{LMd}(\mathfrak{M})) : A \subseteq B\}$ for some $B \in \text{LMd}(\mathfrak{M})) = \mathfrak{G}$. Hence,

$$\psi(X)=\psi\bigl(\{\eta(e):\eta(e)\subseteq B\}\bigr)=\psi\bigl(\{\eta(e):e\in B\}\bigr)=\{\psi(\eta(e)):e\in B\}=B.$$

That is, ψ maps the closed sets X of Geom(LMd(\mathfrak{M})) to the closed sets of \mathfrak{M} . Since each $B \in \text{LMd}(\mathfrak{M})$) = \mathfrak{G} determines a closed set $X = \{A \in \text{Ji}(\text{LMd}(\mathfrak{M})) : A \subseteq B\}$ of Geom(LMd(\mathfrak{M})), the calculation above also shows that all $B \in \mathfrak{G}$ are ψ -images of closed sets of Geom(LMd(\mathfrak{M})). Thus ψ is an isomorphism.

Combining Proposition 7.3 with the dual of Lemma 7.4, we easily obtain the following statement. It asserts that join-distributive lattices and antimatroids are essentially the same mathematical objects.

Corollary 7.5. If L is a join-distributive lattice and $\mathfrak{A} = \langle E, \mathfrak{F} \rangle$ is an antimatroid, then the following three statements hold.

- (i) $\langle \mathfrak{F}, \subseteq \rangle$ is a join-distributive lattice; it is denoted by $\mathrm{LJd}(\mathfrak{A})$.
- (ii) $\langle \operatorname{Mi} L, \{\operatorname{Mi} L \setminus \uparrow x : x \in L\} \rangle$ is an antimatroid; it is denoted by $\operatorname{Amat}(L)$,
- (iii) $\mathrm{LJd}(\mathrm{Amat}(L)) \cong L$ and $\mathrm{Amat}(\mathrm{LJd}(\mathfrak{A})) \cong \mathfrak{A}$.

Now, Corollary 7.5 allows us to translate Theorem 5.2 to a coordinatization of antimatroids, while the coordinatization of convex geometries is reduced to that of antimatroids by Proposition 7.3. A brief translation is exemplified by the following corollary; the full translation and the case of convex geometries are omitted.

Let $\vec{\pi} \in S_n^{k-1}$. As before, the set of suborbital vectors and that of $\vec{\pi}$ eligible tuples are denoted by $B(\vec{\pi})$ and $L(\vec{\pi})$, respectively; see Definition 5.1, and see also (6.3), (6.4), and (6.6). For $\vec{x} \in L(\vec{\pi})$, let $U(\vec{x})$ denote the set

 $\{\vec{y} \in B(\vec{\pi}) : \vec{x} \leq \vec{y}\}$. The convex dimension of an antimatroid \mathfrak{A} is the width of Ji (LJd(\mathfrak{A})).

Corollary 7.6. For each $\vec{\pi} \in S_n^{k-1}$, $\mathfrak{A}(\vec{\pi}) = \langle B(\vec{\pi}), \{U(\vec{x}) : x \in L(\vec{\pi})\} \rangle$ is an antimatroid with convex dimension at most k. Conversely, for each antimatroid \mathfrak{B} of convex dimension k on an n-element set, there exists a $\vec{\pi} \in S_n^{k-1}$ such that \mathfrak{B} is isomorphic to $\mathfrak{A}(\vec{\pi})$.

Acknowledgment. The author is indebted to Anna Romanowska for calling his attention to [1] and [19], and to Kira Adaricheva for her comments that led to [2]. The referee's careful work is also appreciated.

References

- Abels, H.: The geometry of the chamber system of a semimodular lattice. Order 8, 143–158 (1991)
- [2] Adaricheva, K., Czédli, G.: Notes on the description of join-distributive lattices by permutations. http://arxiv.org/abs/1210.3376
- [3] Adaricheva, K., Gorbunov, V.A., Tumanov, V.I.: Join-semidistributive lattices and convex geometries. Advances in Math. 173, 1–49 (2003)
- [4] Armstrong, D.: The Sorting Order on a Coxeter Group. Journal of Combinatorial Theory Series A 116, 1285–1305 (2009)
- [5] Avann, S.P.: Application of the join-irreducible excess function to semimodular lattices. Math. Ann. 142, 345–354 (1961)
- [6] Avann, S.P.: Increases in the join-excess function in a lattice. Math. Ann. 154, 420–426 (1964)
- [7] Burris, S., Sankappanavar, H.P.: A Course in Universal Algebra. Graduate Texts in Mathematics, vol. 78. Springer, New YorkBerlin (1981). The Millennium Edition: http://www.math.uwaterloo.ca/~snburris/htdocs/ualg.html
- [8] Czédli, G., Ozsvárt, L., Udvari, B.: How many ways can two composition series intersect?. Discrete Mathematics 312, 3523–3536 (2012)
- Czédli, G., Schmidt, E.T.: Some results on semimodular lattices. In: Proceedings of the AAA79 Conference in Olomouc, 2010. Contributions to General Algebra, vol. 19, pp. 45–56. Johannes Hein verlag, Klagenfurt (2010)
- [10] Czédli, G., Schmidt, E.T.: The Jordan-Hölder theorem with uniqueness for groups and semimodular lattices. Algebra Universalis 66, 69–79 (2011)
- [11] Czédli, G., Schmidt, E.T.: Composition series in groups and the structure of slim semimodular lattices. Acta Sci.Math. (Szeged) (in press); see also at http://arxiv.org/abs/1208.4749
- [12] Dilworth, R.P.: Lattices with unique irreducible decompositions. Ann. of Math. (2)
 41, 771–777 (1940)
- [13] Edelman, P.H.: Meet-distributive lattices and the anti-exchange closure. Algebra Universalis 10, 290–299 (1980)
- [14] Edelman, P.H.: Abstract convexity and meet-distributive lattices. In: Proc. Conf. Combinatorics and ordered sets (Arcata, Calif., 1985). Contemp. Math. vol. 57, pp. 127–150. Amer. Math. Soc., Providence, RI (1986)
- [15] Edelman, P.H., Jamison, R.E.: The theory of convex geometries. Geom. Dedicata 19, 247-271 (1985)
- [16] Grätzer, G.: Lattice Theory: Foundation. Birkhäuser, Basel (2011)
- [17] Grätzer, G., Knapp, E.: Notes on planar semimodular lattices I. Construction. Acta Sci. Math. (Szeged) 73, 445–462 (2007)
- [18] Grätzer, G., Nation, J B.: A new look at the Jordan-Hölder theorem for semimodular lattices. Algebra Universalis 66, 69–79 (2011)
- [19] Herscovici, D.S.: Semimodular lattices and semibuildings. Journal of Algebraic Combinatorics 7, 39–51 (1998)

- [20] Jamison-Waldner, R.E.: Copoints in antimatroids. Proceedings of the 11th Southeast. Conference, Boca Raton/Florida 1980. Combinatorics, graph theory and computing, vol. II, Congr. Numerantium 29, pp. 535–544 (1980)
- [21] Korte, B., Lovász, L., Schrader, R.: Greedoids, Algorithms and Combinatorics 4. Springer (1991)
- [22] Korte, B., Lovász, L.: Mathematical structures underlying greedy algorithms. Proc. int. FCT-Conf., Szeged, Hungary 1981 Fundamentals of computation theory. Lect. Notes Comput. Sci. vol. 117, pp. 205–209 (1981)
- [23] Korte, B., Lovász, L.: Structural properties of greedoids. Combinatorica 3, 359–374 (1983)
- [24] Monjardet, B.: A use for frequently rediscovering a concept. Order 1, 415–417 (1985)
- [25] Stanley, R.P.: Supersolvable lattices. Algebra Universalis 2, 197–217 (1972)
- [26] Stern, M.: Semimodular Lattices. Theory and Applications. Encyclopedia of Mathematics and its Applications vol. 73. Cambridge University Press (1999)

Gábor Czédli

University of Szeged, Bolyai Institute, Szeged, Aradi vértanúk tere 1, HUNGARY 6720 *e-mail*: czedli@math.u-szeged.hu

URL: http://www.math.u-szeged.hu/~czedli/

20