2-DISTRIBUTIVITY AND LATTICES OF SUBLATTICES CLOSED UNDER TAKING RELATIVE COMPLEMENTS

GÁBOR CZÉDLI

Dedicated to Professor Béla Csákány on his ninetieth birthday

ABSTRACT. For a modular lattice L of finite length, we prove that the distributivity of L is a sufficient condition while its 2-distributivity is a necessary condition that those sublattices of L that are closed under taking relative complements form a ranked lattice of finite length.

Compared to the previous (January 6, 2022) arXiv version, now Theorem 1, Lemma 1, and Corollary 1 are stronger since they do not assume the modularity of L.

For elements u, x, v of a lattice L, let $rc_L(u, x, v) := \{y \in L : x \land y = u \text{ and } x \lor y = v\}$. A sublattice S of L is closed with respect to taking relative complements or, shortly saying, it is an RC-closed sublattice if for all $u, x, v \in S$, we have that $rc_L(u, x, v) \subseteq S$. The set consisting of the empty set and the RC-closed sublattices of L will be denoted by RCSub(L). For any lattice L, the poset $RCSub(L) = (RCSub(L), \subseteq)$ is an algebraic lattice. Following Huhn [7, 8], a lattice L is n-distributive if for all $x, y_0, \ldots, y_n \in L, x \land \bigvee \{y_i : 0 \le i \le n\} = \bigvee_{j=0}^n (x \land \bigvee \{y_i : 0 \le i \le n \text{ and } i \ne j\})$. The length len(L) of a lattice L is the supremum of the lengths of its finite chains; for a finite chain C, len(C) = |C| - 1. A ranked lattice is a lattice in which any two maximal chains of every principal ideal are of the same finite length. Our aim is to prove the following four statements.

Theorem 1. If L is a lattice of finite length, then len(RCSub(L)) = 1 + len(L).

Theorem 2. If L is modular lattice such that $\operatorname{RCSub}(L)$ is a ranked lattice of finite length, then L is a 2-distributive lattice of finite length.

Theorem 3. For every finite distributive lattice L, RCSub(L) is a ranked lattice.

For a lattice L and $X \subseteq L$, $\operatorname{rcg}_L(X)$ stands for the least RC-closed sublattice of L that includes X as a subset; "g" in the acronym comes from "generated".

Lemma 1. If L is a lattice of finite length, Y is an RC-closed sublattice of L, X is a sublattice of Y, and len(X) = len(Y), then $Y = rcg_L(X)$.

Date: January 15, 2022. Check the author's website (later MathSciNet or Zentralblatt) for updates and related papers.

²⁰¹⁰ Mathematics Subject Classification. 06C99, 06D99.

Key words and phrases. Relative complement, distributive lattice, modular lattice, 2distributive lattice, *n*-distributivity, sublattice, ranked lattice.

This research was supported by the National Research, Development and Innovation Fund of Hungary under funding scheme K 134851.

For the particular case where L is distributive, Lemma 1 could be extracted from Section 10 of Grätzer [3]. Letting Y := L, the lemma trivially implies the following.

Corollary 1. If X is a sublattice of a lattice L of finite length such that len(X) = len(L), then L is RC-generated by X.

Proof of Lemma 1. We give a proof by contradiction. With $S := \operatorname{rcg}_L(X)$, suppose that $S \neq Y$. Since $X \subseteq S \subset Y$, $\operatorname{len}(Y) = \operatorname{len}(X) \leq \operatorname{len}(S) \leq \operatorname{len}(Y)$ gives that $\operatorname{len}(S) = \operatorname{len}(Y)$. This allows us to pick a maximal chain K in S such that $\operatorname{len}(K) = \operatorname{len}(Y)$. We have that $0_K = 0_S = 0_Y$, $1_K = 1_S = 1_Y$, and both K and Sare cover-preserving sublattices of Y, that is, for any $x, y \in S$, $x \prec_S y \iff x \prec_Y y$, and similarly for K. For $a \in L$, the principal ideal and the principal filter generated by a are denoted by $\downarrow_L a := \{x \in L : x \leq a\}$ and $\uparrow_L a := \{x \in L : x \geq a\}$, respectively. We write $\downarrow a$ and $\uparrow a$ if L is understood. If $u \leq v$ in Y, then the length of the interval [u, v], understood in Y, will be denoted by $\operatorname{len}_Y([u, v])$. For, say, $u \leq v \in K$, the notation $\operatorname{len}_K([u, v])$ is analogously defined. For $x \in Y$, we let

$$x_{\bullet} := \bigvee (K \cap \downarrow_Y x) \quad \text{and} \quad x^{\bullet} := \bigwedge (K \cap \uparrow_Y x).$$

Since K is finite and so it is a complete sublattice, $x_{\bullet}, x^{\bullet} \in K$ for every $x \in Y$. Choose an element $b \in Y \setminus S$ such that $d := \operatorname{len}_K([b_{\bullet}, b^{\bullet}]) = \operatorname{len}_Y([b_{\bullet}, b^{\bullet}])$ is minimal. Since $K \ni b_{\bullet} \leq b \leq b^{\bullet} \in K$ but $b \notin K$, $d \geq 2$. Hence, we can pick an element $q \in K$ such that $b_{\bullet} < q < b^{\bullet}$. Then $z := b \land q \in Y$. If z was outside S, then $K \ni b_{\bullet} \leq z \leq q \in K$ would give that $b_{\bullet} \leq z_{\bullet} \leq z^{\bullet} \leq q < b^{\bullet}$ and $\operatorname{len}_K([z_{\bullet}, z^{\bullet}]) < \operatorname{len}_K([b_{\bullet}, b^{\bullet}]) = d$ would be a contradiction. Thus, $b \land q = z \in S$. Dually, $b \lor q \in S$ and, of course, $q \in K \subseteq S$. Thus, $b \in \operatorname{rc}_L(b \land q, q, b \lor q) \in S$, contradicting that $b \in Y \setminus S$.

In the following three proofs, we can assume that n := len(L) is at least 2. As always in Lattice Theory, " \subset " is the conjunction of " \subseteq " and " \neq ".

Proof of Theorem 1. Let $0 = c_0 \prec c_1 \cdots \prec c_n = 1$ be a chain in L. Let $X_{-1} := \emptyset$ and, for $i \in \{0, \ldots, n\}$, $X_i := \downarrow c_i$. Since all these X_i belong to RCSub(L) and $X_{-1} \subset X_0 \subset X_1 \subset \cdots \subset X_n = L$, $\operatorname{len}(\operatorname{RCSub}(L)) \ge 1 + n = 1 + \operatorname{len}(L)$. Next, let $Y_{-1} \subset Y_0 \subset \cdots \subset Y_k = L$ be an arbitrary chain in RCSub(L). Clearly, $\operatorname{len}(Y_{i-1}) \le \operatorname{len}(Y_i)$ for all $i \in \{0, 1, \ldots, k\}$. We claim that $\operatorname{len}(Y_{i-1}) < \operatorname{len}(Y_i)$ for all meaningful i. Suppose the contrary. Then $Y_{i-1} \subset Y_i$ and $\operatorname{len}(Y_{i-1}) = \operatorname{len}(Y_i)$ for some i. With reference to Lemma 1 at "=*", $Y_{i-1} = \operatorname{rcg}_L(Y_{i-1}) = * Y_i$, which is a contradiction proving that $\operatorname{len}(Y_{i-1}) < \operatorname{len}(Y_i)$ for all meaningful i. Therefore, since $-1 \le \operatorname{len}(\emptyset) \le \operatorname{len}(Y_{-1})$, we obtain that $k \le 1 + n$. Thus, $\operatorname{len}(\operatorname{RCSub}(L)) \le$ $1 + n = 1 + \operatorname{len}(L)$, completing the proof. \Box

Proof of Theorem 2. Following von Neumann [11] and going also after Herrmann [5], a system $F = (a_i, c_{i,j} : i, j \in \{1, \ldots, n\}, i \neq j)$ of elements of a modular lattice L is a non-trivial normalized (von Neumann) n-frame or, briefly, a von Neumann n-frame if, with the notation $0_F := \bigwedge_{i=1}^n a_i$ and $1_F := \bigvee_{i=1}^n a_i$, we have that $0_F \neq 1_F$, $a_j \land \bigvee_{t\neq j} a_t = 0_F = a_i \land c_{i,j}, c_{i,j} = c_{j,i}, a_i \lor c_{i,j} = a_i \lor a_j$, and $c_{i,k} = (a_i \lor a_k) \land (c_{i,j} \lor c_{j,k})$ for all $\{i, j, k\} \subseteq \{1, \ldots, n\}$ with $|\{i, j, k\}| = 3$. Here $2 \le n \in \mathbb{N}^+ := \{1, 2, 3, \ldots\}$. We know from Huhn [10, Proposition 1.2] that

for $n \in \mathbb{N}^+$, a modular lattice is *n*-distributive if and only if it does not contain a von Neumann (n + 1)-frame. (1) if $F = (a_i, c_{i,j} : i, j \in \{1, \dots, n\}, i \neq j)$ is a von Neumann *n*frame in a modular lattice L and $a'_1 \in L$ such that $0_F < a'_1 < a_1$, then a'_1 belongs to a von Neumann *n*-frame $F = (a'_i, c'_{i,j} : i, j \in \{1, \dots, n\}, i \neq j)$ such that $0_F = 0_{F'} < a'_i < a_i$ for $i \in \{1, \dots, n\}$. (2)

We say that a projective space is *irreducible* (AKA nondegenerate) if each of its lines contains at least three points. It is known (and easy to see) that each point of an irreducible projective plane lies on at least three lines. By Huhn [9, Thm. 1.1],

for $n \in \mathbb{N}^+$, a modular algebraic lattice is *n*-distributive if and only if none of its sublattices is isomorphic to the subspace (3) lattice of an irreducible *n*-dimensional projective geometry.

Assume that L is a modular lattice such that $\operatorname{RCSub}(L)$ is a ranked lattice of finite length. Whenever $\cdots < c_i < c_{i+1} < \ldots$ is a (finite or infinite) chain in L, then $\cdots < \operatorname{rcg}_L(\downarrow c_i) < \operatorname{rcg}_L(\downarrow c_{i+1}) < \ldots$ is a chain in $\operatorname{RCSub}(L)$. Thus, L is of finite length. For the sake of contradiction, suppose that L is not 2-distributive. By (1), Lcontains a von Neumann 3-frame $F = (a_i, c_{i,j} : i \neq j, i, j \in \{1, \dots, 3\})$. Applying (2), repeatedly if necessary, we can assume that $0_F \prec a_i$, for $i \in \{1, 2, 3\}$. The definition of a 3-frame implies that $\{a_1, a_2, a_3\}$ is an independent set of atoms in the filter $\uparrow 0_F$. Hence, $1_F = a_1 \lor a_2 \lor a_3$ is of height 3 in $\uparrow 0_F$, that is, $\operatorname{len}_L([0_F, 1_F]) = 3$. By (1), the interval $I := [0_F, 1_F]$ is not 2-distributive. Using (3), we obtain that the subspace lattice S of a projective plane G is a sublattice of I. Since len(I) =3 = len(S), S is a cover-preserving sublattice of I and L, $0_S = 0_F = 0_I$, and $1_s = 1_I$. Let At(S) and Coat(S) denote the set of atoms and that of coatoms of S, respectively. The are disjoint sets, $S = \{0_S, 1_S\} \cup \operatorname{At}(S) \cup \operatorname{Coat}(S), \operatorname{At}(S) =$ $\{a \in I : 0_I \prec a, a \in S\}$, and dually. A trivial geometric argument shows that for $\forall a \in \operatorname{At}(S) \text{ and } \forall b \in \operatorname{Coat}(S), 1_S \stackrel{*}{=} \bigvee (\operatorname{At}(S) \setminus \{a\}) \text{ and } 0_S \stackrel{**}{=} \bigwedge (\operatorname{Coat}(S) \setminus \{b\}).$ Define $Z_{-1} := \emptyset$, $Z_0 := \{0_S\}$, $Z_1 := \{0_S, 1_S\}$, and $Z_2 := \operatorname{rcg}_L(S)$. We claim that

$$Z_{-1}, Z_0, Z_1, Z_2 = I \in \operatorname{RCSub}(L) \text{ and } Z_{-1} \prec Z_0 \prec Z_1 \prec Z_2 \text{ in } \operatorname{RCSub}(L).$$
(4)

By Lemma 1, $I = \operatorname{rcg}_I(S)$. Since $\operatorname{rcg}_I(S) \subseteq \operatorname{rcg}_L(S)$, we obtain that $I \subseteq \operatorname{rcg}_L(S) = Z_2$. Since every interval is RC-closed, $S \subseteq I \in \operatorname{RCSub}(L)$ yields that $Z_2 = \operatorname{rcg}_L(S) \subseteq \operatorname{rcg}_L(I) = I$. Thus, $Z_2 = I$. Trivially, $Z_{-1} \prec Z_0 \prec Z_1$ and $Z_1 \subset Z_2$. To verify that $Z_1 \prec Z_2 = I$, assume that $Z_1 \subset X \subseteq I$ for some $X \in \operatorname{RCSub}(L)$. Pick an element $u \in X \setminus Z_1$. Then $0_F = 0_S < u < 1_S$. Since len(S) = 3, either u is of height 1, or it is of height 2. First, assume that u is of height 2, that is, $u \prec 1_S$ in I (and in L). If we had that $|\operatorname{At}(S) \setminus |u| \leq 1$, then " $\stackrel{*}{=}$ " would give that $1_S = \bigvee(\operatorname{At}(S) \cap |u|) \leq u$, contradicting that $u < 1_S$. Hence, $|\operatorname{At}(S) \setminus |u| \geq 2$, and we can pick two distinct elements, v_1 and v_2 , from $\operatorname{At}(S) \setminus |u|$. For $s \in \{1, 2\}$, using that $v_s \not\leq u$, $u, v_s \in [0_S, 1_S]$, $0_S \prec v_s$, and $u \prec 1_S$, we obtain that $v_s \in \operatorname{rcL}(0_S, u, 1_S)$. Hence, $v_s \in X$. By modularity (in fact, by semimodularity), $v_1 \prec v_1 \lor v_2 \in X$. Since len(I) = 3, $0_S = 0_I \prec v_1 \prec v_1 \lor v_2 < 1_S = 1_I$. This chain, being in X, shows that $3 \leq \operatorname{len}(X)$. On the other hand, $\operatorname{len}(X) = I = Z_2$. Thus, $Z_1 \prec Z_2$, completing the proof of (4).

By (4), $Z_{-1} \prec \cdots \prec Z_2$ extends to a maximal chain $\vec{Z} : Z_{-1} \prec \cdots \prec Z_k$ of $\operatorname{RCSub}(L)$. Like in the proof of Theorem 1, Lemma 1 applies and we obtain that

G. CZÉDLI

 $-1 \leq \operatorname{len}(Z_{-1}) < \operatorname{len}(Z_0) < \cdots < \operatorname{len}(Z_k)$. In fact, we know even that $\operatorname{len}(Z_1) + 1 = 2 < 3 = \operatorname{len}(Z_2)$. Thus, the maximal chain \vec{Z} consists of at most $1 + \operatorname{len}(L)$ elements, whence $\operatorname{len}(\vec{Z}) \leq \operatorname{len}(L)$. But $\operatorname{RCSub}(L)$ is a ranked lattice of finite length, whereby $\operatorname{len}(\operatorname{RCSub}(L)) = \operatorname{len}(\vec{Z}) \leq \operatorname{len}(L)$, contradicting Theorem 1.

Proof of Theorem 3. Let L be a finite distributive lattice. Since it is a ranked lattice of finite length, it suffices to show that for any $U, V \in \mathrm{RCSub}(L)$,

$$U \prec V \text{ in RCSub}(L) \iff (U \subset V \text{ and } \operatorname{len}(V) = \operatorname{len}(U) + 1).$$
 (5)

To prove the \Rightarrow direction, assume that $U \prec V$. Clearly, $U \subset V$ and $\operatorname{len}(U) \leq \operatorname{len}(V)$. If we had that $\operatorname{len}(U) = \operatorname{len}(V)$, then Lemma 1 would give that $U = \operatorname{rcg}_L(U) = V$, a contradiction. Hence, $\operatorname{len}(U) < \operatorname{len}(V)$. We are going to show by way of contradiction that $\operatorname{len}(V) = \operatorname{len}(U) + 1$. Suppose the contrary; then $k := \operatorname{len}(U) \leq \operatorname{len}(V) - 2$. Take a maximal chain C_0 in U. Since $\operatorname{len}(C_0) = k$ and $\operatorname{len}(V) \geq k + 2$, we can extend C_0 to a chain C of V such that $\operatorname{len}(C) = k + 1$. Let $W := \operatorname{rcg}_L(C)$. It follows from Lemma 1 that $U = \operatorname{rcg}_L(C_0)$. Hence, $U = \operatorname{rcg}_L(C_0) \subseteq \operatorname{rcg}_L(C) = W$. Since $\operatorname{len}(W) \geq \operatorname{len}(C) > \operatorname{len}(U)$, $W \neq U$. Hence, $U \subset W$. The inclusion $C \subseteq V$ gives that $W = \operatorname{rcg}_L(C) \subseteq \operatorname{rcg}_L(V) = V$. Combining $U \subset W \subseteq V$ and $U \prec V$, we obtain that W = V.

Next, we write C in the form $C = \{c_0 < c_1 < \cdots < c_{k+1}\}$. (Note that "<" here need not mean covering in L.) By Birkhoff [1], we can fix a finite Boolean lattice D such that L is a sublattice of D. We define the elements $b_i \in D$ for $i \in \{1, \ldots, k+1\}$ by the rule $b_i \in \operatorname{rc}_D(c_0, c_{i-1}, c_i)$. Since D is a Boolean lattice, b_i exists and it is uniquely determined. We claim that, for $i = 2, 3, \ldots, k+1$,

$$c_0 \notin \{b_1, \dots, b_i\}, \ c_{i-1} = b_1 \lor \dots \lor b_{i-1}, \ \text{and} \ (b_1 \lor \dots \lor b_{i-1}) \land b_i = c_0.$$
 (6)

We show this by induction on *i*. Clearly, $b_1 = c_1 \neq c_0$. Since $b_2 \in \operatorname{rc}_D(c_0, c_1, c_2)$, we have that $b_2 \neq c_0$ since otherwise $c_2 = c_1 \lor b_2 = c_1$ would be a contradiction. Also, $b_2 \in \operatorname{rc}_D(c_0, c_1, c_2)$ gives that $b_1 \land b_2 = c_1 \land b_2 = c_0$. Hence, (6) holds for i = 2. Assume that $2 \leq i < k + 1$ and (6) holds for this *i*. As before, $b_{i+1} \neq c_0$ since otherwise $b_{i+1} \in \operatorname{rc}_D(c_0, c_i, c_{i+1})$ would lead to $c_{i+1} = b_{i+1} \lor c_i = c_i$, which is a contradiction. Using the definition of b_i and the induction hypothesis, we have that $c_i = c_{i-1} \lor b_i = b_1 \lor \cdots \lor b_{i-1} \lor b_i$, that is, the second equality of (6) holds for i + 1. Using this equality and the definition of b_{i+1} , we have that $(b_1 \lor \cdots \lor b_i) \land b_{i+1} = c_i \land b_{i+1} = c_0$. This completes the induction step and proves that (6) holds for $i = 2, 3, \ldots, k + 1$.

By, say, Grätzer [4, Theorem 360] and (6), $\{b_1, b_2, \ldots, b_{k+1}\}$ is a (k+1)-element independent set in the filter $\uparrow_D c_0$. Thus, this set generates a 2^{k+1} -element Boolean sublattice E. Using that any element in an interval of a distributive lattice has at most one relative complement with respect to the interval in question and E as a Boolean lattice is closed with respect to taking relative complements, we obtain that E is RC-closed. It is clear from (6) and $b_{k+1} \in \operatorname{rc}_D(c_0, c_k, c_{k+1})$ that $C \subseteq E$. Hence, $\operatorname{rcg}_D(C) \subseteq \operatorname{rcg}_D(E) = E$. Now let F be a maximal chain in W. Since $W = \operatorname{rcg}_L(C) \subseteq \operatorname{rcg}_D(C) \subseteq E$, we have that F is a chain in E. But $\operatorname{len}(E) = k+1$, implying that $\operatorname{len}(F) \leq k+1$. So $\operatorname{len}(W) = \operatorname{len}(F) \leq k+1$. On the other hand, $k+1 = \operatorname{len}(C) \leq \operatorname{len}(W)$. Thus, $\operatorname{len}(W) = k+1$, which is a contradiction since W = V and $\operatorname{len}(V) = k+2$. This proves the \Rightarrow direction of (5).

To prove the \Leftarrow direction, assume that $U \subset V$ and $\operatorname{len}(V) = \operatorname{len}(U) + 1$. Assume also that $H \in \operatorname{RCSub}(L)$ such that $U \subseteq H \subseteq V$. Clearly, $\operatorname{len}(U) \leq \operatorname{len}(H) \leq$ len(V), whence len(H) is len(U) or len(V). If len(H) = len(U), then Lemma 1 gives that $U = \operatorname{rcg}_L(U) = H$. Similarly, if len(H) = len(V), then the same lemma gives that $H = \operatorname{rcg}_L(H) = V$. Therefore, $U \prec V$, completing the proof.

Some facts are easy to observe; see Czédli [2] for details. For a lattice L, RCSub(L) is Boolean \iff RCSub(L) is semimodular \iff RCSub(L) is lower semimodular. Let B_n stand for the 2^n -element Boolean lattice. Our tools lead to a (bit complicated) formula for $|\text{RCSub}(B_n)|$, which allows us to compute (in 0.05 seconds with computer algebra) that, say, $|\text{RCSub}(B_{57})|$ is the 59-digit number

 $46\,669\,606\,977\,325\,325\,544\,440\,157\,525\,187\,321\,911\,338\,002\,625\,473\,546\,541\,556$

where B_{57} is the Boolean lattice of length 57. It took 24 seconds to obtain that $|\text{RCSub}(B_{999})| \approx 0.566759343075881648 \cdot 10^{1930}$.

References

- Birkhoff, G.: Rings of sets. Duke Math. J. 3, 443–454 (1937)
- [2] Czédli, G.: A property of lattices of sublattices closed under taking relative complements and its connection to 2-distributivity. In preparation
- [3] Grätzer, G.: Lattice Theory. First Concepts and Distributive Lattices
- [4] Grätzer, G.: Lattice Theory: Foundation. Birkhäuser Verlag, Basel (2011)
- [5] Herrmann, C.: On the arithmetic of projective coordinate systems. Transactions of the American Math. Soc. 284, 759–785 (1984)
- [6] Herrmann, C., Huhn, A. P.: Lattices of normal subgroups generated by frames. Lattice theory (Proc. Colloq., Szeged, 1974), pp. 97–136. Colloq. Math. Soc. János Bolyai, Vol. 14, North-Holland, Amsterdam, 1976
- [7] Huhn, A. P.: Schwach distributive Verbände. (German). Acta Fac. Rerum Natur. Univ. Comenianae Math. (Bratislava), 51–56 (1971)
- [8] Huhn, A. P.: Schwach distributive Verbände. I. (German). Acta Sci. Math. (Szeged) 33, 297–305 (1972)
- Huhn, A. P.: Two notes on n-distributive lattices. Lattice theory (Proc. Colloq., Szeged, 1974), pp. 137–147. Colloq. Math. Soc. János Bolyai, Vol. 14, North-Holland, Amsterdam, 1976.
- [10] Huhn, A. P.: n-distributivity and some questions of the equational theory of lattices. Contributions to universal algebra (Colloq., József Attila Univ., Szeged, 1975), pp. 167–178. Colloq. Math. Soc. János Bolyai, Vol. 17, North-Holland, Amsterdam, 1977.
- [11] Von Neumann, J.: Continuous Geometry. Princeton Univ. Press, Princeton, N. J., 1960
 E-mail address: czedli@math.u-szeged.hu Address: University of Szeged, Hungary URL: http://www.math.u-szeged.hu/~czedli/